



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 10:12 PM GMT

PDB ID : 1SL1
Title : Binary 5' complex of T7 DNA polymerase with a DNA primer/template containing a cis-syn thymine dimer on the template
Authors : Li, Y.; Dutta, S.; Doublié, S.; Bdour, H.M.; Taylor, J.S.; Ellenberger, T.
Deposited on : 2004-03-05
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	22	<div> <div>14%</div> <div>23%</div> <div>5%</div> <div>59%</div> </div>
2	T	25	<div> <div>16%</div> <div>24%</div> <div>60%</div> </div>
3	A	698	<div> <div>9%</div> <div>71%</div> <div>24%</div> <div>• •</div> </div>
4	B	108	<div> <div>3%</div> <div>67%</div> <div>29%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MG	A	4003	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6654 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*CP*GP*AP*AP*AP*AP*CP*GP*AP*CP*GP*GP*CP*CP*AP*GP*TP*GP*CP*CP*TP*(2DA))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	P	9	Total	C	N	O	P	0	0	0
			172	82	28	53	9			

- Molecule 2 is a DNA chain called 5'-D(*CP*CP*C*(TTD)P*AP*GP*GP*CP*AP*CP*TP*GP*GP*CP*CP*GP*TP*CP*GP*TP*TP*TP*TP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	10	Total	C	N	O	P	0	0	0
			223	107	42	64	10			

- Molecule 3 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	678	Total	C	N	O	S	0	0	0
			5243	3340	907	972	24			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	LYS	DELETION	UNP P00581
A	?	-	ARG	DELETION	UNP P00581
A	?	-	PHE	DELETION	UNP P00581
A	?	-	GLY	DELETION	UNP P00581
A	?	-	SER	DELETION	UNP P00581
A	?	-	HIS	DELETION	UNP P00581

- Molecule 4 is a protein called Thioredoxin 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	105	Total	C	N	O	S	0	0	0
			776	503	122	148	3			

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Mg 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	213	Total 213	O 213	0	0
6	B	20	Total 20	O 20	0	0
6	T	6	Total 6	O 6	0	0



● Molecule 1: 5'-D(*CP*GP*AP*AP*AP*AP*CP*GP*AP*CP*GP*GP*CP*CP*AP*GP*TP*GP*CP*CP*TP*(2DA))-3'



DC	DC	DC	T5	A6	G7	G8	C9	A10	G14	DC	DC	DG	DT	DC	DG	DT	DT	DT	DT	DC	DG
----	----	----	----	----	----	----	----	-----	-----	----	----	----	----	----	----	----	----	----	----	----	----

A	GLU	A513	A420	P301	M167	M1
	GLN	A514	A421	LYS		
	GLN	L515	P422	ASN	Y170	F18
	VAL	P516		LYS	M171	
	K587	T517	A425	ALA	V172	T28
	W588	R518		GLN	Q173	A29
	K589		R429	ARG		E30
		A521	A430	GLU	V176	
	I593	I525	T431	GLY	V177	R35
	K594	Y526	H432	ARG		P36
D	D597	G527	A433	E311	L181	E49
	G598	F528		P312		
	R599	L529	M436	C313	L185	
	A609	Y530	Q439	D316	P193	G53
	L610	G531	L440		P194	G54
	M611	A532	P441	V321		V57
	T612	G533	O442	A322	T204	
	L613	D534		G323	T205	H62
	A619	K536	P446	A324	E209	Q76
	L620	T537	O447	V326	S210	R79
I	I621	G538	O448		L211	E30
	G622	Q539	R452	P336		
		I540			A220	R85
	I626	A543	G456	R339	A221	
	L638	G544	A457		T222	
		K545	P458	K344	L223	I89
	W642	E546	H459		L224	T91
	D645	E547	K467	Q347	F232	L92
		G548		E348	P233	V93
		R549	Q471	P353	F234	L94
M	M649	E550		T354	D235	
	A650	L551	L474	K355		R111
	M651	K552		V356	T239	
	F652	F555	E480	T357	G270	Y130
	H653	L556	L481	D358		R131
	D654	E557	M488	K359	D273	E134
	G655	D654	A489	G360	M73	
	I656	N558	A489	A361	F274	E138
		T559	R490	P362	G275	
	G659	P560	F491	V363	R276	D141
R		A561	D492	P277	R278	
		I562	D365			R145
	E663	R566	Y496		P284	E149
	P664		E499	V368		
	L665		I500	L369	Q288	Q150
	Q667	I569	L501	G371	L289	G151
		L573		V372	K290	
	E671	V574	D604		T291	D156
		E575	I505			
	F686	SER	E586	V389	V294	M158
L	L689	SER	T507	K394	G295	
		GLN	R508			
	M696	TRP	I509	R404	L297	
	G697	VAL	O510	F298	T298	M164
		ALA	T511	V410	K299	E165
		GLY	A512		V200	E162



● Molecule 4: Thioredoxin 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	104.38 Å 214.78 Å 52.18 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20 36.90 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.4 (50.00-2.20) 98.6 (36.90-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.03 (at 2.20 Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.242 , 0.262 0.240 , 0.258	Depositor DCC
R_{free} test set	2814 reflections (4.72%)	DCC
Wilson B-factor (Å ²)	36.1	Xtriage
Anisotropy	0.256	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 59617 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6654	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 2DA, TTD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	P	0.50	0/180	0.79	0/275
2	T	0.38	0/211	0.76	0/324
3	A	0.34	0/5374	0.57	0/7301
4	B	0.30	0/791	0.55	0/1079
All	All	0.34	0/6556	0.58	0/8979

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	P	0	1
2	T	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	P	19	DC	Sidechain
2	T	8	DG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	172	0	97	11	0
2	T	223	0	122	10	0
3	A	5243	0	4982	141	0
4	B	776	0	761	22	0
5	A	1	0	0	0	0
6	A	213	0	0	4	0
6	B	20	0	0	0	0
6	T	6	0	0	0	0
All	All	6654	0	5962	174	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 174 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:131:ARG:HH11	3:A:131:ARG:HB3	1.20	1.07
4:B:19:ALA:HB3	4:B:23:ILE:HD11	1.43	0.99
3:A:642:TRP:HH2	3:A:649:MET:HE2	1.29	0.95
3:A:18:PHE:H	3:A:76:GLN:HE22	1.23	0.86
3:A:501:LEU:HD21	3:A:689:LEU:HB3	1.67	0.77

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	672/698 (96%)	641 (95%)	22 (3%)	9 (1%)	15	11
4	B	103/108 (95%)	99 (96%)	3 (3%)	1 (1%)	19	16
All	All	775/806 (96%)	740 (96%)	25 (3%)	10 (1%)	15	11

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	156	ASP
3	A	531	GLY
3	A	360	GLY
3	A	514	GLU
3	A	544	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	520/579 (90%)	504 (97%)	16 (3%)	47	59
4	B	78/87 (90%)	75 (96%)	3 (4%)	40	49
All	All	598/666 (90%)	579 (97%)	19 (3%)	46	57

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	481	LEU
3	A	518	ARG
3	A	686	PHE
3	A	429	ARG
4	B	10	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
3	A	343	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	B	98	GLN
3	A	347	GLN
3	A	164	ASN
3	A	510	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	2DA	P	22	1	6,10,23	0.45	0	7,12,34	0.40	0
2	TTD	T	5	2	39,40,46	3.88	9 (23%)	54,67,77	3.66	23 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	2DA	P	22	1	-	0/3/12/19	0/1/1/3
2	TTD	T	5	2	-	0/19/101/110	0/3/6/6

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	T	5	TTD	C5T-C6T	-19.19	1.32	1.55
2	T	5	TTD	C5-C6	-10.10	1.43	1.55
2	T	5	TTD	C2-N3	-3.77	1.31	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	T	5	TTD	O4R-C1R	-2.07	1.37	1.42
2	T	5	TTD	C5M-C5T	2.80	1.59	1.53

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	5	TTD	C5A-C5-C4	-7.70	97.41	108.41
2	T	5	TTD	O5R-PB-O5P	-7.54	80.36	109.62
2	T	5	TTD	O4'-C1'-C2'	-6.60	97.12	106.86
2	T	5	TTD	O4P-PB-O5R	-4.95	83.51	108.46
2	T	5	TTD	C5T-C5-C6	-4.77	82.33	88.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	T	5	TTD	5	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	P	8/22 (36%)	-0.10	0 100 100	42, 49, 60, 68	0
2	T	9/25 (36%)	0.42	0 100 100	35, 49, 64, 68	0
3	A	678/698 (97%)	0.39	62 (9%) 11 11	19, 35, 64, 71	0
4	B	105/108 (97%)	0.19	3 (2%) 55 54	31, 45, 58, 60	0
All	All	800/853 (93%)	0.36	65 (8%) 15 14	19, 37, 63, 71	0

The worst 5 of 65 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	544	GLY	7.8
3	A	548	GLY	6.4
3	A	513	ALA	6.2
3	A	298	PHE	5.2
3	A	313	CYS	5.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	2DA	P	22	10/21	0.88	0.14	-	59,62,64,64	0
2	TTD	T	5	35/41	0.67	0.31	-	71,73,79,80	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	MG	A	4003	1/1	0.55	0.34	8.97	47,47,47,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.