



Full wwPDB NMR Structure Validation Report ⓘ

Apr 27, 2016 – 05:04 AM BST

PDB ID : 2SPZ
Title : STAPHYLOCOCCAL PROTEIN A, Z-DOMAIN, NMR, 10 STRUCTURES
Authors : Montelione, G.T.; Tashiro, M.; Tejero, R.; Lyons, B.A.
Deposited on : 1998-07-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 96%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	58	 50% 31% • 16%

2 Ensemble composition and analysis

This entry contains 10 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:8-A:56 (49)	0.45	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	2, 3, 5, 6, 7
2	1, 8, 10
Single-model clusters	4; 9

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 926 atoms, of which 457 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called IMMUNOGLOBULIN G BINDING PROTEIN A.

Mol	Chain	Residues	Atoms					Trace
1	A	58	Total	C	H	N	O	0
			926	291	457	83	95	

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	VAL	ALA	ENGINEERED	UNP P38507
A	29	ALA	GLY	ENGINEERED	UNP P38507

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: IMMUNOGLOBULIN G BINDING PROTEIN A




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: IMMUNOGLOBULIN G BINDING PROTEIN A




4.2.2 Score per residue for model 2

- Molecule 1: IMMUNOGLOBULIN G BINDING PROTEIN A




4.2.3 Score per residue for model 3

- Molecule 1: IMMUNOGLOBULIN G BINDING PROTEIN A




4.2.4 Score per residue for model 4

- Molecule 1: IMMUNOGLOBULIN G BINDING PROTEIN A




4.2.5 Score per residue for model 5

- Molecule 1: IMMUNOGLOBULIN G BINDING PROTEIN A




4.2.6 Score per residue for model 6

- Molecule 1: IMMUNOGLOBULIN G BINDING PROTEIN A




4.2.7 Score per residue for model 7

- Molecule 1: IMMUNOGLOBULIN G BINDING PROTEIN A




4.2.8 Score per residue for model 8 (medoid)

- Molecule 1: IMMUNOGLOBULIN G BINDING PROTEIN A




4.2.9 Score per residue for model 9

- Molecule 1: IMMUNOGLOBULIN G BINDING PROTEIN A



4.2.10 Score per residue for model 10

- Molecule 1: IMMUNOGLOBULIN G BINDING PROTEIN A



5 Refinement protocol and experimental data overview

The models were refined using the following method: *SIMULATED ANNEALING WITH RESTRAINED MOLECULAR DYNAMICS*.

Of the 40 calculated structures, 10 were deposited, based on the following criterion: *LOWEST CONFORMATIONAL ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CONGEN	refinement	
DIANA	structure solution	
CONGEN	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 4023
Number of chemical shift lists	1
Total number of shifts	928
Number of shifts mapped to atoms	770
Number of unparsed shifts	14
Number of shifts with mapping errors	144
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	96%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.83±0.02	0±0/398 (0.0±0.0%)	1.40±0.09	6±2/540 (1.1±0.4%)
All	All	0.83	0/3980 (0.0%)	1.40	61/5400 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.7±0.5
All	All	0	7

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	30	PHE	CB-CG-CD2	10.38	128.06	120.80	7	7
1	A	30	PHE	CB-CG-CD1	9.54	127.48	120.80	5	7
1	A	14	TYR	CB-CG-CD1	8.88	126.33	121.00	10	5
1	A	14	TYR	CB-CG-CD2	-8.87	115.68	121.00	10	4
1	A	27	ARG	NE-CZ-NH2	8.84	124.72	120.30	5	3
1	A	44	LEU	CB-CG-CD1	8.45	125.37	111.00	6	1
1	A	32	GLN	N-CA-CB	-8.16	95.92	110.60	2	8
1	A	27	ARG	NE-CZ-NH1	7.39	124.00	120.30	3	3
1	A	13	PHE	CB-CG-CD2	-7.23	115.74	120.80	9	4
1	A	9	GLN	N-CA-CB	-6.74	98.46	110.60	6	9
1	A	9	GLN	CA-CB-CG	6.29	127.23	113.40	3	1
1	A	13	PHE	CB-CG-CD1	6.06	125.04	120.80	7	3
1	A	24	GLU	N-CA-CB	-5.89	100.00	110.60	8	3
1	A	30	PHE	CA-CB-CG	-5.73	100.16	113.90	10	2
1	A	9	GLN	CB-CA-C	5.07	120.55	110.40	3	1

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	27	ARG	Sidechain	7

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	392	375	375	5±2
All	All	3920	3750	3750	49

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:19:LEU:HD13	1:A:22:LEU:HD13	0.63	1.69	5	2
1:A:21:ASN:HD22	1:A:56:ALA:HB2	0.62	1.54	3	2
1:A:24:GLU:CD	1:A:24:GLU:H	0.58	2.03	9	4
1:A:21:ASN:HD21	1:A:56:ALA:HB2	0.55	1.61	6	1
1:A:52:ASN:C	1:A:52:ASN:HD22	0.55	2.04	9	1
1:A:24:GLU:H	1:A:24:GLU:CD	0.55	2.05	1	1
1:A:37:ASP:HB3	1:A:40:GLN:CG	0.52	2.33	7	1
1:A:34:LEU:HD22	1:A:44:LEU:HD23	0.52	1.82	7	1
1:A:21:ASN:ND2	1:A:56:ALA:HB3	0.50	2.22	1	1
1:A:26:GLN:HE22	1:A:55:GLN:NE2	0.49	2.04	10	4
1:A:44:LEU:H	1:A:44:LEU:CD1	0.49	2.20	6	1
1:A:21:ASN:HD21	1:A:52:ASN:CG	0.48	2.12	4	1
1:A:26:GLN:NE2	1:A:55:GLN:HE22	0.48	2.06	3	1
1:A:12:ALA:HA	1:A:45:LEU:HD22	0.47	1.85	7	2
1:A:22:LEU:HD11	1:A:26:GLN:NE2	0.46	2.25	5	1
1:A:24:GLU:N	1:A:24:GLU:CD	0.46	2.69	1	1
1:A:16:ILE:O	1:A:27:ARG:NH1	0.46	2.49	1	1
1:A:41:SER:HA	1:A:44:LEU:HG	0.45	1.87	9	2
1:A:37:ASP:HB3	1:A:40:GLN:HG2	0.44	1.90	7	1
1:A:24:GLU:CD	1:A:24:GLU:N	0.44	2.70	10	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:22:LEU:HD11	1:A:26:GLN:HE22	0.44	1.73	5	1
1:A:21:ASN:HD22	1:A:56:ALA:CB	0.43	2.23	3	1
1:A:26:GLN:NE2	1:A:55:GLN:NE2	0.43	2.67	3	3
1:A:25:GLU:H	1:A:25:GLU:CD	0.43	2.17	2	1
1:A:25:GLU:HA	1:A:28:ASN:ND2	0.43	2.28	4	1
1:A:52:ASN:C	1:A:52:ASN:ND2	0.42	2.73	9	1
1:A:9:GLN:NE2	1:A:34:LEU:O	0.42	2.52	10	1
1:A:22:LEU:CD1	1:A:26:GLN:NE2	0.41	2.83	5	1
1:A:21:ASN:ND2	1:A:56:ALA:HB2	0.41	2.30	5	1
1:A:9:GLN:O	1:A:34:LEU:HD11	0.41	2.15	7	1
1:A:25:GLU:CD	1:A:26:GLN:N	0.41	2.73	4	1
1:A:12:ALA:CB	1:A:41:SER:OG	0.41	2.69	6	1
1:A:19:LEU:CD1	1:A:22:LEU:HD13	0.41	2.42	5	1
1:A:41:SER:HA	1:A:44:LEU:CG	0.40	2.46	9	2
1:A:27:ARG:CZ	1:A:27:ARG:HB3	0.40	2.47	9	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	49/58 (84%)	44±2 (89±4%)	4±2 (7±4%)	2±1 (4±1%)	7	34
All	All	490/580 (84%)	436 (89%)	35 (7%)	19 (4%)	7	34

All 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	38	PRO	7
1	A	24	GLU	3
1	A	56	ALA	3
1	A	20	PRO	1
1	A	37	ASP	1
1	A	9	GLN	1
1	A	40	GLN	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	23	ASN	1
1	A	19	LEU	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	42/51 (82%)	35±2 (82±5%)	7±2 (18±5%)	6	41
All	All	420/510 (82%)	346 (82%)	74 (18%)	6	41

All 23 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	8	GLU	9
1	A	52	ASN	6
1	A	35	LYS	6
1	A	11	ASN	5
1	A	50	LYS	5
1	A	24	GLU	5
1	A	32	GLN	4
1	A	16	ILE	4
1	A	15	GLU	4
1	A	44	LEU	4
1	A	49	LYS	3
1	A	25	GLU	3
1	A	34	LEU	2
1	A	38	PRO	2
1	A	27	ARG	2
1	A	55	GLN	2
1	A	20	PRO	2
1	A	40	GLN	1
1	A	23	ASN	1
1	A	26	GLN	1
1	A	22	LEU	1
1	A	9	GLN	1
1	A	43	ASN	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 96% for the well-defined parts and 96% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 4023

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	928
Number of shifts mapped to atoms	770
Number of unparsed shifts	14
Number of shifts with mapping errors	144
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

The following errors were found when reading this chemical shift list.

- Chemical shift has been reported more than once. All 14 occurrences are reported below.

Shift ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
269	A	9	GLN	N	121.6	.	1
270	A	9	GLN	H	8.54	.	1
271	A	9	GLN	C	176.5	.	1
272	A	9	GLN	CA	57.3	.	1
273	A	9	GLN	HA	3.88	.	1
274	A	9	GLN	CB	28.6	.	1
275	A	9	GLN	HB2	1.54	.	2
276	A	9	GLN	HB3	2.25	.	2
277	A	9	GLN	CG	33.2	.	1
278	A	9	GLN	HG2	2.49	.	1
279	A	9	GLN	HG3	2.49	.	1
280	A	9	GLN	CD	178.6	.	1
281	A	9	GLN	HE21	6.96	.	2
282	A	9	GLN	HE22	7.26	.	2

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Residue not found in structure. All 144 occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
A	7	ASN	HB2	2.76	-1.0	2
A	6	LEU	HA	4.25	-1.0	1
A	13	ALA	H	8.28	-1.0	1
A	6	LEU	HG	1.57	-1.0	1
A	11	ASP	HA	4.61	-1.0	1
A	6	LEU	HB2	1.97	-1.0	1
A	10	HIS	HE1	7.17	-1.0	1
A	3	ILE	CB	38.6	-1.0	1
A	5	VAL	CG2	20.7	-1.0	2
A	3	ILE	HD12	0.8	-1.0	1
A	7	ASN	HB3	2.82	-1.0	2
A	11	ASP	N	122.0	-1.0	1
A	11	ASP	CB	41.1	-1.0	1
A	3	ILE	CD1	12.6	-1.0	1
A	6	LEU	HD13	0.86	-1.0	2
A	2	ALA	HA	4.3	-1.0	1
A	11	ASP	C	174.9	-1.0	1
A	6	LEU	N	125.8	-1.0	1
A	6	LEU	HD21	0.91	-1.0	2
A	12	GLU	HB3	2.05	-1.0	2
A	12	GLU	C	175.2	-1.0	1
A	2	ALA	HB1	1.3	-1.0	1
A	10	HIS	N	119.1	-1.0	1
A	12	GLU	CG	36.1	-1.0	1
A	13	ALA	CB	18.4	-1.0	1
A	1	LYS	HB3	1.78	-1.0	2
A	12	GLU	HG2	2.27	-1.0	1
A	11	ASP	HB2	2.59	-1.0	2
A	5	VAL	C	174.2	-1.0	1
A	3	ILE	H	8.02	-1.0	1
A	2	ALA	H	8.4	-1.0	1
A	7	ASN	HD22	7.61	-1.0	2
A	5	VAL	HG23	0.86	-1.0	1
A	5	VAL	CG1	20.2	-1.0	2
A	3	ILE	HD13	0.8	-1.0	1
A	1	LYS	HA	4.35	-1.0	1
A	7	ASN	HD21	6.91	-1.0	2
A	6	LEU	HD12	0.86	-1.0	2

Continued on next page...

Continued from previous page...

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
A	6	LEU	CA	55.0	-1.0	1
A	10	HIS	C	176.4	-1.0	1
A	6	LEU	CG	26.7	-1.0	1
A	8	ALA	HB3	1.35	-1.0	1
A	6	LEU	HD22	0.91	-1.0	2
A	12	GLU	HB2	1.96	-1.0	2
A	5	VAL	HG11	0.86	-1.0	1
A	3	ILE	CG1	26.8	-1.0	1
A	10	HIS	CA	55.2	-1.0	1
A	7	ASN	CA	52.8	-1.0	1
A	1	LYS	HG3	1.44	-1.0	1
A	2	ALA	CB	19.0	-1.0	1
A	5	VAL	CB	32.0	-1.0	1
A	1	LYS	CA	56.0	-1.0	1
A	12	GLU	HG3	2.27	-1.0	1
A	11	ASP	HB3	2.67	-1.0	2
A	1	LYS	CG	24.4	-1.0	1
A	3	ILE	HG13	1.36	-1.0	2
A	6	LEU	C	174.5	-1.0	1
A	1	LYS	HD2	1.68	-1.0	1
A	8	ALA	H	8.2	-1.0	1
A	4	PHE	HA	4.68	-1.0	1
A	5	VAL	HG22	0.86	-1.0	1
A	10	HIS	HB3	3.21	-1.0	2
A	1	LYS	C	174.5	-1.0	1
A	4	PHE	HD1	7.18	-1.0	1
A	8	ALA	HB2	1.35	-1.0	1
A	6	LEU	HD23	0.91	-1.0	2
A	2	ALA	C	175.9	-1.0	1
A	5	VAL	H	7.99	-1.0	1
A	3	ILE	CG2	17.2	-1.0	1
A	7	ASN	N	119.1	-1.0	1
A	5	VAL	HA	4.04	-1.0	1
A	1	LYS	HG2	1.44	-1.0	1
A	2	ALA	CA	51.8	-1.0	1
A	4	PHE	CD2	130.9	-1.0	1
A	12	GLU	CA	56.7	-1.0	1
A	1	LYS	CB	33.0	-1.0	1
A	3	ILE	HG21	0.8	-1.0	1
A	8	ALA	N	123.7	-1.0	1
A	6	LEU	CD1	23.3	-1.0	2

Continued on next page...

Continued from previous page...

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
A	4	PHE	C	173.9	-1.0	1
A	8	ALA	CB	18.6	-1.0	1
A	3	ILE	HG12	1.1	-1.0	2
A	4	PHE	CA	57.1	-1.0	1
A	7	ASN	ND2	112.6	-1.0	1
A	1	LYS	HD3	1.68	-1.0	1
A	5	VAL	HG21	0.86	-1.0	1
A	3	ILE	CA	60.7	-1.0	1
A	3	ILE	HD11	0.8	-1.0	1
A	1	LYS	HE3	2.98	-1.0	1
A	4	PHE	H	8.22	-1.0	1
A	3	ILE	HB	1.75	-1.0	1
A	10	HIS	HB2	3.13	-1.0	2
A	13	ALA	HA	4.26	-1.0	1
A	7	ASN	H	8.42	-1.0	1
A	4	PHE	HD2	7.18	-1.0	1
A	8	ALA	HB1	1.35	-1.0	1
A	7	ASN	HA	4.66	-1.0	1
A	13	ALA	C	176.9	-1.0	1
A	13	ALA	HB3	1.35	-1.0	1
A	8	ALA	HA	4.26	-1.0	1
A	5	VAL	HB	1.97	-1.0	1
A	2	ALA	HB2	1.3	-1.0	1
A	6	LEU	H	8.23	-1.0	1
A	4	PHE	CD1	130.9	-1.0	1
A	7	ASN	C	173.7	-1.0	1
A	5	VAL	N	122.8	-1.0	1
A	3	ILE	HG22	0.8	-1.0	1
A	10	HIS	H	8.37	-1.0	1
A	4	PHE	N	124.0	-1.0	1
A	4	PHE	CB	39.5	-1.0	1
A	13	ALA	HB1	1.35	-1.0	1
A	13	ALA	HB2	1.35	-1.0	1
A	3	ILE	N	119.7	-1.0	1
A	1	LYS	HE2	2.98	-1.0	1
A	3	ILE	HA	4.12	-1.0	1
A	12	GLU	HA	4.22	-1.0	1
A	6	LEU	HD11	0.86	-1.0	2
A	7	ASN	CB	38.5	-1.0	1
A	6	LEU	CB	41.9	-1.0	1
A	2	ALA	HB3	1.3	-1.0	1

Continued on next page...

Continued from previous page...

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
A	3	ILE	C	174.4	-1.0	1
A	5	VAL	HG12	0.86	-1.0	1
A	10	HIS	CB	29.5	-1.0	1
A	5	VAL	CA	61.5	-1.0	1
A	1	LYS	CD	28.7	-1.0	1
A	3	ILE	HG23	0.8	-1.0	1
A	10	HIS	HD2	8.3	-1.0	1
A	12	GLU	H	8.52	-1.0	1
A	13	ALA	CA	52.2	-1.0	1
A	6	LEU	HB3	1.97	-1.0	1
A	13	ALA	N	123.9	-1.0	1
A	1	LYS	HB2	1.75	-1.0	2
A	11	ASP	CA	54.0	-1.0	1
A	5	VAL	HG13	0.86	-1.0	1
A	11	ASP	H	8.35	-1.0	1
A	7	ASN	CG	176.0	-1.0	1
A	8	ALA	C	176.9	-1.0	1
A	2	ALA	N	126.2	-1.0	1
A	12	GLU	N	122.0	-1.0	1
A	12	GLU	CB	29.9	-1.0	1
A	10	HIS	HA	4.67	-1.0	1
A	1	LYS	CE	41.5	-1.0	1
A	6	LEU	CD2	24.8	-1.0	2
A	8	ALA	CA	52.1	-1.0	1

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	70	0.05 ± 0.25	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	70	0.70 ± 0.11	Should be applied
$^{13}\text{C}'$	70	0.90 ± 0.09	Should be applied
^{15}N	66	0.21 ± 0.17	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 96%, i.e. 590 atoms were assigned a chemical shift out of a possible 614. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	241/241 (100%)	96/96 (100%)	98/98 (100%)	47/47 (100%)
Sidechain	321/340 (94%)	195/199 (98%)	114/123 (93%)	12/18 (67%)
Aromatic	28/33 (85%)	16/18 (89%)	12/14 (86%)	0/1 (0%)
Overall	590/614 (96%)	307/313 (98%)	224/235 (95%)	59/66 (89%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 96%, i.e. 713 atoms were assigned a chemical shift out of a possible 741. 0 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	284/284 (100%)	113/113 (100%)	116/116 (100%)	55/55 (100%)
Sidechain	392/415 (94%)	240/244 (98%)	138/148 (93%)	14/23 (61%)
Aromatic	37/42 (88%)	21/23 (91%)	16/18 (89%)	0/1 (0%)
Overall	713/741 (96%)	374/380 (98%)	270/282 (96%)	69/79 (87%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	5	PHE	CD2	122.30	137.34 – 125.84	-8.1
1	A	5	PHE	CD1	122.30	137.63 – 125.43	-7.6

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

