



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:15 PM GMT

PDB ID : 1ST0
Title : Structure of DcpS bound to m7GpppG
Authors : Gu, M.; Fabrega, C.; Liu, S.W.; Liu, H.; Kiledjian, M.; Lima, C.D.
Deposited on : 2004-03-24
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	<div> <div>3%</div> <div> <div></div> <div>70%</div> <div>15%</div> <div>•</div> <div>11%</div> </div> </div>
1	B	337	<div> <div>4%</div> <div> <div></div> <div>67%</div> <div>18%</div> <div>••</div> <div>12%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5647 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called mRNA decapping enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	297	Total	C	N	O	S	0	0	0
			2439	1554	435	447	3			
1	A	300	Total	C	N	O	S	0	0	0
			2458	1565	438	452	3			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	277	ASN	HIS	ENGINEERED	UNP Q96C86
B	277	ASN	HIS	ENGINEERED	UNP Q96C86

- Molecule 2 is YTTRIUM (III) ION (three-letter code: YT3) (formula: Y).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Y	0	0
			3	3		

- Molecule 3 is 7-METHYL-GUANOSINE-5'-TRIPHOSPHATE-5'-GUANOSINE (three-letter code: GTG) (formula: C₂₁H₃₀N₁₀O₁₈P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			52	21	10	18	3		
3	B	1	Total	C	N	O	P	0	0
			52	21	10	18	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	349	Total	O	0	0
			349	349		
4	B	294	Total	O	0	0
			294	294		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: mRNA decapping enzyme



- Molecule 1: mRNA decapping enzyme



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.45Å 94.78Å 178.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 19.83 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.1 (20.00-1.90) 99.2 (19.83-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.187 , 0.231 0.187 , 0.230	Depositor DCC
R_{free} test set	3991 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	29.1	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 79543 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5647	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: YT3, GTG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	6/2515 (0.2%)	1.13	12/3413 (0.4%)
1	B	1.05	2/2495 (0.1%)	1.07	12/3386 (0.4%)
All	All	1.09	8/5010 (0.2%)	1.10	24/6799 (0.4%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	190	VAL	CB-CG2	5.90	1.65	1.52
1	A	177	TYR	CD1-CE1	5.48	1.47	1.39
1	A	200	PHE	CE2-CZ	5.39	1.47	1.37
1	A	269	TYR	CE1-CZ	5.26	1.45	1.38
1	B	86	LYS	CE-NZ	5.21	1.62	1.49
1	A	280	PHE	CD1-CE1	5.17	1.49	1.39
1	A	217	TYR	CE2-CZ	5.09	1.45	1.38
1	B	58	ARG	CG-CD	5.04	1.64	1.51

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	231	ARG	NE-CZ-NH1	12.04	126.32	120.30
1	A	231	ARG	NE-CZ-NH2	-12.02	114.29	120.30
1	B	261	ASP	CB-CG-OD2	8.23	125.70	118.30
1	B	84	LEU	CA-CB-CG	7.88	133.43	115.30
1	B	232	ASP	CB-CG-OD2	7.48	125.03	118.30
1	A	215	ASP	CB-CG-OD2	7.23	124.81	118.30
1	B	76	ASP	CB-CG-OD2	6.92	124.53	118.30
1	A	298	LEU	CB-CG-CD2	6.91	122.75	111.00
1	B	126	ASP	CB-CG-OD2	6.71	124.33	118.30
1	A	325	ASP	CB-CG-OD2	6.66	124.29	118.30
1	A	315	ARG	NE-CZ-NH2	-6.54	117.03	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	74	ASP	CB-CG-OD1	6.39	124.05	118.30
1	B	187	ASP	CB-CG-OD2	6.26	123.93	118.30
1	A	255	ARG	NE-CZ-NH1	6.13	123.37	120.30
1	B	147	ASP	CB-CG-OD2	6.02	123.72	118.30
1	B	214	ASP	CB-CG-OD2	5.83	123.54	118.30
1	B	188	ARG	NE-CZ-NH1	5.72	123.16	120.30
1	A	156	ASP	CB-CG-OD2	5.59	123.33	118.30
1	B	315	ARG	NE-CZ-NH1	5.45	123.02	120.30
1	A	232	ASP	CB-CG-OD2	5.29	123.06	118.30
1	A	205	ASP	CB-CG-OD2	5.22	123.00	118.30
1	A	54	ARG	NE-CZ-NH1	5.21	122.90	120.30
1	B	74	ASP	CB-CG-OD2	5.19	122.97	118.30
1	B	297	LEU	CB-CG-CD1	-5.04	102.43	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2458	0	2445	44	0
1	B	2439	0	2428	46	0
2	A	3	0	0	0	1
3	A	52	0	26	1	0
3	B	52	0	26	3	0
4	A	349	0	0	12	0
4	B	294	0	0	12	0
All	All	5647	0	4925	84	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:102:PRO:HA	4:B:942:HOH:O	1.26	1.33
1:A:254:GLN:HG3	4:A:962:HOH:O	1.48	1.11
1:B:205:ASP:HB2	1:B:219:ILE:CD1	1.84	1.06
1:B:205:ASP:HB2	1:B:219:ILE:HD12	1.11	1.05
1:B:91:VAL:HG23	4:A:829:HOH:O	1.62	0.97
1:B:90:GLN:HG3	4:B:906:HOH:O	1.69	0.93
1:B:205:ASP:OD2	1:B:207:LYS:HG2	1.69	0.92
1:A:40:VAL:HG12	1:A:41:ARG:H	1.36	0.91
1:A:322:ARG:HD3	4:A:836:HOH:O	1.77	0.84
1:A:40:VAL:HG12	1:A:41:ARG:N	1.95	0.81
1:A:245:LEU:HB2	1:A:298:LEU:HD13	1.62	0.80
1:A:234:THR:H	1:A:237:HIS:HD2	1.29	0.78
1:B:205:ASP:CB	1:B:219:ILE:HD12	2.05	0.74
1:B:168:SER:O	1:B:169:GLN:HB2	1.88	0.73
1:A:40:VAL:CG1	1:A:41:ARG:H	2.04	0.71
1:B:163:LEU:O	1:B:167:GLU:HG2	1.92	0.70
1:B:74:ASP:HB3	1:B:76:ASP:H	1.55	0.70
1:A:246:HIS:HE1	4:A:845:HOH:O	1.78	0.65
1:B:51:LYS:HE2	4:B:867:HOH:O	1.97	0.65
1:A:158:TYR:CG	1:A:231:ARG:HD3	2.32	0.64
1:A:234:THR:H	1:A:237:HIS:CD2	2.14	0.63
1:B:169:GLN:OE1	1:B:169:GLN:HA	1.99	0.62
1:B:176:VAL:HG13	1:B:221:ILE:HG21	1.81	0.61
1:B:48:ARG:HD3	1:B:70:GLU:OE1	2.01	0.61
1:A:197:SER:HB3	4:A:926:HOH:O	2.00	0.61
1:A:254:GLN:HG3	4:A:1047:HOH:O	2.00	0.60
1:A:137:GLU:OE1	1:A:141:GLN:NE2	2.36	0.59
1:A:322:ARG:HG2	1:A:325:ASP:HB2	1.85	0.58
1:B:236:GLU:HG3	1:B:236:GLU:O	2.02	0.57
1:B:314:GLN:HG2	4:B:822:HOH:O	2.03	0.56
1:A:331:LEU:O	1:A:335:GLN:HG3	2.05	0.56
1:A:163:LEU:HB3	1:A:164:PRO:HD3	1.88	0.56
1:B:103:GLU:OE2	1:A:122:ARG:NH1	2.39	0.55
1:B:103:GLU:CD	1:A:122:ARG:NH1	2.60	0.55
1:A:40:VAL:CG1	1:A:41:ARG:N	2.63	0.53
1:B:272:SER:HB2	3:B:652:GTG:O3A	2.08	0.53
1:B:336:GLN:HB2	4:B:865:HOH:O	2.09	0.53
1:B:230:LEU:HD12	1:B:233:LEU:HD12	1.91	0.52
1:B:74:ASP:HB2	4:B:887:HOH:O	2.10	0.51
1:A:96:GLN:HG2	4:A:959:HOH:O	2.11	0.51
1:A:51:LYS:HE2	1:A:144:LEU:CD1	2.43	0.49
1:B:314:GLN:NE2	4:B:815:HOH:O	2.44	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:117:HIS:ND1	1:A:126:ASP:OD2	2.46	0.48
1:A:41:ARG:HG2	4:A:1001:HOH:O	2.13	0.48
1:B:204:PRO:HA	1:B:218:LEU:HD23	1.96	0.48
1:A:322:ARG:CD	4:A:836:HOH:O	2.49	0.48
1:B:94:VAL:CG1	1:A:64:LEU:HD11	2.43	0.48
1:A:207:LYS:NZ	3:A:651:GTG:O2G	2.30	0.47
1:B:99:THR:HG22	1:A:47:PHE:O	2.15	0.47
1:B:141:GLN:HG3	4:B:896:HOH:O	2.14	0.47
1:B:40:VAL:HG11	4:B:899:HOH:O	2.15	0.47
4:B:709:HOH:O	1:A:315:ARG:HD2	2.15	0.47
1:A:51:LYS:HE2	1:A:144:LEU:HD13	1.98	0.46
1:B:226:GLY:O	4:B:778:HOH:O	2.21	0.46
1:A:233:LEU:HA	1:A:237:HIS:CD2	2.50	0.46
1:B:95:ALA:O	1:B:99:THR:HG23	2.15	0.46
1:B:103:GLU:O	1:B:103:GLU:CG	2.64	0.45
1:A:158:TYR:CD1	1:A:231:ARG:HD3	2.52	0.45
1:A:231:ARG:HD2	1:A:269:TYR:OH	2.16	0.45
1:A:237:HIS:HE1	4:A:837:HOH:O	1.99	0.45
1:B:82:VAL:HG12	1:B:84:LEU:HD22	1.97	0.44
3:B:652:GTG:H2'	3:B:652:GTG:O1G	2.17	0.44
1:A:138:LYS:HE3	4:A:1029:HOH:O	2.17	0.44
1:B:94:VAL:HG11	1:A:64:LEU:HD11	1.99	0.44
1:B:57:ALA:O	1:B:60:LYS:HE3	2.17	0.44
1:B:103:GLU:HB3	4:B:753:HOH:O	2.16	0.44
1:A:336:GLN:HG2	1:A:337:SER:N	2.33	0.44
1:B:189:ILE:HA	1:B:203:ILE:HG22	1.99	0.44
1:A:106:LEU:HD11	1:A:109:SER:HB2	2.00	0.43
1:B:221:ILE:HG23	1:B:275:HIS:CD2	2.55	0.42
1:A:161:ILE:HD11	1:A:334:ALA:HB2	2.01	0.42
1:B:193:ASN:OD1	1:B:193:ASN:C	2.56	0.42
1:A:167:GLU:HB2	4:A:971:HOH:O	2.19	0.42
1:B:209:ASN:CG	1:B:211:GLN:HG3	2.40	0.42
1:B:209:ASN:OD1	1:B:211:GLN:HG3	2.20	0.41
1:B:304:ASN:OD1	1:A:315:ARG:NH2	2.32	0.41
1:A:146:GLN:HB3	1:A:148:LEU:HG	2.02	0.41
1:A:230:LEU:HD12	1:A:230:LEU:HA	1.81	0.41
1:B:195:ASP:HA	1:B:196:PRO:HD3	1.96	0.41
1:B:168:SER:O	1:B:169:GLN:CB	2.65	0.41
1:A:59:ASP:OD2	1:A:59:ASP:N	2.54	0.41
1:B:219:ILE:HD11	3:B:652:GTG:O3D	2.20	0.41
1:B:208:TRP:CE2	1:B:216:LEU:HA	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:ILE:HG22	1:A:273:TYR:HB3	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:701:YT3:Y	2:A:702:YT3:Y[3_655]	2.07	0.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	298/337 (88%)	285 (96%)	12 (4%)	1 (0%)	46	35
1	B	295/337 (88%)	284 (96%)	10 (3%)	1 (0%)	46	35
All	All	593/674 (88%)	569 (96%)	22 (4%)	2 (0%)	46	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	169	GLN
1	A	145	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/295 (91%)	253 (94%)	16 (6%)	24	12
1	B	267/295 (90%)	247 (92%)	20 (8%)	17	7
All	All	536/590 (91%)	500 (93%)	36 (7%)	20	9

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	40	VAL
1	B	42	LEU
1	B	55	GLU
1	B	72	SER
1	B	84	LEU
1	B	141	GLN
1	B	145	ARG
1	B	169	GLN
1	B	172	SER
1	B	203	ILE
1	B	205	ASP
1	B	207	LYS
1	B	217	TYR
1	B	236	GLU
1	B	245	LEU
1	B	253	LEU
1	B	255	ARG
1	B	270	LEU
1	B	276	LEU
1	B	298	LEU
1	A	58	ARG
1	A	99	THR
1	A	122	ARG
1	A	146	GLN
1	A	147	ASP
1	A	150	LEU
1	A	159	ARG
1	A	161	ILE
1	A	163	LEU
1	A	197	SER
1	A	203	ILE
1	A	217	TYR
1	A	270	LEU
1	A	298	LEU
1	A	336	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	337	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	123	GLN
1	A	146	GLN
1	A	237	HIS
1	A	246	HIS
1	A	249	GLN
1	A	314	GLN
1	A	336	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GTG	A	651	-	43,57,57	1.86	8 (18%)	53,90,90	2.29	20 (37%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GTG	B	652	-	43,57,57	2.00	7 (16%)	53,90,90	2.05	14 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GTG	A	651	-	-	0/24/64/64	0/6/6/6
3	GTG	B	652	-	-	0/24/64/64	0/6/6/6

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	652	GTG	C5E-C4E	-5.09	1.35	1.51
3	A	651	GTG	C5E-C4E	-4.71	1.36	1.51
3	A	651	GTG	C6B-C5B	-3.34	1.34	1.41
3	A	651	GTG	C5B-C4B	-3.10	1.33	1.40
3	B	652	GTG	C8A-N9A	2.19	1.37	1.33
3	A	651	GTG	C8A-N7A	2.49	1.37	1.33
3	B	652	GTG	O4E-C1E	2.93	1.44	1.41
3	A	651	GTG	C6B-N1B	3.29	1.39	1.33
3	B	652	GTG	C8A-N7A	3.32	1.39	1.33
3	B	652	GTG	O4D-C1D	3.43	1.45	1.41
3	A	651	GTG	C8B-N7B	3.50	1.41	1.34
3	A	651	GTG	O4E-C1E	4.67	1.47	1.41
3	A	651	GTG	C4A-N3A	5.17	1.43	1.35
3	B	652	GTG	C8B-N7B	6.48	1.47	1.34
3	B	652	GTG	C4A-N3A	7.11	1.46	1.35

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	651	GTG	N3B-C2B-N1B	-5.40	119.22	127.44
3	B	652	GTG	N3A-C2A-N1A	-5.10	119.67	127.44
3	B	652	GTG	C6A-C5A-C4A	-4.86	115.09	120.90
3	A	651	GTG	PG-O3B-PB	-4.57	119.91	132.73
3	B	652	GTG	C5B-C6B-N1B	-4.43	117.53	123.59
3	A	651	GTG	O4E-C1E-N9B	-4.24	99.23	108.10
3	A	651	GTG	C5B-C6B-N1B	-4.19	117.87	123.59
3	B	652	GTG	C2E-C1E-N9B	-4.18	107.91	114.29
3	B	652	GTG	N3B-C2B-N1B	-4.10	121.20	127.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	651	GTG	C4E-O4E-C1E	-3.86	105.47	109.72
3	A	651	GTG	C5A-C6A-N1A	-3.86	118.31	123.59
3	A	651	GTG	O4D-C1D-N9A	-2.87	102.09	108.10
3	A	651	GTG	C2D-C1D-N9A	-2.80	110.01	114.29
3	B	652	GTG	C1D-N9A-C4A	-2.76	122.78	126.94
3	A	651	GTG	C6A-C5A-C4A	-2.68	117.70	120.90
3	A	651	GTG	C1E-N9B-C4B	-2.61	123.00	126.94
3	A	651	GTG	N3A-C2A-N1A	-2.49	123.65	127.44
3	B	652	GTG	PB-O3A-PA	-2.46	125.83	132.73
3	B	652	GTG	C4E-O4E-C1E	-2.18	107.32	109.72
3	A	651	GTG	O3E-C3E-C4E	-2.17	104.55	111.05
3	A	651	GTG	O2G-PG-O1G	2.14	124.12	112.53
3	A	651	GTG	N2B-C2B-N3B	2.31	122.23	117.80
3	A	651	GTG	O4E-C4E-C5E	2.36	117.75	109.32
3	B	652	GTG	O2G-PG-O3B	2.37	115.86	105.09
3	B	652	GTG	O5E-C5E-C4E	2.54	118.49	109.12
3	A	651	GTG	O3A-PA-O5D	2.78	110.31	102.94
3	A	651	GTG	O4E-C4E-C3E	2.83	110.84	105.15
3	A	651	GTG	C2E-C1E-N9B	2.84	118.64	114.29
3	B	652	GTG	O4E-C4E-C3E	2.93	111.05	105.15
3	B	652	GTG	O4E-C1E-N9B	3.43	115.27	108.10
3	B	652	GTG	C6B-N1B-C2B	3.89	121.33	115.94
3	A	651	GTG	C6B-N1B-C2B	4.16	121.71	115.94
3	B	652	GTG	C6A-N1A-C2A	4.88	122.71	115.94
3	A	651	GTG	C6A-N1A-C2A	5.05	122.94	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	651	GTG	1	0
3	B	652	GTG	3	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	300/337 (89%)	-0.07	9 (3%) 54 57	17, 30, 55, 79	0
1	B	297/337 (88%)	0.15	13 (4%) 38 41	21, 37, 60, 76	0
All	All	597/674 (88%)	0.04	22 (3%) 45 49	17, 35, 57, 79	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	337	SER	7.9
1	A	38	ALA	6.6
1	A	336	GLN	4.7
1	B	71	ALA	4.5
1	B	73	GLY	4.4
1	A	39	PRO	4.2
1	B	171	LEU	4.1
1	A	145	ARG	4.0
1	B	74	ASP	3.9
1	B	169	GLN	3.8
1	B	72	SER	3.7
1	A	111	ASP	3.6
1	B	336	GLN	3.4
1	A	146	GLN	3.2
1	B	211	GLN	3.0
1	B	170	SER	2.6
1	B	219	ILE	2.5
1	A	168	SER	2.4
1	B	168	SER	2.2
1	B	183	LYS	2.1
1	B	141	GLN	2.1
1	A	113	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GTG	B	652	52/52	0.93	0.17	0.59	35,52,89,90	11
3	GTG	A	651	52/52	0.98	0.07	-0.53	17,25,39,41	0
2	YT3	A	702	1/1	0.98	0.13	-	94,94,94,94	0
2	YT3	A	703	1/1	0.98	0.20	-	66,66,66,66	0
2	YT3	A	701	1/1	0.98	0.09	-	55,55,55,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.