



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:04 PM GMT

PDB ID : 3STC
Title : Crystal structure of loop 7 truncated mutant of 3-deoxy-D-manno-octulosonate 8-phosphate synthase (KDO8PS) from *Neisseria meningitidis*
Authors : Allison, T.M.; Jameson, G.B.; Parker, E.J.
Deposited on : 2011-07-09
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5832 (1.94-1.90)
Clashscore	102246	6540 (1.94-1.90)
Ramachandran outliers	100387	6464 (1.94-1.90)
Sidechain outliers	100360	6465 (1.94-1.90)
RSRZ outliers	91569	5846 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	<div> <div>12%</div> <div>81% 11% 7%</div> </div>
1	B	269	<div> <div>4%</div> <div>84% 10% 6%</div> </div>
1	C	269	<div> <div>%</div> <div>86% 10% ..</div> </div>
1	D	269	<div> <div>6%</div> <div>86% 9% ..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CL	D	270	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8613 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-dehydro-3-deoxyphosphooctonate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	0	1	0
			1908	1229	317	351	11			
1	B	254	Total	C	N	O	S	0	5	0
			1969	1268	328	362	11			
1	C	260	Total	C	N	O	S	0	4	0
			2028	1303	343	371	11			
1	D	259	Total	C	N	O	S	0	2	0
			2001	1287	337	366	11			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLN	DELETION	UNP Q9JZ55
A	?	-	THR	DELETION	UNP Q9JZ55
A	?	-	ARG	DELETION	UNP Q9JZ55
A	?	-	ASP	DELETION	UNP Q9JZ55
A	?	-	ALA	DELETION	UNP Q9JZ55
A	?	-	GLY	DELETION	UNP Q9JZ55
A	?	-	SER	DELETION	UNP Q9JZ55
A	?	-	ALA	DELETION	UNP Q9JZ55
A	?	-	ALA	DELETION	UNP Q9JZ55
A	?	-	SER	DELETION	UNP Q9JZ55
A	?	-	GLY	DELETION	UNP Q9JZ55
B	?	-	GLN	DELETION	UNP Q9JZ55
B	?	-	THR	DELETION	UNP Q9JZ55
B	?	-	ARG	DELETION	UNP Q9JZ55
B	?	-	ASP	DELETION	UNP Q9JZ55
B	?	-	ALA	DELETION	UNP Q9JZ55
B	?	-	GLY	DELETION	UNP Q9JZ55
B	?	-	SER	DELETION	UNP Q9JZ55
B	?	-	ALA	DELETION	UNP Q9JZ55
B	?	-	ALA	DELETION	UNP Q9JZ55
B	?	-	SER	DELETION	UNP Q9JZ55

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	GLY	DELETION	UNP Q9JZ55
C	?	-	GLN	DELETION	UNP Q9JZ55
C	?	-	THR	DELETION	UNP Q9JZ55
C	?	-	ARG	DELETION	UNP Q9JZ55
C	?	-	ASP	DELETION	UNP Q9JZ55
C	?	-	ALA	DELETION	UNP Q9JZ55
C	?	-	GLY	DELETION	UNP Q9JZ55
C	?	-	SER	DELETION	UNP Q9JZ55
C	?	-	ALA	DELETION	UNP Q9JZ55
C	?	-	ALA	DELETION	UNP Q9JZ55
C	?	-	SER	DELETION	UNP Q9JZ55
C	?	-	GLY	DELETION	UNP Q9JZ55
D	?	-	GLN	DELETION	UNP Q9JZ55
D	?	-	THR	DELETION	UNP Q9JZ55
D	?	-	ARG	DELETION	UNP Q9JZ55
D	?	-	ASP	DELETION	UNP Q9JZ55
D	?	-	ALA	DELETION	UNP Q9JZ55
D	?	-	GLY	DELETION	UNP Q9JZ55
D	?	-	SER	DELETION	UNP Q9JZ55
D	?	-	ALA	DELETION	UNP Q9JZ55
D	?	-	ALA	DELETION	UNP Q9JZ55
D	?	-	SER	DELETION	UNP Q9JZ55
D	?	-	GLY	DELETION	UNP Q9JZ55

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Cl 2 2	0	0
2	A	1	Total Cl 1 1	0	0
2	D	2	Total Cl 2 2	0	0
2	C	2	Total Cl 2 2	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total Na 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	128	Total 128	O 128	0	0
4	B	205	Total 205	O 205	0	0
4	C	204	Total 204	O 204	0	0
4	D	162	Total 162	O 162	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase




- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase




- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase



- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.78Å 85.76Å 163.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.56 – 1.91 36.56 – 1.91	Depositor EDS
% Data completeness (in resolution range)	98.9 (36.56-1.91) 99.0 (36.56-1.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.90 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.6.0094	Depositor
R, R_{free}	0.207 , 0.245 0.210 , 0.248	Depositor DCC
R_{free} test set	4415 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	21.9	Xtriage
Anisotropy	0.199	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.7	EDS
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 88211 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8613	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/1944	0.60	1/2631 (0.0%)
1	B	0.53	0/2015	0.62	0/2728
1	C	0.50	0/2071	0.64	2/2802 (0.1%)
1	D	0.48	0/2041	0.58	1/2762 (0.0%)
All	All	0.50	0/8071	0.61	4/10923 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	ARG	NE-CZ-NH2	-5.46	117.57	120.30
1	D	218	ARG	NE-CZ-NH2	-5.18	117.71	120.30
1	C	218	ARG	NE-CZ-NH2	-5.16	117.72	120.30
1	C	241	LEU	CA-CB-CG	5.13	127.11	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1908	0	1947	17	0
1	B	1969	0	2022	23	0
1	C	2028	0	2086	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2001	0	2057	20	0
2	A	1	0	0	1	0
2	B	2	0	0	1	0
2	C	2	0	0	1	0
2	D	2	0	0	2	0
3	C	1	0	0	0	0
4	A	128	0	0	3	0
4	B	205	0	0	4	0
4	C	204	0	0	5	0
4	D	162	0	0	0	0
All	All	8613	0	8112	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:72:GLU:HG2	4:C:356:HOH:O	1.77	0.83
1:B:242:PRO:HB2	1:B:245:LEU:HD12	1.64	0.78
1:D:204:ARG:HD3	1:D:240:ALA:O	1.94	0.68
1:A:206:GLN:HB3	1:B:179:LEU:HD22	1.77	0.66
1:B:28:LEU:HD11	1:B:80:LYS:HG2	1.77	0.66
1:B:13[A]:ASN:ND2	4:B:687:HOH:O	2.30	0.65
1:A:60:ARG:NH1	1:C:120:ASP:OD1	2.23	0.64
1:A:127:LYS:NZ	4:A:342:HOH:O	2.28	0.64
1:D:137:PRO:HB2	1:D:140:LEU:HD13	1.78	0.63
1:A:256:LEU:C	1:A:256:LEU:HD23	2.22	0.59
1:A:22:ILE:HD11	1:A:25:LEU:HD23	1.84	0.58
1:B:119:THR:HG23	1:D:63:ILE:HG22	1.87	0.56
1:A:179:LEU:HD22	1:B:206:GLN:HB3	1.88	0.56
1:B:63:ILE:HD11	1:D:154:GLU:HG3	1.87	0.55
1:D:139:PHE:CD2	1:D:140:LEU:HD12	2.41	0.55
1:D:224:LEU:C	1:D:224:LEU:HD12	2.27	0.54
1:C:195:PHE:CE2	1:C:197[B]:VAL:HG12	2.44	0.52
1:B:242:PRO:CB	1:B:245:LEU:HD12	2.36	0.51
1:B:241:LEU:HD22	1:B:242:PRO:HD2	1.92	0.50
1:B:118:GLN:NE2	4:B:533:HOH:O	2.42	0.50
1:B:208:LEU:HD21	1:B:252:ARG:HD3	1.95	0.49
1:B:198:THR:HB	1:B:225:GLU:HB2	1.93	0.49
1:B:13[A]:ASN:CG	4:B:687:HOH:O	2.51	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:24[B]:VAL:CG1	1:B:68:GLY:HA2	2.44	0.47
1:B:109:ILE:HG13	1:B:125:MET:HE2	1.96	0.47
1:C:267:THR:HG22	4:C:476:HOH:O	2.15	0.47
1:B:28:LEU:HD11	1:B:80:LYS:CG	2.45	0.47
1:B:63:ILE:HA	1:D:119:THR:HG21	1.96	0.47
1:A:224:LEU:HD12	1:A:224:LEU:C	2.35	0.47
1:C:13:ASN:HD21	1:C:265:ILE:HD11	1.78	0.47
1:A:195:PHE:CE2	1:A:197[B]:VAL:HG12	2.50	0.47
1:C:140:LEU:HD11	1:C:144:GLN:HB3	1.97	0.46
1:A:196:ASP:HA	1:A:223:PHE:HB3	1.98	0.46
1:B:138:GLN:HG2	2:B:270:CL:CL	2.53	0.45
1:D:204:ARG:HB2	1:D:238:PRO:O	2.16	0.45
1:A:43:ARG:NH1	4:A:469:HOH:O	2.49	0.45
1:C:206:GLN:HB3	1:D:179:LEU:HD22	1.98	0.45
1:C:107:ASP:O	1:C:130[A]:ASN:ND2	2.51	0.44
1:C:138:GLN:HG2	2:C:270:CL:CL	2.54	0.44
1:A:180:GLY:HA3	4:A:307:HOH:O	2.16	0.44
1:D:139:PHE:CE2	1:D:140:LEU:CD1	3.00	0.44
1:B:13[A]:ASN:OD1	1:B:265:ILE:HD11	2.18	0.44
1:B:63:ILE:HG22	1:D:119:THR:HG23	2.00	0.43
1:C:82:LYS:HE2	4:C:358:HOH:O	2.18	0.43
1:A:138:GLN:HG2	2:A:270:CL:CL	2.55	0.43
1:C:136:LYS:O	1:C:165:ARG:HD2	2.19	0.43
1:B:52:LYS:HE2	1:B:223:PHE:CZ	2.54	0.43
1:C:210:LEU:HB2	1:D:179:LEU:HD21	1.99	0.43
1:C:45:LEU:O	1:C:254:LYS:HE3	2.19	0.43
1:C:140:LEU:HD12	4:C:481:HOH:O	2.18	0.43
1:D:139:PHE:CD2	1:D:140:LEU:CD1	3.02	0.43
1:A:45:LEU:O	1:A:254:LYS:HE3	2.19	0.42
1:D:138:GLN:NE2	2:D:270:CL:CL	2.74	0.42
1:C:203:ARG:NH2	4:C:302:HOH:O	2.52	0.42
1:D:100:GLN:O	1:D:104:GLU:HG2	2.19	0.42
1:A:92:ASP:N	1:A:92:ASP:OD1	2.50	0.42
1:A:28:LEU:HD11	1:A:32:LEU:HD11	2.02	0.42
1:B:13[B]:ASN:OD1	4:B:687:HOH:O	0.42	0.42
1:D:38:TYR:HB3	1:D:49:TYR:CZ	2.55	0.42
1:B:226:SER:CB	1:B:241:LEU:H	2.31	0.42
1:C:197[B]:VAL:CG1	1:C:222:LEU:HD11	2.50	0.41
1:C:241:LEU:HD13	1:C:242:PRO:O	2.20	0.41
1:A:141:SER:HB3	1:C:171:TYR:HB3	2.01	0.41
1:A:110:GLN:NE2	1:A:135:LYS:HE3	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:165:ARG:NH2	2:D:270:CL:CL	2.91	0.41
1:D:98:GLN:O	1:D:102:VAL:HG23	2.21	0.41
1:C:266:LEU:HD23	1:D:208:LEU:HD22	2.02	0.40
1:D:256:LEU:HD23	1:D:256:LEU:C	2.42	0.40
1:C:169:PHE:HB2	1:C:175:VAL:HG12	2.03	0.40
1:C:267:THR:O	1:C:267:THR:HG23	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/269 (91%)	240 (98%)	6 (2%)	0	100	100
1	B	255/269 (95%)	249 (98%)	6 (2%)	0	100	100
1	C	260/269 (97%)	256 (98%)	3 (1%)	1 (0%)	39	27
1	D	257/269 (96%)	254 (99%)	2 (1%)	1 (0%)	39	27
All	All	1018/1076 (95%)	999 (98%)	17 (2%)	2 (0%)	52	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	218	ARG
1	D	218	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/230 (91%)	204 (97%)	6 (3%)	50	39
1	B	219/230 (95%)	213 (97%)	6 (3%)	52	42
1	C	227/230 (99%)	222 (98%)	5 (2%)	60	52
1	D	223/230 (97%)	217 (97%)	6 (3%)	52	42
All	All	879/920 (96%)	856 (97%)	23 (3%)	52	44

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	GLU
1	A	62	SER
1	A	140	LEU
1	A	165	ARG
1	A	244	HIS
1	A	266	LEU
1	B	60	ARG
1	B	104	GLU
1	B	146	LYS
1	B	165	ARG
1	B	239	SER
1	B	241	LEU
1	C	13	ASN
1	C	165	ARG
1	C	201	LEU
1	C	229	ASP
1	C	241	LEU
1	D	40	GLU
1	D	59	ASN
1	D	117	ARG
1	D	118	GLN
1	D	165	ARG
1	D	239	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	147	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	249/269 (92%)	0.77	32 (12%) 5 5	14, 33, 69, 95	1 (0%)
1	B	254/269 (94%)	0.19	11 (4%) 39 43	14, 24, 43, 63	0
1	C	260/269 (96%)	0.02	4 (1%) 76 79	15, 23, 44, 72	0
1	D	259/269 (96%)	0.31	15 (5%) 26 30	15, 30, 59, 78	0
All	All	1022/1076 (94%)	0.32	62 (6%) 25 28	14, 27, 58, 95	1 (0%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	243	LEU	7.4
1	A	66	TYR	7.3
1	B	244	HIS	6.8
1	A	266	LEU	6.5
1	B	242	PRO	5.8
1	B	245	LEU	5.2
1	B	243	LEU	4.6
1	D	238	PRO	4.6
1	B	241	LEU	4.6
1	A	224	LEU	4.5
1	B	240	ALA	4.4
1	A	244	HIS	4.4
1	A	69	VAL	4.3
1	A	64	HIS	4.2
1	A	29	ASP	4.0
1	A	61	SER	4.0
1	A	202	GLY	4.0
1	A	265	ILE	3.9
1	B	239	SER	3.9
1	C	238	PRO	3.9
1	D	64	HIS	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	228	PRO	3.7
1	D	76	LYS	3.2
1	A	68	GLY	3.2
1	A	73	GLU	3.2
1	B	172	ASP	3.2
1	B	226	SER	3.1
1	D	66	TYR	3.1
1	D	227	HIS	3.1
1	B	205	ALA	2.9
1	D	63	ILE	2.9
1	A	222	LEU	2.8
1	A	83	ALA	2.7
1	A	140	LEU	2.7
1	A	67	ARG	2.7
1	D	205	ALA	2.7
1	A	76	LYS	2.6
1	D	27	SER	2.6
1	A	62	SER	2.6
1	D	80	LYS	2.5
1	A	223	PHE	2.5
1	A	114	PHE	2.4
1	A	65	SER	2.4
1	A	28	LEU	2.3
1	A	204	ARG	2.3
1	C	61	SER	2.3
1	A	80	LYS	2.3
1	B	33	GLN	2.3
1	A	30	SER	2.2
1	A	197[A]	VAL	2.2
1	A	74	GLY	2.2
1	D	30	SER	2.2
1	D	239	SER	2.2
1	D	73	GLU	2.2
1	A	27	SER	2.2
1	C	59	ASN	2.1
1	A	33	GLN	2.1
1	D	228	PRO	2.0
1	A	19	PHE	2.0
1	D	72	GLU	2.0
1	D	29	ASP	2.0
1	A	225	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NA	C	272	1/1	0.94	0.11	0.34	32,32,32,32	0
2	CL	D	271	1/1	0.97	0.05	-2.19	32,32,32,32	0
2	CL	C	270	1/1	0.96	0.06	-2.25	34,34,34,34	0
2	CL	B	270	1/1	0.98	0.07	-2.59	30,30,30,30	0
2	CL	A	270	1/1	0.92	0.07	-2.60	44,44,44,44	0
2	CL	C	271	1/1	0.98	0.05	-2.64	27,27,27,27	0
2	CL	D	270	1/1	0.98	0.04	-4.12	40,40,40,40	0
2	CL	B	271	1/1	0.97	0.06	-	25,25,25,25	1

6.5 Other polymers [i](#)

There are no such residues in this entry.