



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:04 PM GMT

PDB ID : 3STI
Title : Crystal structure of the protease domain of DegQ from Escherichia coli
Authors : Sawa, J.; Malet, H.; Krojer, T.; Canellas, F.; Ehrmann, M.; Clausen, T.
Deposited on : 2011-07-11
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	<div> <div></div> <div>66% 9% 25%</div> </div>
1	B	245	<div> <div>2%</div> <div>64% 11% 25%</div> </div>
1	C	245	<div> <div>2%</div> <div>64% 9% 25%</div> </div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 4016 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protease degQ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	184	Total	C	N	O	S	0	0	0
			1341	847	230	262	2			
1	B	184	Total	C	N	O	S	0	0	0
			1342	847	233	260	2			
1	C	183	Total	C	N	O	S	0	0	0
			1333	839	232	260	2			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	238	LEU	-	EXPRESSION TAG	UNP P39099
A	239	GLU	-	EXPRESSION TAG	UNP P39099
A	240	HIS	-	EXPRESSION TAG	UNP P39099
A	241	HIS	-	EXPRESSION TAG	UNP P39099
A	242	HIS	-	EXPRESSION TAG	UNP P39099
A	243	HIS	-	EXPRESSION TAG	UNP P39099
A	244	HIS	-	EXPRESSION TAG	UNP P39099
A	245	HIS	-	EXPRESSION TAG	UNP P39099
B	238	LEU	-	EXPRESSION TAG	UNP P39099
B	239	GLU	-	EXPRESSION TAG	UNP P39099
B	240	HIS	-	EXPRESSION TAG	UNP P39099
B	241	HIS	-	EXPRESSION TAG	UNP P39099
B	242	HIS	-	EXPRESSION TAG	UNP P39099
B	243	HIS	-	EXPRESSION TAG	UNP P39099
B	244	HIS	-	EXPRESSION TAG	UNP P39099
B	245	HIS	-	EXPRESSION TAG	UNP P39099
C	238	LEU	-	EXPRESSION TAG	UNP P39099
C	239	GLU	-	EXPRESSION TAG	UNP P39099
C	240	HIS	-	EXPRESSION TAG	UNP P39099
C	241	HIS	-	EXPRESSION TAG	UNP P39099
C	242	HIS	-	EXPRESSION TAG	UNP P39099
C	243	HIS	-	EXPRESSION TAG	UNP P39099
C	244	HIS	-	EXPRESSION TAG	UNP P39099

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	245	HIS	-	EXPRESSION TAG	UNP P39099


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Protease degQ



• Molecule 1: Protease degQ



• Molecule 1: Protease degQ



4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	70.87Å 70.87Å 152.01Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.80 – 2.60 19.80 – 2.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (19.80-2.60) 90.0 (19.80-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.59Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.222 , 0.257 0.206 , 0.232	Depositor DCC
R_{free} test set	1204 reflections (5.38%)	DCC
Wilson B-factor (Å ²)	49.5	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.0	EDS
Estimated twinning fraction	0.077 for -h,-k,l 0.288 for h,-h-k,-l 0.086 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.24$	Xtriage
Outliers	0 of 23603 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4016	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/1352	0.71	0/1829
1	B	0.50	0/1352	0.70	0/1827
1	C	0.48	0/1344	0.69	0/1817
All	All	0.50	0/4048	0.70	0/5473

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1341	0	1384	8	0
1	B	1342	0	1388	10	0
1	C	1333	0	1374	12	0
All	All	4016	0	4146	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:177:GLN:HE21	1:C:215:ILE:HG21	1.60	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:177:GLN:HE21	1:B:215:ILE:HG21	1.65	0.59
1:C:95:ASN:H	1:C:95:ASN:HD22	1.56	0.54
1:C:105:ILE:HD11	1:C:115:LEU:HD23	1.91	0.52
1:B:64:LEU:HD22	1:B:148:PHE:H	1.76	0.51
1:B:70:ILE:HG21	1:B:78:LEU:HD12	1.93	0.50
1:C:159:VAL:HG22	1:C:176:ILE:HD11	1.93	0.50
1:B:28:SER:HB3	1:B:93:GLN:HB3	1.94	0.49
1:B:81:ASN:HD22	1:B:109:ASP:HA	1.78	0.49
1:A:28:SER:HB3	1:A:93:GLN:HB3	1.96	0.47
1:A:194:LEU:HB2	1:C:12:LEU:HD22	1.96	0.47
1:A:174:ASN:HB3	1:A:223:MET:SD	2.56	0.46
1:B:64:LEU:CD2	1:B:148:PHE:H	2.27	0.46
1:C:172:LEU:HD21	1:C:217:PHE:HE2	1.81	0.45
1:B:147:PRO:C	1:B:149:GLY:H	2.20	0.45
1:C:70:ILE:HG13	1:C:129:ILE:HD11	1.99	0.45
1:A:22:VAL:HG11	1:A:192:LEU:HD11	1.99	0.45
1:B:172:LEU:HB3	1:B:175:PHE:HD2	1.83	0.44
1:A:70:ILE:HG13	1:A:129:ILE:HD11	1.98	0.44
1:B:80:ASN:HB2	1:B:83:VAL:HG23	2.01	0.43
1:A:105:ILE:HD11	1:A:117:GLN:HE21	1.83	0.42
1:C:155:THR:HG21	1:C:181:SER:HB2	2.00	0.42
1:C:143:ALA:HB3	1:C:155:THR:OG1	2.20	0.42
1:C:147:PRO:C	1:C:149:GLY:H	2.24	0.41
1:A:158:ILE:HG21	1:C:15:LEU:HD12	2.03	0.41
1:B:15:LEU:HD12	1:C:158:ILE:HG21	2.03	0.40
1:A:205:ILE:H	1:A:205:ILE:HG13	1.76	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	176/245 (72%)	169 (96%)	6 (3%)	1 (1%)	30	56
1	B	174/245 (71%)	166 (95%)	6 (3%)	2 (1%)	17	36
1	C	175/245 (71%)	169 (97%)	6 (3%)	0	100	100
All	All	525/735 (71%)	504 (96%)	18 (3%)	3 (1%)	30	56

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	173	GLU
1	A	149	GLY
1	B	166	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/193 (76%)	137 (94%)	9 (6%)	23	45
1	B	146/193 (76%)	137 (94%)	9 (6%)	23	45
1	C	145/193 (75%)	136 (94%)	9 (6%)	23	45
All	All	437/579 (76%)	410 (94%)	27 (6%)	23	45

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	MET
1	A	21	LYS
1	A	89	LYS
1	A	107	SER
1	A	127	ILE
1	A	148	PHE
1	A	184	ARG
1	A	205	ILE
1	A	213	VAL
1	B	18	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	21	LYS
1	B	64	LEU
1	B	127	ILE
1	B	184	ARG
1	B	192	LEU
1	B	202	ASN
1	B	206	LEU
1	B	229	GLN
1	C	18	MET
1	C	19	LEU
1	C	95	ASN
1	C	113	ILE
1	C	120	ASN
1	C	160	SER
1	C	172	LEU
1	C	184	ARG
1	C	215	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	GLN
1	B	81	ASN
1	B	202	ASN
1	C	81	ASN
1	C	95	ASN
1	C	177	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	184/245 (75%)	0.07	1 (0%) 91 90	29, 46, 67, 77	0
1	B	184/245 (75%)	0.21	5 (2%) 58 51	29, 50, 79, 105	0
1	C	183/245 (74%)	0.21	4 (2%) 65 59	27, 51, 78, 95	0
All	All	551/735 (74%)	0.16	10 (1%) 71 66	27, 49, 76, 105	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	167	LEU	5.1
1	C	206	LEU	4.1
1	B	164	ARG	4.1
1	C	174	ASN	2.6
1	B	165	SER	2.5
1	C	227	LEU	2.4
1	C	167	LEU	2.3
1	A	184	ARG	2.1
1	B	30	ARG	2.0
1	B	174	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.