Full wwPDB X-ray Structure Validation Report (i)

Jan 31, 2016 – 10:16 PM GMT

PDB ID	:	1SUZ
Title	:	The structure of K92A EcoRV bound to cognate DNA and Mg2+
Authors	:	Horton, N.C.; Perona, J.J.
Deposited on	:	2004-03-26
Resolution	:	1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	C	11	27% 64% 9%
1	D	11	27% 55% 9% 9%
2	A	244	77% 21% •
2	B	244	83% 14% • •

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4565 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*C*AP*AP*GP*AP*TP*AP*TP*CP*TP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	C	10	202	99	36	58	9	0	0	0
1	D	10	201	98	37	57	9	0	0	0

- Molecule 2 is a protein called Type II restriction enzyme EcoRV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	243	1942	1252	320	369	1	0	0	0
2	B	237	1913	1234	313	365	1	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	92	ALA	LYS	ENGINEERED	UNP P04390
B	92	ALA	LYS	ENGINEERED	UNP P04390

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	B	1	1	1	0	0
3	A	1	1	1	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
4	B	1	1	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
4	A	1	1	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	147	147	147	0	0
			Total	O		
5	B	113	113	113	0	0
			Total	O		
5	C	23	23	23	0	0
			Total	O		
5	D	20	20	20	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 5'-D(*C*AP*AP*GP*AP*TP*AP*TP*CP*TP*T)-3'

Chain C:

- Molecule 1: 5'-D(*C*AP*AP*GP*AP*TP*AP*TP*CP*TP*T)-3'

Chain D:

- Molecule 2: Type II restriction enzyme EcoRV

Chain A:



- Molecule 2: Type II restriction enzyme EcoRV

Chain B:



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	47.41Å 48.79Å 63.40Å 97.10° 108.80° 107.30°	Depositor
Resolution (Å)	20.00 – 1.80	Depositor
% Data completeness (in resolution range)	0.9 (20.00-1.80)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.197 , 0.244	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4565	wwPDB-VP
Average B, all atoms (Å^2)	23.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	C	0.92	0/226	1.10	0/347
1	D	1.01	0/225	1.13	2/345 (0.6%)
2	A	0.95	0/1992	0.89	1/2709 (0.0%)
2	B	0.93	0/1962	0.87	0/2666
All	All	0.94	0/4405	0.91	3/6067 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
2	B	0	1
All	All	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	8	DT	O5'-P-OP2	-5.49	100.76	105.70
1	D	7	DA	OP2-P-O3'	5.34	116.95	105.20
2	A	134	ILE	N-CA-C	-5.13	97.15	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	95	TYR	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	C	2	DA	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	202	0	116	8	0
1	D	201	0	115	10	0
2	A	1942	0	1826	49	0
2	B	1913	0	1802	25	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	147	0	0	0	0
5	B	113	0	0	0	0
5	C	23	0	0	2	0
5	D	20	0	0	0	0
All	All	4565	0	3859	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:69:GLN:HE22	2:B:36:ASP:HB2	1.40	0.86
1:C:8:DT:H4'	2:A:37:THR:HG21	1.62	0.82
1:D:1:DC:H3'	2:A:180:LEU:HD11	1.63	0.79
2:B:8:ILE:HG12	2:B:170:LEU:HB3	1.66	0.78
2:A:113:PHE:HA	2:A:116:ASN:O	1.87	0.74
5:C:40:HOH:O	2:A:37:THR:HG23	1.89	0.73
1:D:2:DA:P	2:A:180:LEU:HD21	2.33	0.68
2:A:29:LYS:HE2	2:A:150:THR:HG21	1.78	0.65
1:C:8:DT:H4'	2:A:37:THR:CG2	2.26	0.65
2:A:85:LYS:HA	2:A:129:ILE:HG23	1.81	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:103:ILE:HD12	2:A:137:VAL:HG21	1.83	0.60
2:A:143:THR:HG23	2:A:147:SER:OG	2.02	0.59
2:A:116:ASN:O	2:A:118:THR:N	2.35	0.59
2:A:103:ILE:CD1	2:A:137:VAL:HG21	2.35	0.57
1:D:9:DC:H4'	2:A:69:GLN:HG3	1.86	0.57
2:A:49:ARG:HG2	2:A:75:PHE:HZ	1.70	0.56
2:A:148:LEU:HD13	2:B:53:ASN:OD1	2.05	0.55
2:B:147:SER:HB2	2:B:160:PRO:HG3	1.88	0.54
2:A:41:SER:O	2:A:45:GLU:HG3	2.08	0.53
2:A:153:ILE:HD13	2:B:153:ILE:HG23	1.89	0.53
2:B:14:GLU:HG2	2:B:51:ILE:HD11	1.91	0.52
2:B:8:ILE:HG12	2:B:170:LEU:CB	2.37	0.52
2:A:35:SER:HB3	2:A:162:PRO:HB2	1.92	0.51
2:A:40:LEU:HD23	2:A:138:TYR:CE2	2.46	0.51
2:A:45:GLU:O	2:A:49:ARG:HG3	2.11	0.50
2:A:49:ARG:HG2	2:A:75:PHE:CZ	2.47	0.50
2:A:96:THR:HG22	2:A:139:THR:HA	1.94	0.49
2:B:36:ASP:O	2:B:40:LEU:HD23	2.12	0.49
2:B:29:LYS:HE2	2:B:150:THR:HG21	1.95	0.49
2:A:14:GLU:HG2	2:A:51:ILE:HD11	1.95	0.48
1:C:11:DT:H3	1:D:2:DA:H61	1.59	0.48
1:D:1:DC:H2"	1:D:2:DA:C8	2.49	0.48
1:C:6:DT:O4'	2:A:70:ASN:HA	2.14	0.47
2:A:49:ARG:HB2	2:A:50:PRO:HD3	1.95	0.47
2:A:148:LEU:HD13	2:B:53:ASN:CG	2.35	0.47
1:C:9:DC:H2'	1:C:10:DT:H72	1.96	0.47
2:B:39:VAL:O	2:B:43:ILE:HD13	2.16	0.46
2:B:141:VAL:HG23	2:B:141:VAL:O	2.15	0.46
1:C:11:DT:H3	1:D:2:DA:N6	2.13	0.46
1:D:8:DT:H4'	2:B:37:THR:OG1	2.15	0.46
2:A:154:ASN:OD1	2:A:155:GLU:HG3	2.15	0.46
2:B:41:SER:O	2:B:45:GLU:HG3	2.16	0.46
1:C:5:DA:OP1	2:A:119:LYS:HD3	2.16	0.46
2:B:35:SER:HA	2:B:162:PRO:HB2	1.98	0.46
2:A:35:SER:CB	2:A:162:PRO:HB2	2.46	0.45
2:A:91:ILE:HD13	2:A:134:ILE:HB	1.97	0.45
1:D:2:DA:OP2	2:A:180:LEU:HD11	2.16	0.45
2:B:137:VAL:HG22	2:B:167:LYS:HB2	1.98	0.45
2:A:85:LYS:HA	2:A:129:ILE:CG2	2.46	0.45
2:A:69:GLN:NE2	2:B:36:ASP:HB2	2.18	0.45
2:B:11:LEU:HD23	2:B:51:ILE:HD13	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:112:SER:CB	2:A:119:LYS:HD2	2.46	0.44
5:C:172:HOH:O	2:A:119:LYS:HD3	2.16	0.44
2:A:153:ILE:HD13	2:B:153:ILE:CG2	2.46	0.44
2:B:103:ILE:CD1	2:B:137:VAL:HG11	2.48	0.44
2:B:170:LEU:HD23	2:B:170:LEU:C	2.37	0.44
1:D:5:DA:H2'	1:D:6:DT:C6	2.53	0.43
2:A:3:LEU:HD12	2:A:55:ILE:HG22	2.00	0.43
2:B:49:ARG:HB2	2:B:50:PRO:HD3	2.01	0.43
2:A:121:ILE:HG12	2:A:123:TYR:O	2.18	0.43
1:D:2:DA:OP1	2:A:180:LEU:HD21	2.19	0.43
2:A:170:LEU:C	2:A:170:LEU:HD13	2.39	0.43
2:A:23:ILE:HG22	2:B:23:ILE:HG22	2.01	0.42
2:A:85:LYS:HG2	2:A:129:ILE:HG21	2.02	0.42
2:A:16:GLN:HE21	2:A:16:GLN:HB2	1.68	0.42
2:A:20:VAL:HG12	2:A:33:LEU:HD11	2.02	0.41
2:B:4:ARG:O	2:B:8:ILE:HG13	2.20	0.41
2:A:220:GLU:HB2	2:A:226:ARG:HG2	2.03	0.41
2:A:11:LEU:HD23	2:A:51:ILE:HD13	2.02	0.41
1:C:5:DA:H2'	1:C:6:DT:C6	2.55	0.40
2:B:206:PHE:CE1	2:B:233:ILE:HD13	2.56	0.40
2:A:112:SER:HA	2:A:119:LYS:HD2	2.03	0.40
2:A:238:ASN:ND2	2:A:242:ARG:HH12	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	239/244 (98%)	231 (97%)	7 (3%)	1 (0%)	39	23
2	B	231/244 (95%)	222 (96%)	8 (4%)	1 (0%)	39	23
All	All	470/488 (96%)	453 (96%)	15 (3%)	2 (0%)	39	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	117	ASN
2	B	117	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	200/219 (91%)	191 (96%)	9 (4%)	34	16
2	B	199/219 (91%)	194 (98%)	5 (2%)	55	39
All	All	399/438 (91%)	385 (96%)	14 (4%)	43	25

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	16	GLN
2	A	36	ASP
2	A	40	LEU
2	A	57	GLU
2	A	71	HIS
2	A	143	THR
2	A	180	LEU
2	A	188	ASN
2	A	245	LYS
2	B	37	THR
2	B	40	LEU
2	B	71	HIS
2	B	157	ASN
2	B	188	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	16	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	69	GLN
2	A	97	ASN
2	A	238	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates (i)

EDS was not executed - this section will therefore be empty.

6.4 Ligands (i)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers (i)

EDS was not executed - this section will therefore be empty.