



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 17, 2016 – 11:20 AM EDT

PDB ID : 5SW6
Title : Crystal structure of an oxoferryl species of catalase-peroxidase KatG at pH5.6
Authors : Loewen, P.C.
Deposited on : 2016-08-08
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	728	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>8%</div> <div>..</div> </div> </div>
1	B	728	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	MPD	A	807	-	-	-	X
7	MPD	B	806	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 12683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catalase-peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	713	Total	C	N	O	S	0	7	0
			5534	3498	984	1038	14			
1	B	713	Total	C	N	O	S	0	5	0
			5518	3486	981	1037	14			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is OXYGEN MOLECULE (three-letter code: OXY) (formula: O₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O	0	0
			2	2		
4	A	1	Total	O	0	0
			2	2		
4	B	1	Total	O	0	0
			2	2		

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		
5	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 6 is OXYGEN ATOM (three-letter code: O) (formula: O).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	O	0	0
			1	1		
6	A	1	Total	O	0	0
			1	1		

- Molecule 7 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			8	6	2		
7	A	1	Total	C	O	0	0
			8	6	2		
7	B	1	Total	C	O	0	0
			8	6	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	725	Total	O	0	0
			725	725		
8	B	776	Total	O	0	0
			776	776		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Catalase-peroxidase



• Molecule 1: Catalase-peroxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	100.21 Å 114.66 Å 174.38 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 20.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.8 (20.00-1.90) 98.9 (20.00-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.97 (at 1.90 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.141 , 0.175 0.154 , 0.183	Depositor DCC
R_{free} test set	15522 reflections (11.03%)	DCC
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.087	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12683	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, OXY, NA, PO4, O, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.33	24/5694 (0.4%)	1.22	43/7742 (0.6%)
1	B	1.32	20/5669 (0.4%)	1.25	59/7708 (0.8%)
All	All	1.33	44/11363 (0.4%)	1.24	102/15450 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (44) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	726	GLU	CG-CD	10.09	1.67	1.51
1	B	178	GLU	CD-OE2	8.81	1.35	1.25
1	A	725	GLN	CD-OE1	8.61	1.42	1.24
1	B	198	GLU	CG-CD	8.49	1.64	1.51
1	A	726	GLU	CD-OE2	7.78	1.34	1.25
1	B	382	ARG	CZ-NH1	7.72	1.43	1.33
1	A	725	GLN	CG-CD	7.70	1.68	1.51
1	B	410	GLU	CG-CD	7.65	1.63	1.51
1	A	349	GLU	CD-OE2	-7.24	1.17	1.25
1	B	529	GLU	CG-CD	6.98	1.62	1.51
1	A	584	GLU	CG-CD	6.90	1.62	1.51
1	B	128	GLU	CD-OE2	-6.88	1.18	1.25
1	A	178	GLU	CD-OE1	-6.84	1.18	1.25
1	A	407	GLU	CD-OE2	6.80	1.33	1.25
1	A	532	ARG	NE-CZ	-6.60	1.24	1.33
1	A	198	GLU	CG-CD	6.54	1.61	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	442	GLU	CD-OE2	-6.13	1.19	1.25
1	A	736	TRP	CG-CD1	-5.94	1.28	1.36
1	B	512	ASP	CG-OD2	5.86	1.38	1.25
1	B	514	GLU	CD-OE2	-5.85	1.19	1.25
1	B	36	GLY	N-CA	5.84	1.54	1.46
1	B	161	ARG	CZ-NH1	5.83	1.40	1.33
1	B	343	GLU	CD-OE1	5.80	1.32	1.25
1	A	343	GLU	CD-OE1	5.79	1.32	1.25
1	A	349	GLU	CD-OE1	-5.79	1.19	1.25
1	B	337	TRP	CD1-NE1	5.79	1.47	1.38
1	B	82	ARG	CZ-NH2	-5.74	1.25	1.33
1	B	434	TYR	CE2-CZ	-5.74	1.31	1.38
1	B	161	ARG	CD-NE	-5.73	1.36	1.46
1	A	68	TYR	CE1-CZ	5.70	1.46	1.38
1	B	159	TYR	CE2-CZ	-5.68	1.31	1.38
1	A	330	TRP	CG-CD1	-5.68	1.28	1.36
1	A	285	HIS	C-O	5.51	1.33	1.23
1	B	686	GLU	CG-CD	5.51	1.60	1.51
1	A	410	GLU	CD-OE2	5.50	1.31	1.25
1	B	434	TYR	CG-CD1	-5.48	1.32	1.39
1	A	560	GLU	CD-OE2	-5.47	1.19	1.25
1	B	434	TYR	CE1-CZ	-5.43	1.31	1.38
1	B	730	ARG	CZ-NH2	5.40	1.40	1.33
1	A	502	GLY	C-O	5.36	1.32	1.23
1	A	161	ARG	CD-NE	-5.25	1.37	1.46
1	A	434	TYR	CE2-CZ	-5.24	1.31	1.38
1	A	647	GLN	CG-CD	5.16	1.62	1.51
1	A	153	TRP	CG-CD1	-5.06	1.29	1.36

All (102) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	161	ARG	NE-CZ-NH1	21.29	130.94	120.30
1	A	161	ARG	NE-CZ-NH2	-20.84	109.88	120.30
1	B	161	ARG	NE-CZ-NH2	-17.95	111.33	120.30
1	A	161	ARG	NE-CZ-NH1	16.92	128.76	120.30
1	B	382	ARG	NE-CZ-NH1	10.54	125.57	120.30
1	A	426[A]	ARG	NE-CZ-NH2	-9.15	115.72	120.30
1	B	506	ARG	NE-CZ-NH1	8.97	124.79	120.30
1	A	82	ARG	NE-CZ-NH1	8.83	124.72	120.30
1	A	263	ARG	NE-CZ-NH2	-8.81	115.89	120.30
1	A	376	ASP	CB-CG-OD2	-8.74	110.44	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	132	ARG	NE-CZ-NH2	-8.71	115.94	120.30
1	A	433	ARG	NE-CZ-NH1	8.62	124.61	120.30
1	B	178	GLU	OE1-CD-OE2	8.42	133.41	123.30
1	B	633	VAL	CG1-CB-CG2	-8.36	97.53	110.90
1	B	433	ARG	NE-CZ-NH1	8.28	124.44	120.30
1	A	587	ASP	CB-CG-OD1	8.26	125.73	118.30
1	A	223	ARG	NE-CZ-NH2	8.24	124.42	120.30
1	B	268	ASP	CB-CG-OD1	8.19	125.67	118.30
1	A	60	ASP	CB-CG-OD2	-8.06	111.05	118.30
1	A	48	ASP	CB-CG-OD1	8.04	125.53	118.30
1	B	82	ARG	NE-CZ-NH1	8.00	124.30	120.30
1	B	48	ASP	CB-CG-OD1	7.99	125.49	118.30
1	B	393	ARG	NE-CZ-NH2	-7.85	116.37	120.30
1	B	263	ARG	NE-CZ-NH1	7.84	124.22	120.30
1	B	713	ARG	NE-CZ-NH2	-7.69	116.45	120.30
1	A	532	ARG	NE-CZ-NH2	-7.61	116.49	120.30
1	A	198	GLU	OE1-CD-OE2	-7.57	114.21	123.30
1	A	54	ARG	NE-CZ-NH1	7.45	124.03	120.30
1	A	701	ARG	NE-CZ-NH1	7.42	124.01	120.30
1	A	60	ASP	CB-CG-OD1	7.37	124.94	118.30
1	B	40	ARG	NE-CZ-NH1	7.35	123.98	120.30
1	A	218	ARG	NE-CZ-NH2	-7.29	116.66	120.30
1	B	249	ASP	CB-CG-OD1	7.21	124.79	118.30
1	B	688	ARG	NE-CZ-NH1	7.19	123.90	120.30
1	A	649	ARG	NE-CZ-NH1	7.15	123.87	120.30
1	B	382	ARG	NE-CZ-NH2	-7.06	116.77	120.30
1	B	704	LEU	CB-CG-CD2	7.06	123.01	111.00
1	B	713	ARG	NE-CZ-NH1	6.97	123.78	120.30
1	A	198	GLU	CG-CD-OE1	6.96	132.22	118.30
1	B	198	GLU	OE1-CD-OE2	-6.90	115.02	123.30
1	B	649	ARG	NE-CZ-NH2	-6.89	116.85	120.30
1	B	149	ARG	NE-CZ-NH2	-6.87	116.87	120.30
1	B	76	ASP	CB-CG-OD2	-6.84	112.14	118.30
1	A	40	ARG	NE-CZ-NH1	6.83	123.71	120.30
1	A	602	ARG	NE-CZ-NH2	-6.76	116.92	120.30
1	A	395	ASP	CB-CG-OD1	6.75	124.37	118.30
1	B	649	ARG	NE-CZ-NH1	6.74	123.67	120.30
1	A	161	ARG	CD-NE-CZ	6.74	133.04	123.60
1	A	76	ASP	CB-CG-OD2	-6.68	112.29	118.30
1	A	649	ARG	NE-CZ-NH2	-6.60	117.00	120.30
1	A	434	TYR	CB-CG-CD1	6.59	124.96	121.00
1	B	83	ASP	CB-CG-OD2	6.48	124.13	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	459	ASP	CB-CG-OD1	6.44	124.10	118.30
1	B	688	ARG	NE-CZ-NH2	-6.37	117.12	120.30
1	B	529	GLU	OE1-CD-OE2	-6.33	115.70	123.30
1	B	268	ASP	CB-CG-OD2	-6.31	112.62	118.30
1	A	268	ASP	CB-CG-OD2	-6.30	112.63	118.30
1	B	198	GLU	CG-CD-OE1	6.28	130.85	118.30
1	B	587	ASP	CB-CG-OD1	6.27	123.94	118.30
1	B	730	ARG	NE-CZ-NH2	6.23	123.42	120.30
1	A	670	ASP	CB-CG-OD2	-6.22	112.70	118.30
1	A	93	ASP	CB-CG-OD1	6.08	123.77	118.30
1	B	218	ARG	NE-CZ-NH2	-5.97	117.32	120.30
1	B	428	MET	CA-CB-CG	-5.89	103.28	113.30
1	B	434	TYR	CB-CG-CD1	5.87	124.52	121.00
1	A	713	ARG	NE-CZ-NH2	-5.83	117.39	120.30
1	B	459	ASP	CB-CG-OD1	5.81	123.53	118.30
1	B	255	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	B	310	LYS	CB-CG-CD	-5.77	96.60	111.60
1	B	60	ASP	CB-CG-OD2	-5.76	113.12	118.30
1	B	670	ASP	CB-CG-OD2	-5.75	113.12	118.30
1	A	255	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	B	532	ARG	NE-CZ-NH2	-5.67	117.47	120.30
1	B	121	ASP	CB-CG-OD2	-5.65	113.22	118.30
1	A	504	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	B	54	ARG	NE-CZ-NH1	5.55	123.07	120.30
1	B	225	LEU	CB-CG-CD2	-5.54	101.59	111.00
1	B	121	ASP	CB-CG-OD1	5.51	123.26	118.30
1	A	610	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	B	549	ASP	CB-CG-OD1	5.45	123.21	118.30
1	B	580	ASP	CB-CG-OD1	-5.43	113.41	118.30
1	A	268	ASP	CB-CG-OD1	5.43	123.19	118.30
1	A	713	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	B	540	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	B	223	ARG	NE-CZ-NH1	-5.35	117.62	120.30
1	B	653	PHE	CB-CG-CD1	-5.35	117.06	120.80
1	A	459	ASP	CB-CG-OD2	-5.29	113.54	118.30
1	A	540	ARG	NE-CZ-NH1	5.28	122.94	120.30
1	B	653	PHE	CB-CG-CD2	5.26	124.48	120.80
1	A	40	ARG	NE-CZ-NH2	-5.26	117.67	120.30
1	A	121	ASP	CB-CG-OD1	5.25	123.02	118.30
1	A	506	ARG	NE-CZ-NH1	5.21	122.90	120.30
1	B	683	ASP	CB-CG-OD1	5.17	122.96	118.30
1	B	149	ARG	NE-CZ-NH1	5.17	122.89	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	108	ARG	NE-CZ-NH1	5.17	122.89	120.30
1	B	76	ASP	CB-CG-OD1	5.17	122.95	118.30
1	B	683	ASP	CB-CG-OD2	-5.14	113.67	118.30
1	A	683	ASP	CB-CG-OD1	5.09	122.88	118.30
1	B	587	ASP	CB-CG-OD2	-5.09	113.72	118.30
1	B	701	ARG	NE-CZ-NH1	5.07	122.84	120.30
1	B	249	ASP	CB-CG-OD2	-5.07	113.74	118.30
1	A	255	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	161	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5534	0	5366	15	0
1	B	5518	0	5337	21	0
2	A	43	0	30	0	0
2	B	43	0	30	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	0	1	0
4	B	2	0	0	0	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	16	0	28	5	0
7	B	8	0	14	2	0
8	A	725	0	0	8	1
8	B	776	0	0	10	1
All	All	12683	0	10805	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119[B]:THR:HG22	1:B:593:VAL:HG21	1.54	0.90
1:A:255:ARG:NH2	8:A:901:HOH:O	2.06	0.88
1:A:119[B]:THR:HG21	8:A:1061:HOH:O	1.77	0.83
1:B:512:ASP:OD1	8:B:901:HOH:O	2.05	0.74
1:A:119[B]:THR:HG23	1:A:593:VAL:HG11	1.73	0.71
1:A:593:VAL:HG13	8:A:1011:HOH:O	1.92	0.70
7:A:807:MPD:O4	7:A:807:MPD:CM	2.43	0.67
7:B:806:MPD:O4	7:B:806:MPD:H12	1.94	0.66
7:A:808:MPD:H11	7:A:808:MPD:H53	1.77	0.65
1:B:119[B]:THR:HG23	8:B:1092:HOH:O	1.94	0.65
7:B:806:MPD:O4	7:B:806:MPD:C1	2.46	0.64
1:B:696:LYS:HE2	8:B:922:HOH:O	2.02	0.60
1:B:62:MET:HE3	8:B:1587:HOH:O	2.04	0.57
4:A:803:OXY:O2	8:A:902:HOH:O	2.18	0.57
1:B:711[A]:GLN:NE2	8:B:907:HOH:O	2.40	0.55
1:B:540:ARG:CZ	1:B:540:ARG:HA	2.38	0.54
1:A:161:ARG:NH2	8:A:906:HOH:O	2.36	0.54
1:B:403[A]:ARG:NH1	8:B:913:HOH:O	2.42	0.53
1:A:111:TRP:HZ3	1:A:238:TYR:HH	1.52	0.52
1:A:504:ARG:HD2	8:A:976:HOH:O	2.09	0.52
1:A:696:LYS:HE2	8:A:1183:HOH:O	2.11	0.51
1:B:255:ARG:HD2	8:B:1345:HOH:O	2.11	0.50
1:A:633[A]:VAL:CG2	1:A:719:TYR:CZ	2.95	0.49
1:A:119[B]:THR:CG2	1:A:593:VAL:HG11	2.42	0.48
1:B:662:ASN:H	1:B:725:GLN:HE22	1.59	0.48
1:B:725:GLN:NE2	8:B:923:HOH:O	2.48	0.47
1:A:313:TYR:CZ	1:A:314:ARG:HD3	2.50	0.47
1:B:313:TYR:CZ	1:B:314:ARG:HD3	2.51	0.46
1:B:633:VAL:CG2	1:B:719:TYR:CZ	2.99	0.46
1:B:431:ARG:HD2	1:B:447:GLN:OE1	2.16	0.45
7:A:808:MPD:HM2	8:A:1126:HOH:O	2.16	0.45
7:A:807:MPD:HM1	7:A:807:MPD:O4	2.18	0.43
1:B:696:LYS:CE	8:B:922:HOH:O	2.66	0.42
1:B:255:ARG:HB2	8:B:1586:HOH:O	2.20	0.42
1:A:726:GLU:HG2	1:A:730:ARG:NH1	2.35	0.42
7:A:808:MPD:C1	7:A:808:MPD:H53	2.48	0.42
1:B:540:ARG:NH1	1:B:540:ARG:HA	2.35	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:602:ARG:HB3	1:A:645:VAL:HG23	2.01	0.41
1:A:313:TYR:CE2	1:A:314:ARG:HD3	2.55	0.41
1:B:69:ALA:O	1:B:73:GLU:HG2	2.20	0.41
1:B:49:LEU:HD12	1:B:711[B]:GLN:CD	2.41	0.41
1:B:269:GLU:OE1	1:B:404:ARG:NH2	2.48	0.41
1:B:428:MET:O	1:B:433:ARG:HD3	2.21	0.41
1:A:242:GLU:O	1:A:248:PRO:HA	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:1470:HOH:O	8:B:1563:HOH:O[4_445]	1.97	0.23

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	717/728 (98%)	706 (98%)	11 (2%)	0	100	100
1	B	714/728 (98%)	704 (99%)	10 (1%)	0	100	100
All	All	1431/1456 (98%)	1410 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	556/561 (99%)	548 (99%)	8 (1%)	74	71
1	B	553/561 (99%)	548 (99%)	5 (1%)	84	83
All	All	1109/1122 (99%)	1096 (99%)	13 (1%)	78	76

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	ARG
1	A	118	ARG
1	A	141	ASP
1	A	540	ARG
1	A	608	LYS
1	A	610	ARG
1	A	649	ARG
1	A	726	GLU
1	B	40	ARG
1	B	118	ARG
1	B	141	ASP
1	B	608	LYS
1	B	649	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	HIS
1	A	70	GLN
1	A	85	HIS
1	A	725	GLN
1	B	46	GLN
1	B	55	HIS
1	B	544	GLN
1	B	725	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 4 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	801	1,6	24,50,50	1.21	3 (12%)	16,82,82	1.92	4 (25%)
4	OXY	A	803	-	1,1,1	0.05	0	0,0,0	0.00	-
4	OXY	A	804	-	1,1,1	0.07	0	0,0,0	0.00	-
5	PO4	A	805	-	4,4,4	0.23	0	6,6,6	0.29	0
7	MPD	A	807	-	6,7,7	1.53	2 (33%)	6,10,10	2.04	3 (50%)
7	MPD	A	808	-	6,7,7	0.75	0	6,10,10	0.76	0
2	HEM	B	801	1,6	24,50,50	1.38	3 (12%)	16,82,82	2.86	7 (43%)
4	OXY	B	804	-	1,1,1	0.15	0	0,0,0	0.00	-
5	PO4	B	805	-	4,4,4	0.54	0	6,6,6	0.30	0
7	MPD	B	806	-	6,7,7	0.51	0	6,10,10	1.61	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	801	1,6	-	0/6/54/54	0/0/8/8
4	OXY	A	803	-	-	0/0/0/0	0/0/0/0
4	OXY	A	804	-	-	0/0/0/0	0/0/0/0
5	PO4	A	805	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MPD	A	807	-	-	0/5/5/5	0/0/0/0
7	MPD	A	808	-	-	0/5/5/5	0/0/0/0
2	HEM	B	801	1,6	-	0/6/54/54	0/0/8/8
4	OXY	B	804	-	-	0/0/0/0	0/0/0/0
5	PO4	B	805	-	-	0/0/0/0	0/0/0/0
7	MPD	B	806	-	-	0/5/5/5	0/0/0/0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	HEM	CAD-C3D	-3.10	1.48	1.52
2	B	801	HEM	C1B-NB	-2.89	1.32	1.36
7	A	807	MPD	O2-C2	-2.54	1.37	1.44
2	B	801	HEM	CMD-C2D	-2.15	1.47	1.51
2	A	801	HEM	CMB-C2B	2.04	1.56	1.51
2	A	801	HEM	C3C-C2C	2.19	1.43	1.40
7	A	807	MPD	C5-C4	2.31	1.62	1.51
2	A	801	HEM	C4C-NC	2.83	1.40	1.36

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	HEM	CMA-C3A-C4A	-5.95	118.19	128.31
2	A	801	HEM	CAA-CBA-CGA	-4.49	104.04	112.78
2	B	801	HEM	CAA-CBA-CGA	-4.48	104.06	112.78
2	B	801	HEM	C3B-CAB-CBB	-4.20	117.96	126.40
2	A	801	HEM	C3B-CAB-CBB	-3.69	118.97	126.40
2	A	801	HEM	CMB-C2B-C3B	2.01	129.02	125.09
7	B	806	MPD	O2-C2-CM	2.14	115.88	108.01
2	B	801	HEM	CMB-C2B-C3B	2.21	129.41	125.09
7	B	806	MPD	C1-C2-C3	2.23	121.92	109.98
7	A	807	MPD	O4-C4-C5	2.39	121.39	109.47
7	A	807	MPD	CM-C2-C1	2.40	116.14	110.41
2	B	801	HEM	CAD-CBD-CGD	2.48	117.61	112.78
7	A	807	MPD	CM-C2-C3	2.63	124.02	109.98
2	A	801	HEM	CBD-CAD-C3D	2.88	117.52	112.47
2	B	801	HEM	C3B-C4B-NB	3.38	113.58	109.21
2	B	801	HEM	CMA-C3A-C2A	4.64	134.95	125.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	803	OXY	1	0
7	A	807	MPD	2	0
7	A	808	MPD	3	0
7	B	806	MPD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	713/728 (97%)	-0.37	16 (2%) 65 68	11, 18, 37, 68	0
1	B	713/728 (97%)	-0.46	14 (1%) 68 71	11, 17, 35, 70	0
All	All	1426/1456 (97%)	-0.42	30 (2%) 67 70	11, 18, 36, 70	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	748	ALA	6.4
1	A	748	ALA	6.0
1	A	541	GLY	5.1
1	B	679	ALA	4.6
1	B	65	ASP	4.5
1	A	64	LYS	4.1
1	B	64	LYS	4.0
1	A	610	ARG	4.0
1	B	610	ARG	3.8
1	A	540	ARG	3.7
1	B	680	ALA	3.5
1	A	65	ASP	3.5
1	A	608	LYS	3.3
1	B	44	PRO	3.2
1	B	608	LYS	3.2
1	B	540	ARG	3.2
1	A	63	GLY	2.8
1	A	647	GLN	2.8
1	A	679	ALA	2.7
1	A	378	SER	2.7
1	B	541	GLY	2.6
1	A	454	ASP	2.5
1	B	691	ALA	2.4
1	A	222	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	215	PRO	2.3
1	A	691	ALA	2.2
1	B	378	SER	2.2
1	B	55	HIS	2.2
1	A	680	ALA	2.1
1	B	45	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	MPD	A	807	8/8	0.82	0.21	9.91	44,47,66,67	0
7	MPD	B	806	8/8	0.91	0.14	3.17	45,50,55,56	0
3	NA	B	802	1/1	0.99	0.09	1.77	14,14,14,14	0
3	NA	A	802	1/1	0.99	0.06	-0.69	13,13,13,13	0
2	HEM	A	801	43/43	0.99	0.06	-1.06	12,14,16,17	0
2	HEM	B	801	43/43	0.99	0.06	-1.11	11,12,13,14	0
5	PO4	A	805	5/5	0.95	0.20	-	44,56,58,63	0
7	MPD	A	808	8/8	0.85	0.18	-	37,39,47,49	0
6	O	A	806	1/1	0.99	0.19	-	24,24,24,24	0
4	OXY	B	804	2/2	0.95	0.22	-	35,35,35,36	0
4	OXY	A	804	2/2	0.96	0.17	-	32,32,32,42	0
6	O	B	803	1/1	0.98	0.08	-	19,19,19,19	0
5	PO4	B	805	5/5	0.92	0.26	-	53,54,67,70	0
4	OXY	A	803	2/2	0.98	0.24	-	28,28,28,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.