



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 10:21 PM GMT

PDB ID : 1SX4
Title : GroEL-GroES-ADP7
Authors : Chaudhry, C.; Horwich, A.L.; Brunger, A.T.; Adams, P.D.
Deposited on : 2004-03-30
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	524	<div> <div>3%</div> <div>53%</div> <div>38%</div> <div>8%</div> </div>
1	B	524	<div> <div>5%</div> <div>56%</div> <div>35%</div> <div>8%</div> </div>
1	C	524	<div> <div>4%</div> <div>55%</div> <div>35%</div> <div>9%</div> </div>
1	D	524	<div> <div>3%</div> <div>56%</div> <div>37%</div> <div>6%</div> </div>
1	E	524	<div> <div>5%</div> <div>55%</div> <div>37%</div> <div>7%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	524	
1	G	524	
1	H	524	
1	I	524	
1	J	524	
1	K	524	
1	L	524	
1	M	524	
1	N	524	
2	O	97	
2	P	97	
2	Q	97	
2	R	97	
2	S	97	
2	T	97	
2	U	97	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 59457 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called groEL protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	B	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	C	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	D	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	E	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	F	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	G	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	H	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	I	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	J	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	K	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	L	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	M	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			
1	N	524	Total	C	N	O	S	0	0	0
			3856	2397	665	774	20			

- Molecule 2 is a protein called groES protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	O	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	P	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	Q	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	R	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	S	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	T	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	U	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		
3	E	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		
3	F	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	E	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	F	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	G	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	12	Total	O	0	0
			12	12		
5	B	10	Total	O	0	0
			10	10		
5	C	11	Total	O	0	0
			11	11		
5	D	11	Total	O	0	0
			11	11		
5	E	10	Total	O	0	0
			10	10		

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	F	8	Total 8	O 8	0	0
5	G	13	Total 13	O 13	0	0
5	H	12	Total 12	O 12	0	0
5	I	19	Total 19	O 19	0	0
5	J	14	Total 14	O 14	0	0
5	K	14	Total 14	O 14	0	0
5	L	16	Total 16	O 16	0	0
5	M	10	Total 10	O 10	0	0
5	N	21	Total 21	O 21	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: groEL protein







• Molecule 1: groEL protein




• Molecule 1: groEL protein







• Molecule 1: groEL protein




• Molecule 1: groEL protein







• Molecule 1: groEL protein




• Molecule 1: groEL protein







• Molecule 1: groEL protein



• Molecule 1: groEL protein






- Molecule 1: groEL protein




- Molecule 1: groEL protein







• Molecule 1: groEL protein




• Molecule 1: groEL protein







• Molecule 2: groES protein




• Molecule 2: groES protein




• Molecule 2: groES protein




• Molecule 2: groES protein




- Molecule 2: groES protein



- Molecule 2: groES protein



- Molecule 2: groES protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	255.26Å 265.25Å 184.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 3.00 40.07 – 2.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-3.00) 96.7 (40.07-2.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.258 , 0.287 0.246 , 0.273	Depositor DCC
R_{free} test set	12046 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	50.3	Xtriage
Anisotropy	0.614	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 56.0	EDS
Estimated twinning fraction	0.003 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 242645 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	59457	wwPDB-VP
Average B, all atoms (Å ²)	2.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.26	27/3884 (0.7%)	1.17	27/5243 (0.5%)
1	B	1.09	12/3884 (0.3%)	1.10	20/5243 (0.4%)
1	C	1.20	17/3884 (0.4%)	1.20	31/5243 (0.6%)
1	D	1.22	19/3884 (0.5%)	1.14	15/5243 (0.3%)
1	E	1.01	11/3884 (0.3%)	1.07	22/5243 (0.4%)
1	F	0.98	12/3884 (0.3%)	1.06	21/5243 (0.4%)
1	G	1.16	18/3884 (0.5%)	1.16	28/5243 (0.5%)
1	H	1.06	7/3884 (0.2%)	1.11	25/5243 (0.5%)
1	I	1.18	10/3884 (0.3%)	1.16	17/5243 (0.3%)
1	J	1.07	9/3884 (0.2%)	1.15	26/5243 (0.5%)
1	K	0.95	4/3884 (0.1%)	1.06	21/5243 (0.4%)
1	L	0.87	1/3884 (0.0%)	1.03	20/5243 (0.4%)
1	M	1.08	16/3884 (0.4%)	1.13	19/5243 (0.4%)
1	N	1.09	13/3884 (0.3%)	1.13	26/5243 (0.5%)
2	O	0.65	0/732	0.91	3/983 (0.3%)
2	P	0.61	0/732	0.89	3/983 (0.3%)
2	Q	0.66	0/732	0.92	3/983 (0.3%)
2	R	0.73	0/732	0.92	3/983 (0.3%)
2	S	0.67	0/732	0.92	3/983 (0.3%)
2	T	0.65	0/732	0.91	2/983 (0.2%)
2	U	0.65	0/732	0.90	5/983 (0.5%)
All	All	1.06	176/59500 (0.3%)	1.10	340/80283 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	I	0	2
1	J	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	M	0	1
All	All	0	4

The worst 5 of 176 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	130	GLU	CD-OE1	9.80	1.36	1.25
1	I	461	GLU	CD-OE1	9.36	1.35	1.25
1	N	483	GLU	CD-OE2	8.57	1.35	1.25
1	I	484	GLU	CD-OE1	8.36	1.34	1.25
1	D	71	ALA	CA-CB	-8.26	1.35	1.52

The worst 5 of 340 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	N	83	ASP	CB-CG-OD2	10.21	127.49	118.30
1	G	13	ARG	NE-CZ-NH2	-10.02	115.29	120.30
1	D	421	ARG	NE-CZ-NH1	9.99	125.29	120.30
1	J	428	ASP	CB-CG-OD2	9.67	127.00	118.30
1	C	452	ARG	NE-CZ-NH1	9.51	125.05	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	I	32	GLY	Peptide
1	I	415	GLY	Peptide
1	J	415	GLY	Peptide
1	M	415	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3856	0	3976	190	0
1	B	3856	0	3976	188	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3856	0	3976	191	0
1	D	3856	0	3976	180	0
1	E	3856	0	3976	172	0
1	F	3856	0	3976	168	0
1	G	3856	0	3976	185	0
1	H	3856	0	3976	168	0
1	I	3856	0	3976	175	0
1	J	3856	0	3976	162	0
1	K	3856	0	3976	171	0
1	L	3856	0	3976	184	1
1	M	3856	0	3976	190	0
1	N	3856	0	3976	173	0
2	O	728	0	762	24	0
2	P	728	0	762	19	0
2	Q	728	0	762	20	0
2	R	728	0	762	15	0
2	S	728	0	762	17	0
2	T	728	0	762	16	0
2	U	728	0	762	16	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
4	A	27	0	12	1	0
4	B	27	0	12	2	0
4	C	27	0	12	1	0
4	D	27	0	12	1	0
4	E	27	0	12	1	0
4	F	27	0	12	0	0
4	G	27	0	12	1	0
5	A	12	0	0	1	0
5	B	10	0	0	6	0
5	C	11	0	0	1	0
5	D	11	0	0	2	0
5	E	10	0	0	2	0
5	F	8	0	0	1	0
5	G	13	0	0	2	0
5	H	12	0	0	7	0
5	I	19	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	J	14	0	0	4	0
5	K	14	0	0	2	0
5	L	16	0	0	3	0
5	M	10	0	0	1	0
5	N	21	0	0	8	0
All	All	59457	0	61082	2532	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 2532 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:73:MET:SD	1:I:73:MET:CG	2.01	1.48
1:A:73:MET:CE	1:A:73:MET:SD	2.01	1.46
1:J:288:MET:SD	1:J:288:MET:CE	2.02	1.46
1:I:114:MET:CE	1:I:114:MET:SD	2.02	1.46
1:A:16:MET:SD	1:A:16:MET:CE	2.02	1.45

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:354:GLU:OE1	1:L:354:GLU:OE1[2_655]	1.71	0.49

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	522/524 (100%)	478 (92%)	40 (8%)	4 (1%)	24	66
1	B	522/524 (100%)	477 (91%)	39 (8%)	6 (1%)	17	58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	522/524 (100%)	475 (91%)	41 (8%)	6 (1%)	17	58
1	D	522/524 (100%)	475 (91%)	41 (8%)	6 (1%)	17	58
1	E	522/524 (100%)	477 (91%)	40 (8%)	5 (1%)	19	61
1	F	522/524 (100%)	478 (92%)	38 (7%)	6 (1%)	17	58
1	G	522/524 (100%)	473 (91%)	44 (8%)	5 (1%)	19	61
1	H	522/524 (100%)	473 (91%)	39 (8%)	10 (2%)	10	43
1	I	522/524 (100%)	474 (91%)	35 (7%)	13 (2%)	7	34
1	J	522/524 (100%)	473 (91%)	41 (8%)	8 (2%)	13	50
1	K	522/524 (100%)	471 (90%)	42 (8%)	9 (2%)	11	46
1	L	522/524 (100%)	476 (91%)	36 (7%)	10 (2%)	10	43
1	M	522/524 (100%)	469 (90%)	42 (8%)	11 (2%)	9	40
1	N	522/524 (100%)	470 (90%)	40 (8%)	12 (2%)	8	36
2	O	95/97 (98%)	75 (79%)	12 (13%)	8 (8%)	1	5
2	P	95/97 (98%)	75 (79%)	12 (13%)	8 (8%)	1	5
2	Q	95/97 (98%)	75 (79%)	12 (13%)	8 (8%)	1	5
2	R	95/97 (98%)	74 (78%)	13 (14%)	8 (8%)	1	5
2	S	95/97 (98%)	75 (79%)	12 (13%)	8 (8%)	1	5
2	T	95/97 (98%)	73 (77%)	14 (15%)	8 (8%)	1	5
2	U	95/97 (98%)	73 (77%)	14 (15%)	8 (8%)	1	5
All	All	7973/8015 (100%)	7159 (90%)	647 (8%)	167 (2%)	9	40

5 of 167 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	PHE
1	B	44	PHE
1	B	483	GLU
1	C	9	GLY
1	C	44	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/404 (100%)	345 (85%)	59 (15%)	4	18
1	B	404/404 (100%)	345 (85%)	59 (15%)	4	18
1	C	404/404 (100%)	345 (85%)	59 (15%)	4	18
1	D	404/404 (100%)	346 (86%)	58 (14%)	4	18
1	E	404/404 (100%)	346 (86%)	58 (14%)	4	18
1	F	404/404 (100%)	345 (85%)	59 (15%)	4	18
1	G	404/404 (100%)	344 (85%)	60 (15%)	4	17
1	H	404/404 (100%)	345 (85%)	59 (15%)	4	18
1	I	404/404 (100%)	342 (85%)	62 (15%)	3	16
1	J	404/404 (100%)	344 (85%)	60 (15%)	4	17
1	K	404/404 (100%)	345 (85%)	59 (15%)	4	18
1	L	404/404 (100%)	341 (84%)	63 (16%)	3	16
1	M	404/404 (100%)	345 (85%)	59 (15%)	4	18
1	N	404/404 (100%)	342 (85%)	62 (15%)	3	16
2	O	80/80 (100%)	65 (81%)	15 (19%)	2	10
2	P	80/80 (100%)	64 (80%)	16 (20%)	1	8
2	Q	80/80 (100%)	64 (80%)	16 (20%)	1	8
2	R	80/80 (100%)	64 (80%)	16 (20%)	1	8
2	S	80/80 (100%)	65 (81%)	15 (19%)	2	10
2	T	80/80 (100%)	63 (79%)	17 (21%)	1	6
2	U	80/80 (100%)	66 (82%)	14 (18%)	2	12
All	All	6216/6216 (100%)	5271 (85%)	945 (15%)	3	16

5 of 945 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	141	SER
1	I	483	GLU
2	Q	30	SER
1	H	311	LYS
1	I	79	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 174 such

sidechains are listed below:

Mol	Chain	Res	Type
1	G	153	ASN
1	H	436	GLN
2	O	68	ASN
1	G	326	ASN
1	H	21	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 7 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ADP	A	600	3	22,29,29	1.50	5 (22%)	27,45,45	3.57	12 (44%)
4	ADP	B	600	3	22,29,29	1.50	5 (22%)	27,45,45	3.23	12 (44%)
4	ADP	C	600	3	22,29,29	1.31	4 (18%)	27,45,45	3.37	13 (48%)
4	ADP	D	600	3	22,29,29	1.66	4 (18%)	27,45,45	3.69	12 (44%)
4	ADP	E	600	3	22,29,29	1.31	4 (18%)	27,45,45	3.09	9 (33%)
4	ADP	F	600	3	22,29,29	1.11	2 (9%)	27,45,45	3.05	10 (37%)
4	ADP	G	600	3	22,29,29	1.41	3 (13%)	27,45,45	2.90	13 (48%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	A	600	3	-	0/12/32/32	0/3/3/3
4	ADP	B	600	3	-	0/12/32/32	0/3/3/3
4	ADP	C	600	3	-	0/12/32/32	0/3/3/3
4	ADP	D	600	3	-	0/12/32/32	0/3/3/3
4	ADP	E	600	3	-	0/12/32/32	0/3/3/3
4	ADP	F	600	3	-	0/12/32/32	0/3/3/3
4	ADP	G	600	3	-	0/12/32/32	0/3/3/3

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	600	ADP	O4'-C1'	-4.30	1.35	1.41
4	A	600	ADP	O4'-C4'	-3.16	1.37	1.45
4	D	600	ADP	O4'-C4'	-3.13	1.37	1.45
4	E	600	ADP	O4'-C1'	-3.06	1.37	1.41
4	G	600	ADP	O4'-C4'	-3.03	1.38	1.45

The worst 5 of 81 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	600	ADP	N3-C2-N1	-14.68	117.66	128.89
4	A	600	ADP	N3-C2-N1	-12.51	119.32	128.89
4	F	600	ADP	N3-C2-N1	-11.41	120.16	128.89
4	C	600	ADP	N3-C2-N1	-11.15	120.36	128.89
4	B	600	ADP	N3-C2-N1	-10.98	120.49	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	600	ADP	1	0
4	B	600	ADP	2	0
4	C	600	ADP	1	0
4	D	600	ADP	1	0
4	E	600	ADP	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	600	ADP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	524/524 (100%)	-0.02	18 (3%)	49	21	2, 2, 3, 5	0
1	B	524/524 (100%)	0.11	24 (4%)	36	14	2, 2, 3, 5	0
1	C	524/524 (100%)	-0.05	20 (3%)	44	18	2, 2, 3, 5	0
1	D	524/524 (100%)	-0.09	16 (3%)	52	24	2, 2, 3, 5	0
1	E	524/524 (100%)	0.16	25 (4%)	34	14	2, 2, 3, 5	0
1	F	524/524 (100%)	0.18	25 (4%)	34	14	2, 2, 3, 5	0
1	G	524/524 (100%)	0.12	34 (6%)	22	8	2, 2, 3, 5	0
1	H	524/524 (100%)	-0.17	4 (0%)	87	67	2, 2, 3, 5	0
1	I	524/524 (100%)	-0.24	3 (0%)	90	73	2, 2, 3, 5	0
1	J	524/524 (100%)	-0.17	2 (0%)	93	80	2, 2, 3, 5	0
1	K	524/524 (100%)	0.01	11 (2%)	67	36	2, 2, 3, 5	0
1	L	524/524 (100%)	0.02	10 (1%)	70	41	2, 2, 3, 5	0
1	M	524/524 (100%)	-0.09	9 (1%)	73	45	2, 2, 3, 5	0
1	N	524/524 (100%)	-0.06	7 (1%)	79	53	2, 2, 3, 5	0
2	O	97/97 (100%)	0.91	18 (18%)	2	1	2, 2, 2, 3	0
2	P	97/97 (100%)	0.79	16 (16%)	2	1	2, 2, 2, 3	0
2	Q	97/97 (100%)	0.61	11 (11%)	7	2	2, 2, 2, 3	0
2	R	97/97 (100%)	0.76	15 (15%)	3	1	2, 2, 2, 3	0
2	S	97/97 (100%)	0.76	15 (15%)	3	1	2, 2, 2, 3	0
2	T	97/97 (100%)	0.77	18 (18%)	2	1	2, 2, 2, 3	0
2	U	97/97 (100%)	0.74	12 (12%)	5	2	2, 2, 2, 3	0
All	All	8015/8015 (100%)	0.05	313 (3%)	43	18	2, 2, 3, 5	0

The worst 5 of 313 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	208	PRO	10.6
1	C	361	ASP	10.4
1	A	361	ASP	10.1
1	E	361	ASP	8.0
1	B	361	ASP	7.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	ADP	D	600	27/27	0.96	0.18	0.62	2,2,4,5	0
4	ADP	A	600	27/27	0.95	0.18	0.55	2,2,3,6	0
4	ADP	B	600	27/27	0.94	0.20	0.52	2,2,4,5	0
4	ADP	F	600	27/27	0.93	0.21	0.20	2,2,3,5	0
4	ADP	E	600	27/27	0.93	0.20	-0.09	2,2,3,5	0
4	ADP	G	600	27/27	0.96	0.17	-0.14	2,2,4,5	0
4	ADP	C	600	27/27	0.97	0.17	-0.49	2,2,4,5	0
3	MG	D	601	1/1	0.97	0.38	-	2,2,2,2	0
3	MG	C	601	1/1	0.90	0.27	-	2,2,2,2	0
3	MG	E	601	1/1	0.96	0.27	-	2,2,2,2	0
3	MG	B	601	1/1	0.92	0.30	-	2,2,2,2	0
3	MG	A	601	1/1	0.94	0.34	-	2,2,2,2	0
3	MG	G	601	1/1	0.97	0.29	-	2,2,2,2	0
3	MG	F	601	1/1	0.97	0.37	-	2,2,2,2	0

6.5 Other polymers [i](#)

There are no such residues in this entry.