



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:05 PM GMT

PDB ID : 3SX3
Title : Crystal structure of AAB+UDP+Gal with glycerol as the cryoprotectant
Authors : Johal, A.R.; Evans, S.V.
Deposited on : 2011-07-14
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1278 (1.48-1.44)
Clashscore	102246	1336 (1.48-1.44)
Ramachandran outliers	100387	1320 (1.48-1.44)
Sidechain outliers	100360	1320 (1.48-1.44)
RSRZ outliers	91569	1279 (1.48-1.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 2708 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Histo-blood group ABO system transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	291	Total	C	N	O	S	0	1	0
			2397	1551	418	414	14			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	62	PHE	-	EXPRESSION TAG	UNP P16442
A	63	MET	-	EXPRESSION TAG	UNP P16442
A	266	MET	LEU	SEE REMARK 999	UNP P16442
A	268	ALA	GLY	SEE REMARK 999	UNP P16442

- Molecule 2 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C₉H₁₄N₂O₁₂P₂).




- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SUGAR (BETA-D-GALACTOSE) (three-letter code: GAL) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	260	Total	O	0	0
			260	260		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histo-blood group ABO system transferase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	52.39 Å 149.12 Å 79.46 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.45 19.86 – 1.45	Depositor EDS
% Data completeness (in resolution range)	96.3 (20.00-1.45) 96.3 (19.86-1.45)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.76 (at 1.45 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.198 , 0.227 0.189 , 0.218	Depositor DCC
R_{free} test set	2704 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 50.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 53445 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2708	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, GOL, UDP, MN, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.38	8/2468 (0.3%)	1.34	24/3348 (0.7%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	135	GLU	CG-CD	8.78	1.65	1.51
1	A	135	GLU	CD-OE2	8.15	1.34	1.25
1	A	135	GLU	CD-OE1	-7.05	1.17	1.25
1	A	206	TYR	CE2-CZ	6.43	1.47	1.38
1	A	126	TYR	CG-CD2	6.09	1.47	1.39
1	A	107	GLU	CD-OE2	-5.94	1.19	1.25
1	A	100	PHE	CE1-CZ	5.04	1.47	1.37
1	A	334	ALA	CA-CB	5.00	1.62	1.52

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	266	MET	CG-SD-CE	-13.90	77.95	100.20
1	A	288	MET	CG-SD-CE	-9.96	84.27	100.20
1	A	211	ASP	CB-CG-OD1	8.87	126.29	118.30
1	A	282	ARG	NE-CZ-NH1	8.72	124.66	120.30
1	A	213	ASP	CB-CG-OD1	8.62	126.06	118.30
1	A	326	ASP	CB-CG-OD2	-8.43	110.72	118.30
1	A	324	LEU	CB-CG-CD2	-8.34	96.81	111.00
1	A	191	MET	CG-SD-CE	-8.21	87.07	100.20
1	A	340	ARG	NE-CZ-NH1	7.58	124.09	120.30
1	A	110	ARG	NE-CZ-NH1	6.75	123.67	120.30
1	A	191	MET	CA-CB-CG	-6.75	101.83	113.30
1	A	340	ARG	NE-CZ-NH2	-6.17	117.22	120.30
1	A	213	ASP	OD1-CG-OD2	-6.14	111.64	123.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	312	ARG	NE-CZ-NH1	-6.10	117.25	120.30
1	A	324	LEU	CB-CG-CD1	5.95	121.11	111.00
1	A	68	ARG	NE-CZ-NH2	5.90	123.25	120.30
1	A	187	ARG	CB-CG-CD	5.84	126.79	111.60
1	A	246	TYR	CB-CG-CD1	-5.43	117.74	121.00
1	A	205	ASP	CB-CG-OD1	-5.42	113.42	118.30
1	A	135	GLU	CG-CD-OE1	-5.41	107.48	118.30
1	A	244	PHE	CB-CG-CD2	5.40	124.58	120.80
1	A	187	ARG	NE-CZ-NH2	5.10	122.85	120.30
1	A	180	ARG	NE-CZ-NH1	5.04	122.82	120.30
1	A	109	PHE	CB-CG-CD1	-5.01	117.29	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2397	0	2386	38	0
2	A	25	0	11	2	0
3	A	6	0	8	0	0
4	A	12	0	12	0	0
5	A	7	0	9	1	0
6	A	1	0	0	0	0
7	A	260	0	0	10	0
All	All	2708	0	2426	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118:LEU:HG	7:A:470:HOH:O	1.15	1.30
1:A:324:LEU:HG	7:A:469:HOH:O	1.10	1.27

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:GLU:HA	1:A:278:GLN:HE22	1.15	1.07
1:A:63:MET:SD	1:A:112:GLN:NE2	2.31	1.02
1:A:135:GLU:OE1	7:A:547:HOH:O	1.88	0.91
1:A:75:LYS:HB2	1:A:78:THR:OG1	1.84	0.77
1:A:348:HIS:HE1	2:A:1:UDP:O1B	1.67	0.77
1:A:197:GLU:HA	1:A:278:GLN:NE2	1.99	0.74
1:A:193:SER:C	1:A:197:GLU:HG2	2.16	0.66
1:A:286:GLN:HA	1:A:289:MET:HE2	1.78	0.65
1:A:197:GLU:CA	1:A:278:GLN:HE22	2.02	0.63
1:A:348:HIS:CE1	2:A:1:UDP:O1B	2.51	0.63
1:A:101:ASN:HD22	1:A:101:ASN:C	2.02	0.63
1:A:101:ASN:HD21	1:A:103:ASP:HB2	1.65	0.62
1:A:193:SER:O	1:A:197:GLU:HG2	2.03	0.59
1:A:193:SER:C	1:A:197:GLU:OE1	2.42	0.57
1:A:62:PHE:N	7:A:499:HOH:O	2.37	0.56
1:A:266:MET:CB	1:A:324:LEU:HD23	2.37	0.55
1:A:183:ASP:O	1:A:187:ARG:HG2	2.05	0.55
1:A:106:ASN:OD1	1:A:145:HIS:HE1	1.90	0.54
1:A:217:ARG:NH2	7:A:462:HOH:O	1.92	0.54
1:A:305:HIS:HE1	7:A:15:HOH:O	1.90	0.54
1:A:266:MET:HG3	1:A:324:LEU:HD23	1.90	0.53
1:A:110:ARG:HH22	1:A:113:ASN:HD22	1.57	0.53
1:A:79:PRO:HB3	7:A:554:HOH:O	2.11	0.51
1:A:242:GLU:CD	1:A:242:GLU:H	2.12	0.50
1:A:145:HIS:HD2	7:A:38:HOH:O	1.94	0.49
1:A:305:HIS:HD2	7:A:52:HOH:O	1.97	0.47
1:A:110:ARG:HH22	1:A:113:ASN:ND2	2.14	0.45
1:A:135:GLU:CG	7:A:547:HOH:O	2.64	0.45
1:A:266:MET:CG	1:A:324:LEU:HD23	2.47	0.45
1:A:257:PRO:HG2	1:A:260:GLU:OE2	2.17	0.44
1:A:196:CYS:SG	1:A:196:CYS:O	2.76	0.43
1:A:266:MET:HB3	1:A:324:LEU:CD2	2.49	0.42
1:A:205:ASP:CG	5:A:4:PEG:H11	2.40	0.41
1:A:261:GLY:HA2	1:A:311:LEU:HD13	2.02	0.40
1:A:266:MET:CB	1:A:324:LEU:CD2	2.99	0.40
1:A:101:ASN:C	1:A:101:ASN:ND2	2.72	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288 / 293 (98%)	282 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260 / 261 (100%)	253 (97%)	7 (3%)	52	15

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	ASN
1	A	113	ASN
1	A	266	MET
1	A	269	PHE
1	A	302	ASP
1	A	329	LEU
1	A	348	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	101	ASN
1	A	113	ASN
1	A	145	HIS
1	A	278	GLN
1	A	286	GLN
1	A	294	ASN
1	A	305	HIS
1	A	348	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	UDP	A	1	6	18,26,26	1.91	4 (22%)	26,40,40	1.77	3 (11%)
4	GAL	A	2	-	12,12,12	0.69	0	17,17,17	1.16	3 (17%)
3	GOL	A	355	-	5,5,5	1.30	1 (20%)	5,5,5	1.17	0
5	PEG	A	4	-	6,6,6	1.10	0	5,5,5	1.95	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	A	1	6	-	0/12/32/32	0/2/2/2
4	GAL	A	2	-	-	0/2/22/22	0/1/1/1
3	GOL	A	355	-	-	0/4/4/4	0/0/0/0
5	PEG	A	4	-	-	0/4/4/4	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	UDP	C6-C5	2.18	1.42	1.38
2	A	1	UDP	O4'-C1'	2.20	1.44	1.41
3	A	355	GOL	O2-C2	2.70	1.51	1.43
2	A	1	UDP	C4-N3	3.06	1.38	1.33
2	A	1	UDP	C6-N1	5.33	1.43	1.35

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	UDP	C4'-O4'-C1'	-3.40	105.98	109.72
4	A	2	GAL	O2-C2-C3	-2.53	104.65	110.34
2	A	1	UDP	PA-O3A-PB	-2.44	124.50	132.67
4	A	2	GAL	O1-C1-O5	-2.02	104.73	110.25
5	A	4	PEG	C3-O2-C2	2.24	122.93	113.31
4	A	2	GAL	O5-C1-C2	2.40	113.62	109.80
5	A	4	PEG	O2-C2-C1	3.04	124.42	110.43
2	A	1	UDP	C4-N3-C2	6.15	120.23	114.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	UDP	2	0
5	A	4	PEG	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	291/293 (99%)	0.98	35 (12%) 6 5	9, 16, 31, 58	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	354	PRO	9.8
1	A	196	CYS	7.3
1	A	198	ARG	6.5
1	A	80	CYS	6.3
1	A	349	GLN	5.6
1	A	350	ALA	5.4
1	A	161	ARG	5.3
1	A	64	VAL	5.0
1	A	62	PHE	4.9
1	A	351	VAL	4.8
1	A	348	HIS	4.6
1	A	197	GLU	4.5
1	A	63	MET	4.4
1	A	181	TRP	4.3
1	A	178	TYR	4.2
1	A	78	THR	3.8
1	A	177	ALA	3.6
1	A	256	ILE	3.1
1	A	353	ASN	2.9
1	A	345	PRO	2.9
1	A	347	ASN	2.9
1	A	146	ARG	2.8
1	A	184	VAL	2.7
1	A	201	LEU	2.6
1	A	328	GLN	2.4
1	A	311	LEU	2.4
1	A	151	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	175	VAL	2.3
1	A	289	MET	2.3
1	A	176	ARG	2.3
1	A	293	ALA	2.2
1	A	191	MET	2.2
1	A	294	ASN	2.1
1	A	122	ALA	2.1
1	A	147	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	A	355	6/6	0.81	0.15	1.22	26,29,33,36	0
4	GAL	A	2	12/12	0.87	0.19	0.88	21,24,32,35	0
2	UDP	A	1	25/25	0.94	0.10	-0.91	13,16,20,24	0
5	PEG	A	4	7/7	0.68	0.22	-	36,40,45,46	0
6	MN	A	356	1/1	0.99	0.04	-	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.