wwPDB NMR Structure Validation Summary Report (i)

Dec 12, 2016 – 03:42 PM EST

PDB ID	:	5T1O
Title	:	Solution-state NMR and SAXS structural ensemble of NPr (1-85) in complex with EIN-Ntr (170-424)
Authors	:	Strickland, M.; Stanley, A.M.; Wang, G.; Schwieters, C.D.; Buchanan, S.; Peterkofsky, A.; Tjandra, N.
Deposited on	:	2016-08-19

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20028442
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 33%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	85	34% 64% .
2	B	256	30% 48% . 20%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:1-A:85, B:170-B:187, B:202-B:311, B:322-B:397 (289)	0.49	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	4, 5, 8, 9, 10, 11, 13, 14, 15, 16, 17, 20
2	1, 2, 3, 7
3	12, 19
Single-model clusters	6; 18

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 5287 atoms, of which 2646 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Phosphocarrier protein NPr.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	85	1298	402	653	108	130	5	0

- Molecule 2 is a protein called Phosphoenolpyruvate-protein phosphotransferase PtsP.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
2	B	255	3989	1256	1993	354	383	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	169	GLY	-	expression tag	UNP P37177
B	356	GLN	HIS	engineered mutation	UNP P37177

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Phosphocarrier protein NPr

Chain A: 34% 64% •



- Molecule 2: Phosphoenolpyruvate-protein phosphotransferase PtsP

Chain B: 30% 48% • 20%



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 8. Colouring as in section 4.1 above.

- Molecule 1: Phosphocarrier protein NPr

Chain A: 38% 46% 16%



• Molecule 2: Phosphoenolpyruvate-protein phosphotransferase PtsP

Chain B:



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *SIMULATED ANNEALING*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *STRUCTURES WITH THE LOWEST ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	refinement	2.37.7
ANALYSIS	structure solution	2.4.2
TOPSPIN	structure solution	3.0
GAUSSIAN	structure solution	9
NMRPIPE	structure solution	8.2
TALOS-N	structure solution	4.12

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	5t1o_cs.cif
Number of chemical shift lists	2
Total number of shifts	1391
Number of shifts mapped to atoms	1391
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	33%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	645	653	653	65±15
2	B	1587	1608	1607	123±14
All	All	44640	45220	45200	3678

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 41.

5 of 2021 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:15:MET:CA	1:A:19:PRO:HG2	1.27	1.60	18	1
1:A:15:MET:HA	1:A:19:PRO:CD	1.27	1.56	18	1
1:A:10:THR:O	1:A:11:ASN:OD1	1.24	1.55	8	3
1:A:53:LEU:O	1:A:53:LEU:HD22	1.23	1.32	16	2
1:A:15:MET:HA	1:A:19:PRO:CG	1.20	1.67	18	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	83/85 (98%)	73±1 (88±1%)	8±1 (9±1%)	2±1 (3±1%)	9 43
2	B	203/256 (79%)	181±4 (89±2%)	17±4 (8±2%)	5±2 (2±1%)	12 49
All	All	5720/6820 (84%)	5078 (89%)	496 (9%)	146 (3%)	11 47

5 of 27 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	380	HIS	18
1	A	32	ALA	18
2	B	234	ALA	13
2	B	351	GLY	12
1	A	17	ALA	11

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	68/68 (100%)	54±3 (79±4%)	14±3 (21±4%)	4 33
2	B	160/205 (78%)	127±4 (79±3%)	33±4 (21±3%)	4 34
All	All	4560/5460 (84%)	3610 (79%)	950 (21%)	4 34

5 of 195 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	203	LEU	13
2	B	291	LEU	13
1	A	48	SER	12
1	A	80	ILE	12
2	B	303	GLN	12

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 33% for the well-defined parts and 34% for the entire structure.

7.1 Chemical shift list 1

File name: 5t1o_cs.cif

Chemical shift list name: *EIN_NPr_Complex.str*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1194
Number of shifts mapped to atoms	1194
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	249	0.13 ± 0.10	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	229	1.10 ± 0.07	Should be applied
$^{13}C'$	250	-0.23 ± 0.09	None needed (< 0.5 ppm)
^{15}N	232	0.70 ± 0.22	Should be applied

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 27%, i.e. 962 atoms were assigned a chemical shift out of a possible 3521. 0 out of 58 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	778/1427 (55%)	188/569 (33%)	402/578 (70%)	188/280 (67%)
Sidechain	182/1911 (10%)	0/1106 (0%)	182/708 (26%)	0/97 (0%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	2/183 (1%)	1/99 (1%)	0/78 (0%)	1/6 (17%)
Overall	962/3521 (27%)	189/1774 (11%)	584/1364 (43%)	189/383 (49%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:



7.2 Chemical shift list 2

File name: 5t1o_cs.cif

Chemical shift list name: *NPr_EIN_Complex.str*

7.2.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	197
Number of shifts mapped to atoms	197
Number of unparsed shifts	0

Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.2.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	51	0.61 ± 0.31	None needed (imprecise)
$^{13}C_{\beta}$	42	0.70 ± 0.08	Should be applied
$^{13}C'$	0	—	None (insufficient data)
^{15}N	52	0.17 ± 0.40	None needed (< 0.5 ppm)

7.2.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 6%, i.e. 197 atoms were assigned a chemical shift out of a possible 3521. 0 out of 58 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	155/1427 (11%)	52/569 (9%)	51/578 (9%)	52/280 (19%)
Sidechain	42/1911 (2%)	0/1106 (0%)	42/708 (6%)	0/97 (0%)
Aromatic	0/183 (0%)	0/99 (0%)	0/78 (0%)	0/6 (0%)
Overall	197/3521 (6%)	52/1774 (3%)	93/1364 (7%)	52/383 (14%)

7.2.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.2.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:


RCI
0.3
0.2
0.1
0
1:MET
4:LYS
7:VAL
10:THR
13:LEU
16:HIS
19:PRO
22:LYS
25:GLU
28:GLN
31:ASP
34:VAL
37:ARG
40:GLU
43:GLU
46:ALA
49:VAL
52:LEU
55:LEU
58:ALA
61:ARG
64:GLU
67:ALA
70:PRO
73:GLU
76:LEU
79:VAL
82:LEU
85:SER