



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:10 PM GMT

PDB ID : 3T3X  
Title : 1.57 Å structure of Friedreich's ataxia frataxin variant R165C  
Authors : Bridwell-Rabb, J.; Winn, A.M.; Barondeau, D.P.  
Deposited on : 2011-07-25  
Resolution : 1.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3815 (1.60-1.56)
Clashscore	102246	4131 (1.60-1.56)
Ramachandran outliers	100387	4021 (1.60-1.56)
Sidechain outliers	100360	4018 (1.60-1.56)
RSRZ outliers	91569	3823 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	129	<div> <div>4%</div> <div>78%</div> <div>17%</div> <div>5%</div> </div>
1	B	129	<div> <div>25%</div> <div>76%</div> <div>17%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	258	-	-	-	X
2	SO4	B	340	-	-	-	X

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2082 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Frataxin, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	122	Total	C	N	O	S	0	13	0
			992	632	149	208	3			
1	B	121	Total	C	N	O	S	0	4	0
			953	607	147	197	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	165	CYS	ARG	ENGINEERED MUTATION	UNP Q16595
B	165	CYS	ARG	ENGINEERED MUTATION	UNP Q16595

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	O S	0	0
			5	4 1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	102	Total	O	0	0
			102	102		
3	B	25	Total	O	0	0
			25	25		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Frataxin, mitochondrial



- Molecule 1: Frataxin, mitochondrial



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.02Å 32.12Å 92.08Å 90.00° 90.16° 90.00°	Depositor
Resolution (Å)	19.50 – 1.57 19.46 – 1.57	Depositor EDS
% Data completeness (in resolution range)	98.2 (19.50-1.57) 98.2 (19.46-1.57)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.19 (at 1.57Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.216 , 0.251 0.214 , 0.252	Depositor DCC
$R_{free}$ test set	1761 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.6	Xtriage
Anisotropy	0.211	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 52.9	EDS
Estimated twinning fraction	0.024 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	4 of 35083 reflections (0.011%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2082	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.42	4/1062 (0.4%)	1.46	17/1442 (1.2%)
1	B	1.05	0/987	1.04	2/1340 (0.1%)
All	All	1.26	4/2049 (0.2%)	1.28	19/2782 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	123	TYR	CE2-CZ	6.30	1.46	1.38
1	A	111	GLU	CD-OE2	5.93	1.32	1.25
1	A	123	TYR	CD2-CE2	5.73	1.48	1.39
1	A	166	TYR	CB-CG	-5.60	1.43	1.51

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	139	ASP	CB-CG-OD2	10.82	128.04	118.30
1	A	123	TYR	CB-CG-CD1	-10.65	114.61	121.00
1	A	143	TYR	CB-CG-CD2	-6.70	116.98	121.00
1	A	199	ASP	CB-CG-OD2	6.44	124.10	118.30
1	A	123	TYR	CB-CG-CD2	6.37	124.82	121.00
1	A	175	TYR	CB-CG-CD2	-6.14	117.31	121.00
1	A	123	TYR	CZ-CE2-CD2	-6.12	114.29	119.80

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	167	ASP	CB-CG-OD1	-5.78	113.10	118.30
1	A	104[A]	ASP	CB-CG-OD2	5.55	123.30	118.30
1	A	104[B]	ASP	CB-CG-OD2	5.55	123.30	118.30
1	A	190	LEU	CB-CG-CD1	-5.45	101.73	111.00
1	A	193	ALA	O-C-N	5.36	131.28	122.70
1	B	97	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	A	104[A]	ASP	CB-CG-OD1	-5.27	113.56	118.30
1	A	104[B]	ASP	CB-CG-OD1	-5.27	113.56	118.30
1	B	122	ASP	CB-CG-OD1	5.20	122.98	118.30
1	A	115	ASP	N-CA-CB	-5.14	101.36	110.60
1	A	132	LEU	CB-CG-CD1	-5.08	102.37	111.00
1	A	110	PHE	CB-CG-CD2	-5.06	117.26	120.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	209	ASP	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	992	0	925	9	0
1	B	953	0	904	17	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	102	0	0	1	2
3	B	25	0	0	2	0
All	All	2082	0	1829	26	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146[B]:ASN:OD1	3:A:350:HOH:O	1.54	1.23
1:A:153:GLN:NE2	1:A:165[C]:CYS:SG	2.44	0.91
1:A:146[C]:ASN:HD21	1:A:155[C]:TRP:HB3	1.59	0.66
1:B:156:LEU:HD22	1:B:182:LEU:HD12	1.78	0.65
1:B:139:ASP:OD1	1:B:139:ASP:N	2.13	0.65
1:B:151:ASN:HB3	3:B:318:HOH:O	1.95	0.65
1:A:153:GLN:CD	1:A:165[C]:CYS:SG	2.78	0.62
1:B:153:GLN:NE2	1:B:165[B]:CYS:SG	2.73	0.62
1:A:92:GLU:O	1:A:96:GLU:HG3	2.04	0.57
1:A:146[C]:ASN:ND2	1:A:155[C]:TRP:HB3	2.22	0.53
1:B:132:LEU:HD23	1:B:145:ILE:HD12	1.90	0.53
1:A:153:GLN:CD	1:A:165[C]:CYS:HG	2.14	0.50
1:B:148[A]:GLN:HG3	1:B:155:TRP:CD2	2.47	0.49
1:B:149:THR:O	1:B:150:PRO:C	2.51	0.48
1:B:153:GLN:CD	1:B:165[B]:CYS:SG	2.95	0.46
1:B:92:GLU:O	1:B:96:GLU:HG2	2.18	0.44
1:B:151:ASN:CA	3:B:318:HOH:O	2.65	0.44
1:B:145:ILE:HG12	1:B:156:LEU:HD13	2.00	0.42
1:B:142:THR:O	1:B:159:PRO:HD3	2.19	0.42
1:A:155[C]:TRP:CH2	1:A:157:SER:HB2	2.55	0.41
1:B:175:TYR:CE2	1:B:177:HIS:HB2	2.55	0.41
1:B:145:ILE:HG12	1:B:156:LEU:CD1	2.50	0.41
1:B:148[A]:GLN:HG3	1:B:155:TRP:CE3	2.56	0.41
1:B:95:TYR:CE2	1:B:149:THR:HA	2.57	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:307:HOH:O	3:A:310:HOH:O[4_456]	1.26	0.94
3:A:276:HOH:O	3:A:278:HOH:O[2_556]	1.98	0.22

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/129 (102%)	127 (97%)	4 (3%)	0	100	100
1	B	122/129 (95%)	118 (97%)	4 (3%)	0	100	100
All	All	253/258 (98%)	245 (97%)	8 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	116/109 (106%)	115 (99%)	1 (1%)	84	69
1	B	106/109 (97%)	105 (99%)	1 (1%)	84	69
All	All	222/218 (102%)	220 (99%)	2 (1%)	82	69

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	148	GLN
1	B	139	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	258	-	4,4,4	0.47	0	6,6,6	0.83	0
2	SO4	B	340	-	4,4,4	0.92	0	6,6,6	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	258	-	-	0/0/0/0	0/0/0/0
2	SO4	B	340	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	122/129 (94%)	0.33	5 (4%) 41 40	6, 10, 21, 34	0
1	B	121/129 (93%)	1.61	32 (26%) 1 1	17, 28, 38, 50	0
All	All	243/258 (94%)	0.97	37 (15%) 3 3	6, 20, 37, 50	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	179	GLY	7.0
1	B	90	LEU	6.8
1	B	210	ALA	6.5
1	B	178	ASP	5.7
1	B	176	SER	5.5
1	B	175	TYR	5.2
1	B	139	ASP	5.1
1	A	89	SER	5.0
1	B	155	TRP	4.8
1	B	180	VAL	4.0
1	B	163	PRO	3.9
1	B	151	ASN	3.8
1	B	138	GLY	3.7
1	B	127	PHE	3.5
1	A	209	ASP	3.5
1	B	165[A]	CYS	3.4
1	B	129	SER	3.3
1	B	156	LEU	3.3
1	B	177	HIS	3.2
1	B	150	PRO	3.1
1	B	182	LEU	3.1
1	B	106	LEU	3.0
1	B	162	GLY	2.9
1	B	160	SER	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	128	GLY	2.8
1	B	92	GLU	2.7
1	B	197	LYS	2.6
1	B	115	ASP	2.5
1	B	173	TRP	2.4
1	B	145	ILE	2.4
1	B	148[A]	GLN	2.3
1	A	155[B]	TRP	2.3
1	A	99	ALA	2.2
1	B	110	PHE	2.2
1	A	182	LEU	2.1
1	B	161	SER	2.1
1	B	117	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	258	5/5	0.96	0.14	3.89	25,27,28,28	0
2	SO4	B	340	5/5	0.95	0.19	2.70	26,26,27,30	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.