wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 05:13 PM BST

PDB ID	:	1T4Z
Title	:	Solution structure of the N-terminal domain of Synechococcus elongatus SasA (25-structures ensemble)
Authors	:	Vakonakis, I.; Klewer, D.A.; LiWang, A.C.
Deposited on	:	2004-04-30

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 79%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	105	32% 46% 6% 16%

2 Ensemble composition and analysis (i)

This entry contains 25 models. Model 11 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:11-A:98 (88)	0.25	11

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 7, 9, 11, 19, 21, 22, 24
2	2, 4, 6, 13, 18
3	8, 15, 16, 25
4	10, 12
5	5, 17
Single-model clusters	3; 14; 20; 23

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1663 atoms, of which 852 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Adaptive-response sensory-kinase sasA.

Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
1	A	105	1663	522	852	139	150	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	CLONING ARTIFACT	UNP Q06904
A	2	SER	-	CLONING ARTIFACT	UNP Q06904
A	103	GLY	-	CLONING ARTIFACT	UNP Q06904
A	104	ILE	-	CLONING ARTIFACT	UNP Q06904
A	105	PHE	-	CLONING ARTIFACT	UNP Q06904

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Adaptive-response sensory-kinase sasA

Chain A: 32% 46% 6% 16%

G1 S2 S3 L4 S5 P6 Q7 A8 L9 A10 L13 L14 L15 Q16 L17 F18 V19 D20 T21 R22 S25 V29 Q30 R31 N34 I35 L36 A37 A38 V39 E40 A41 I45 S46 L47 Q48 V49 I50 N51 Q55 L58 V59 Y62 R63 L64 V65 V66 A69 L70 V71 K72

I73 R78 Q79 V80 L81 S82 G83 I84 D85 L86 Q89 L90 A91 N92 Q93 L94 P95 Q96 W97 L98 V99 Q100 Q101 E102 G103 I104 F105

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 11. Colouring as in section 4.1 above.

- Molecule 1: Adaptive-response sensory-kinase sasA

Chain A: 40% 38% 6% 16%

G1 S2 S3 L4 S5 P6 Q7 A8 L9 A10 Q11 P12 L13 L14 L15 Q16 L17 D20 T21 R22 S25 I28 V29 Q30 N34 I35 L36 A37 A38 V39 E40 A41 I45 S46 L47 Q48 Q55 Y62 R63 L64 V65 V66 L70 V71 K72 I73 R78 Q79 S82 G83

I84 D85 L86 Q89 L90 A91 N92 Q93 L94 P95 V99 Q100 Q101 E102 G103 I104 F105

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *distance geometry, simulated annealing, slow refinement in the presence of database potentials.*

Of the 50 calculated structures, 25 were deposited, based on the following criterion: *structures with the lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
XPLOR-NIH	structure solution	2.9.1
XPLOR-NIH	refinement	2.9.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5141
Number of chemical shift lists	1
Total number of shifts	1130
Number of shifts mapped to atoms	1130
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	79%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.6±0.6
All	All	0	91

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	22	ARG	Sidechain	24
1	A	63	ARG	Sidechain	23
1	A	31	ARG	Sidechain	23
1	A	78	ARG	Sidechain	21

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	689	731	731	70±9
All	All	17225	18275	18275	1743

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 49.

5 of 351 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:25:SER:O	1:A:29:VAL:HG23	0.91	1.65	19	25
1:A:36:LEU:O	1:A:39:VAL:HG12	0.90	1.65	20	20
1:A:50:ILE:HG22	1:A:55:GLN:NE2	0.89	1.82	6	4
1:A:58:LEU:HD12	1:A:58:LEU:C	0.87	1.90	3	5
1:A:15:LEU:HD13	1:A:16:GLN:N	0.84	1.88	13	25

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	88/105 (84%)	82±1 (93±1%)	3±1 (4±1%)	3±1 (3±1%)	8	38
All	All	2200/2625 (84%)	2040 (93%)	85 (4%)	75 (3%)	8	38

5 of 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	84	ILE	25
1	A	40	GLU	17
1	A	41	ALA	14
1	A	39	VAL	9
1	A	64	LEU	4

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	79/92 (86%)	73±1 (93±1%)	6±1 (7±1%)	22	68
All	All	1975/2300 (86%)	1831 (93%)	144 (7%)	22	68

5 of 20 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	73	ILE	25
1	A	47	LEU	24
1	A	58	LEU	23
1	A	55	GLN	10
1	A	65	VAL	8

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 79% for the well-defined parts and 76% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 5141

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1130
Number of shifts mapped to atoms	1130
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	103	-0.17 ± 0.10	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	99	0.16 ± 0.05	None needed (< 0.5 ppm)
$^{13}C'$	0	—	—
^{15}N	99	0.38 ± 0.19	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 79%, i.e. 870 atoms were assigned a chemical shift out of a possible 1108. 13 out of 27 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	338/426 (79%)	169/169 (100%)	88/176 (50%)	81/81 (100%)
Sidechain	496/638 (78%)	296/369 (80%)	196/241 (81%)	4/28 (14%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	36/44 (82%)	18/23 (78%)	17/19 (89%)	1/2 (50%)
Overall	870/1108 (79%)	483/561 (86%)	301/436 (69%)	86/111 (77%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: