



wwPDB X-ray Structure Validation Summary Report

Feb 1, 2016 – 01:14 PM GMT

PDB ID : 3TAL
Title : Crystal structure of NurA with manganese
Authors : Chae, J.; Kim, Y.C.; Cho, Y.
Deposited on : 2011-08-04
Resolution : 3.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1112 (3.20-3.12)
Clashscore	102246	1249 (3.20-3.12)
Ramachandran outliers	100387	1222 (3.20-3.12)
Sidechain outliers	100360	1221 (3.20-3.12)
RSRZ outliers	91569	1117 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	471	
1	B	471	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	454	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6861 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA double-strand break repair protein nurA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	424	3420	2188	587	638	7	0	0	0
1	B	421	3411	2186	584	634	7	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MSE	-	EXPRESSION TAG	UNP Q8U1N8
A	-18	GLY	-	EXPRESSION TAG	UNP Q8U1N8
A	-17	SER	-	EXPRESSION TAG	UNP Q8U1N8
A	-16	SER	-	EXPRESSION TAG	UNP Q8U1N8
A	-15	HIS	-	EXPRESSION TAG	UNP Q8U1N8
A	-14	HIS	-	EXPRESSION TAG	UNP Q8U1N8
A	-13	HIS	-	EXPRESSION TAG	UNP Q8U1N8
A	-12	HIS	-	EXPRESSION TAG	UNP Q8U1N8
A	-11	HIS	-	EXPRESSION TAG	UNP Q8U1N8
A	-10	HIS	-	EXPRESSION TAG	UNP Q8U1N8
A	-9	SER	-	EXPRESSION TAG	UNP Q8U1N8
A	-8	SER	-	EXPRESSION TAG	UNP Q8U1N8
A	-7	GLY	-	EXPRESSION TAG	UNP Q8U1N8
A	-6	LEU	-	EXPRESSION TAG	UNP Q8U1N8
A	-5	VAL	-	EXPRESSION TAG	UNP Q8U1N8
A	-4	PRO	-	EXPRESSION TAG	UNP Q8U1N8
A	-3	ARG	-	EXPRESSION TAG	UNP Q8U1N8
A	-2	GLY	-	EXPRESSION TAG	UNP Q8U1N8
A	-1	SER	-	EXPRESSION TAG	UNP Q8U1N8
A	0	HIS	-	EXPRESSION TAG	UNP Q8U1N8
B	-19	MSE	-	EXPRESSION TAG	UNP Q8U1N8
B	-18	GLY	-	EXPRESSION TAG	UNP Q8U1N8
B	-17	SER	-	EXPRESSION TAG	UNP Q8U1N8
B	-16	SER	-	EXPRESSION TAG	UNP Q8U1N8
B	-15	HIS	-	EXPRESSION TAG	UNP Q8U1N8

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	EXPRESSION TAG	UNP Q8U1N8
B	-13	HIS	-	EXPRESSION TAG	UNP Q8U1N8
B	-12	HIS	-	EXPRESSION TAG	UNP Q8U1N8
B	-11	HIS	-	EXPRESSION TAG	UNP Q8U1N8
B	-10	HIS	-	EXPRESSION TAG	UNP Q8U1N8
B	-9	SER	-	EXPRESSION TAG	UNP Q8U1N8
B	-8	SER	-	EXPRESSION TAG	UNP Q8U1N8
B	-7	GLY	-	EXPRESSION TAG	UNP Q8U1N8
B	-6	LEU	-	EXPRESSION TAG	UNP Q8U1N8
B	-5	VAL	-	EXPRESSION TAG	UNP Q8U1N8
B	-4	PRO	-	EXPRESSION TAG	UNP Q8U1N8
B	-3	ARG	-	EXPRESSION TAG	UNP Q8U1N8
B	-2	GLY	-	EXPRESSION TAG	UNP Q8U1N8
B	-1	SER	-	EXPRESSION TAG	UNP Q8U1N8
B	0	HIS	-	EXPRESSION TAG	UNP Q8U1N8

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Mn 2 2	0	0
2	A	2	Total Mn 2 2	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	13	Total	O	0	0
			13	13		
4	B	7	Total	O	0	0
			7	7		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA double-strand break repair protein nurA



- Molecule 1: DNA double-strand break repair protein nurA



D193	Y260	L325	I388	PRO
N194	L261	I326	E389	LEU
V195	GLY	GLY	K390	GLU
V196	I266	HIS	I391	
E197	Y267	GLU	L392	
K198	D268	K330	P393	
L199	A269	E331	L394	
R200	L270	G332	L395	
K201	H271	Y333	L396	
Y202	M272	L334	H397	
Y203	T273	E335	H398	
L204	L274	I336	K399	
D205	S275	V340	A400	
T206	Y276	V341	GLY	
K207	L277	P342	GLY	
Y208	E278	P343	TYR	
ILE	Y279	K344	L404	
ALA	L280	K345	R405	
TYR	Y281	S346	P406	
GLY	S282	F347	L407	
SER	L283	P348	Q408	
GLY	D284	R349	L409	
K215	K285	D349	A410	
V216	L286	F350	H411	
K217	L287	L351	H412	
V218	L287	L352	G413	
V219	E288	S353	V414	
K220	M281	K354	K415	
K223	L292	F355	L416	
A224	A293	R356	S417	
L225	Y294	N357	Y418	
G226	L295	L358	K419	
Y227	A296	E359	E420	
S228	K297	K360	A421	
P229	F299	L361	R422	
L232	Y300	I362	H423	
L234	T301	D363	T424	
E235	K302	K364	L425	
V236	A305	G365	E426	
S240	R306	H367	A427	
R241	THR	L368	L428	
G242	LEU	A369	L429	
K243	GLY	V370	L432	
S244	V310	V371	R433	
V245	E311	R372	M434	
D246	K312	F373	R435	
E247	V313	E374	D436	
L248	D314	Q375	P437	
L249	L317	G376	A438	
Q250	L318	D377	L439	
K256	D319	V378	K440	
V257	A320	I379	ILE	
E258	L319	V380	PHE	
L259	Q321	M381	VAL	
K256	V321	L382	LYS	
V257	I322	Q383	TYR	
E258	R323	S384	GLY	
L259	T324	T385	ARG	
			SER	

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.80Å 114.65Å 121.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.48 – 3.15 34.37 – 3.10	Depositor EDS
% Data completeness (in resolution range)	96.8 (29.48-3.15) 94.4 (34.37-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 3.12Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.218 , 0.285 0.214 , 0.280	Depositor DCC
R_{free} test set	1583 reflections (10.07%)	DCC
Wilson B-factor (Å ²)	97.2	Xtrriage
Anisotropy	0.133	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 89.1	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Outliers	1 of 16840 reflections (0.006%)	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6861	wwPDB-VP
Average B, all atoms (Å ²)	125.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/3470	0.60	0/4670
1	B	0.32	0/3459	0.62	1/4652 (0.0%)
All	All	0.33	0/6929	0.61	1/9322 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	274	LEU	CA-CB-CG	5.97	129.03	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	228	SER	Peptide
1	B	417	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3420	0	3526	400	0
1	B	3411	0	3532	474	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	6	0	8	3	0
4	A	13	0	0	3	0
4	B	7	0	0	1	0
All	All	6861	0	7066	829	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 60.

The worst 5 of 829 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:172:HIS:ND1	1:B:191:LEU:HD11	1.57	1.20
1:A:54:ARG:HB3	1:A:101:MSE:HE3	1.29	1.14
1:B:98:ARG:HH22	1:B:137:PRO:HB3	0.99	1.10
1:A:244:SER:HA	1:B:244:SER:HA	1.24	1.09
1:A:240:SER:HB3	1:A:243:LYS:HD2	1.31	1.08

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	418/471 (89%)	370 (88%)	46 (11%)	2 (0%)	34	76
1	B	411/471 (87%)	345 (84%)	62 (15%)	4 (1%)	19	63
All	All	829/942 (88%)	715 (86%)	108 (13%)	6 (1%)	26	71

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	29	ILE
1	A	310	VAL
1	B	27	GLU
1	B	312	ILE
1	B	230	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	378/410 (92%)	325 (86%)	53 (14%)	4 19
1	B	380/410 (93%)	318 (84%)	62 (16%)	3 13
All	All	758/820 (92%)	643 (85%)	115 (15%)	3 16

5 of 115 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	428	LEU
1	B	126	ASP
1	B	407	LEU
1	B	20	LEU
1	B	101	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	397	HIS
1	B	28	GLN
1	B	398	HIS
1	A	271	HIS
1	B	185	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	454	-	5,5,5	0.32	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	454	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	454	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	417/471 (88%)	-0.16	10 (2%) 62 45	52, 103, 201, 270	0
1	B	414/471 (87%)	-0.11	9 (2%) 65 49	72, 130, 200, 374	0
All	All	831/942 (88%)	-0.13	19 (2%) 64 47	52, 118, 201, 374	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	34	ASN	6.5
1	A	301	THR	6.1
1	B	35	SER	4.9
1	A	338	HIS	3.1
1	B	318	LEU	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	A	454	6/6	0.80	0.86	13.80	95,114,126,129	0
2	MN	A	453	1/1	0.90	0.10	-	131,131,131,131	0
2	MN	B	453	1/1	0.29	0.26	-	235,235,235,235	0
2	MN	B	452	1/1	0.94	0.20	-	103,103,103,103	0
2	MN	A	452	1/1	0.99	0.28	-	93,93,93,93	0

6.5 Other polymers [i](#)

There are no such residues in this entry.