



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:18 PM GMT

PDB ID : 3THV
Title : Crystal Structure of Bacillus DNA Polymerase I Large Fragment Bound to DNA and ddATP-dT in Closed Conformation
Authors : Wang, W.; Beese, L.S.
Deposited on : 2011-08-19
Resolution : 1.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3202 (1.64-1.60)
Clashscore	102246	3500 (1.64-1.60)
Ramachandran outliers	100387	3411 (1.64-1.60)
Sidechain outliers	100360	3410 (1.64-1.60)
RSRZ outliers	91569	3207 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	592	<div> <div>3%</div> <div>90%</div> <div>8%</div> <div>.</div> </div>
1	D	592	<div> <div>92%</div> <div>6%</div> <div>.</div> </div>
2	B	9	<div> <div>67%</div> <div>33%</div> </div>
2	E	9	<div> <div>67%</div> <div>33%</div> </div>
3	C	13	<div> <div>54%</div> <div>38%</div> <div>8%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	13	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MG	D	2	-	-	-	X
6	SO4	D	3	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 21942 atoms, of which 9980 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase I.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	579	Total	C	H	N	O	S	0	2	0
			9378	2959	4724	809	869	17			
1	D	579	Total	C	H	N	O	S	0	12	0
			9447	2978	4756	814	882	17			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	456	GLU	ALA	SEE REMARK 999	UNP C9RTX7
A	505	LYS	GLU	SEE REMARK 999	UNP C9RTX7
A	598	ALA	ASP	ENGINEERED MUTATION	UNP C9RTX7
A	710	TYR	PHE	ENGINEERED MUTATION	UNP C9RTX7
A	823	HIS	ARG	SEE REMARK 999	UNP C9RTX7
D	456	GLU	ALA	SEE REMARK 999	UNP C9RTX7
D	505	LYS	GLU	SEE REMARK 999	UNP C9RTX7
D	598	ALA	ASP	ENGINEERED MUTATION	UNP C9RTX7
D	710	TYR	PHE	ENGINEERED MUTATION	UNP C9RTX7
D	823	HIS	ARG	SEE REMARK 999	UNP C9RTX7


- Molecule 2 is a DNA chain called 5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(2DA))-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	9	Total	C	H	N	O	P	0	0	0
			277	86	101	31	51	8			
2	E	9	Total	C	H	N	O	P	0	0	0
			277	86	101	31	51	8			

- Molecule 3 is a DNA chain called 5'-D(*C*AP*TP*TP*TP*GP*AP*GP*TP*CP*AP*GP*G)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	12	Total	C	H	N	O	P	0	0	0
			384	119	137	46	71	11			
3	F	12	Total	C	H	N	O	P	0	0	0
			384	119	137	46	71	11			

- Molecule 4 is 2',3'-DIDEOXYADENOSINE TRIPHOSPHATE (three-letter code: DDS) (formula: C₁₀H₁₆N₅O₁₁P₃).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	P	0	0
			41	10	12	5	11	3		
4	D	1	Total	C	H	N	O	P	0	0
			41	10	12	5	11	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	D	1	Total	Mg	0	0
			1	1		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	582	Total	O	0	0
			582	582		
7	D	859	Total	O	0	0
			859	859		
7	B	40	Total	O	0	0
			40	40		
7	C	80	Total	O	0	0
			80	80		
7	E	46	Total	O	0	0
			46	46		
7	F	89	Total	O	0	0
			89	89		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase I




- Molecule 1: DNA polymerase I



- Molecule 2: 5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(2DA))-3'



- Molecule 2: 5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(2DA))-3'



- Molecule 3: 5'-D(*C*AP*TP*TP*TP*GP*AP*GP*TP*CP*AP*GP*G)-3'





- Molecule 3: 5'-D(*C*AP*TP*TP*TP*GP*AP*GP*TP*CP*AP*GP*G)-3'

Chain F: 54% 38% 8%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	93.81Å 109.27Å 150.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.41 – 1.61 88.41 – 1.61	Depositor EDS
% Data completeness (in resolution range)	93.0 (88.41-1.61) 93.0 (88.41-1.61)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 1.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.193 , 0.224 0.198 , 0.227	Depositor DCC
R_{free} test set	8429 reflections (4.77%)	DCC
Wilson B-factor (Å ²)	17.0	Xtriage
Anisotropy	0.070	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 48.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 185311 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	21942	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DDS, MG, 2DA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/4747	0.51	0/6416
1	D	0.53	0/4822	0.64	0/6515
2	B	0.88	0/173	1.79	4/264 (1.5%)
2	E	1.05	0/173	1.91	3/264 (1.1%)
3	C	0.94	0/277	1.60	4/427 (0.9%)
3	F	1.12	0/277	1.60	3/427 (0.7%)
All	All	0.54	0/10469	0.76	14/14313 (0.1%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	8	DT	O4'-C1'-N1	-11.03	100.28	108.00
2	B	23	DT	O4'-C1'-N1	-9.04	101.67	108.00
2	E	23	DT	O4'-C1'-N1	-8.98	101.72	108.00
3	F	2	DT	O4'-C1'-N1	8.52	113.96	108.00
2	E	27	DT	O4'-C4'-C3'	-7.66	101.40	106.00
2	B	26	DC	O4'-C1'-N1	7.10	112.97	108.00
2	B	27	DT	O4'-C4'-C3'	-6.43	101.93	104.50
3	F	8	DT	O4'-C1'-N1	-6.32	103.58	108.00
3	C	12	DG	O4'-C1'-N9	6.16	112.31	108.00
3	F	2	DT	O4'-C1'-C2'	-6.02	101.08	105.90
2	B	23	DT	N3-C2-O2	-5.43	119.04	122.30
3	C	2	DT	O4'-C1'-N1	5.38	111.77	108.00
3	C	11	DG	O4'-C1'-N9	-5.32	104.28	108.00
2	E	26	DC	N1-C2-O2	-5.11	115.84	118.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4654	4724	4703	34	0
1	D	4691	4756	4690	26	0
2	B	176	101	103	0	0
2	E	176	101	103	0	0
3	C	247	137	138	1	0
3	F	247	137	138	3	0
4	A	29	12	12	0	0
4	D	29	12	12	1	0
5	A	1	0	0	0	0
5	D	1	0	0	0	0
6	A	5	0	0	0	0
6	D	10	0	0	0	0
7	A	582	0	0	8	1
7	B	40	0	0	0	0
7	C	80	0	0	0	0
7	D	859	0	0	12	1
7	E	46	0	0	0	0
7	F	89	0	0	2	0
All	All	11962	9980	9899	63	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:679:LEU:HD23	7:D:1137:HOH:O	1.84	0.76
1:D:653:ASP:HB2	7:D:1696:HOH:O	1.95	0.67
1:D:754:VAL:HG12	1:D:758:LYS:HE2	1.77	0.66
1:D:466:ARG:NH2	7:D:1078:HOH:O	2.27	0.63
1:A:848:VAL:HB	1:A:849:PRO:HD3	1.79	0.63
1:D:754:VAL:CG1	1:D:758:LYS:HE2	2.29	0.62
1:A:694:GLU:HG2	1:A:694:GLU:O	2.00	0.61
1:A:431:LYS:NZ	7:A:1179:HOH:O	2.36	0.59
1:D:456:GLU:HG2	7:D:1245:HOH:O	2.05	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:509:GLU:HG3	7:D:984:HOH:O	2.04	0.56
1:A:459:ARG:HB3	1:A:460:PRO:HD3	1.88	0.56
1:A:843:ARG:NH1	7:A:1364:HOH:O	2.40	0.55
1:D:315:LYS:HG3	7:D:978:HOH:O	2.09	0.53
3:F:9:DC:H2'	7:F:1621:HOH:O	2.08	0.53
1:A:692:VAL:HB	1:A:696:GLU:HB2	1.90	0.53
1:D:653:ASP:OD2	7:D:1695:HOH:O	2.19	0.53
1:D:748:ARG:HD3	7:D:1097:HOH:O	2.10	0.51
1:D:500:LEU:HD12	1:D:503:MET:HE3	1.93	0.51
1:A:504:GLY:HA2	1:A:588:ILE:HD12	1.91	0.51
3:F:9:DC:H2''	3:F:10:DA:C8	2.47	0.50
1:D:744:PRO:O	1:D:748:ARG:HG3	2.12	0.49
1:D:558:ALA:O	1:D:562:GLU:HG3	2.12	0.49
1:A:325:GLU:OE1	1:A:435:ARG:NH1	2.30	0.48
1:A:440:GLU:HB3	1:A:441:PRO:HD3	1.95	0.48
1:A:411:ALA:O	1:A:415:LYS:HG2	2.14	0.47
1:A:825:LEU:HD11	1:A:835:GLU:HB3	1.96	0.47
1:A:325:GLU:CD	1:A:325:GLU:H	2.16	0.47
1:A:670:ASN:ND2	1:A:742:SER:O	2.48	0.47
1:D:820:LEU:HD21	1:D:843:ARG:CZ	2.45	0.47
1:A:683:THR:O	1:A:687:MET:HG3	2.15	0.46
1:D:866:TYR:HE1	7:D:1057:HOH:O	1.99	0.46
1:A:415:LYS:NZ	1:A:420:GLU:OE2	2.49	0.46
1:D:658:GLU:CD	4:D:202:DDS:H2'A	2.35	0.46
1:A:381:LYS:NZ	7:A:1633:HOH:O	2.48	0.46
1:A:767:LEU:HG	1:A:802:ASP:HB3	1.98	0.45
1:A:680:ASP:HB3	1:A:683:THR:OG1	2.17	0.45
1:A:814:ARG:HG2	7:A:1350:HOH:O	2.15	0.45
1:A:668:ASP:OD1	1:A:745:GLY:N	2.50	0.45
1:A:693:SER:C	1:A:695:ASP:H	2.20	0.44
1:D:847:LEU:HD23	1:D:847:LEU:C	2.38	0.44
1:A:700:ASN:ND2	7:A:1437:HOH:O	2.51	0.44
1:A:767:LEU:O	1:A:768:HIS:HB2	2.18	0.44
1:D:645:SER:O	1:D:646:ASP:HB2	2.18	0.43
1:D:757:ALA:HB3	1:D:776:ILE:HD13	2.01	0.43
1:A:711:GLY:HA2	3:C:3:DT:C2	2.53	0.43
1:A:684:LYS:HA	1:A:687:MET:HE3	1.99	0.42
1:D:534:LEU:HD11	1:D:574:ILE:HD13	2.00	0.42
1:D:423:ARG:HG2	1:D:423:ARG:HH11	1.84	0.42
1:A:689:ILE:HD11	1:A:708:VAL:HG11	2.01	0.42
1:D:691:GLN:HA	1:D:691:GLN:OE1	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:843:ARG:HD3	7:A:1053:HOH:O	2.20	0.42
1:A:687:MET:SD	1:A:694:GLU:HB2	2.60	0.42
1:D:298:LYS:CD	7:D:1066:HOH:O	2.68	0.42
1:A:429:TYR:O	1:A:435:ARG:HA	2.19	0.42
1:A:534:LEU:HD11	1:A:574:ILE:HD13	2.01	0.41
1:A:468:ASN:O	1:A:469:GLU:HB2	2.20	0.41
3:F:11:DG:OP1	7:F:566:HOH:O	2.21	0.41
1:D:306:ARG:NH2	7:D:1495:HOH:O	2.47	0.41
1:A:794:THR:N	1:A:795:PRO:CD	2.83	0.41
1:A:734:GLU:HG2	7:A:949:HOH:O	2.20	0.41
1:D:712:ILE:HA	1:D:716:ILE:HG22	2.02	0.41
1:D:302:THR:HG21	7:D:1488:HOH:O	2.20	0.41
1:A:738:ARG:NE	7:A:1502:HOH:O	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:1190:HOH:O	7:D:947:HOH:O[4_445]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	579/592 (98%)	563 (97%)	15 (3%)	1 (0%)	52	27
1	D	589/592 (100%)	577 (98%)	11 (2%)	1 (0%)	52	27
All	All	1168/1184 (99%)	1140 (98%)	26 (2%)	2 (0%)	52	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	694	GLU
1	D	628	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	497/507 (98%)	496 (100%)	1 (0%)	95	90
1	D	507/507 (100%)	504 (99%)	3 (1%)	90	81
All	All	1004/1014 (99%)	1000 (100%)	4 (0%)	93	88

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	681	ILE
1	D	303	LEU
1	D	728	SER
1	D	730	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	2DA	B	29	3,2	14,22,23	0.68	0	13,31,34	1.09	1 (7%)
2	2DA	E	29	3,2	14,22,23	0.74	0	13,31,34	1.05	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2DA	B	29	3,2	-	0/3/18/19	0/3/3/3
2	2DA	E	29	3,2	-	0/3/18/19	0/3/3/3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	E	29	2DA	C3'-C2'-C1'	-2.57	99.83	102.71
2	B	29	2DA	O4'-C4'-C5'	-2.23	106.25	109.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
6	SO4	A	2	-	4,4,4	0.17	0	6,6,6	0.25	0
4	DDS	A	201	5	23,31,31	1.24	2 (8%)	28,48,48	3.06	8 (28%)
6	SO4	D	1	-	4,4,4	0.34	0	6,6,6	0.29	0
4	DDS	D	202	5	23,31,31	1.07	2 (8%)	28,48,48	2.25	2 (7%)
6	SO4	D	3	-	4,4,4	0.16	0	6,6,6	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SO4	A	2	-	-	0/0/0/0	0/0/0/0
4	DDS	A	201	5	-	0/18/31/31	0/3/3/3
6	SO4	D	1	-	-	0/0/0/0	0/0/0/0
4	DDS	D	202	5	-	0/18/31/31	0/3/3/3
6	SO4	D	3	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	201	DDS	C3'-C2'	-2.21	1.47	1.54
4	D	202	DDS	C3'-C2'	-2.15	1.47	1.54
4	D	202	DDS	C6-N6	2.54	1.42	1.34
4	A	201	DDS	C6-N6	3.28	1.45	1.34

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	201	DDS	N3-C2-N1	-12.33	119.46	128.89
4	D	202	DDS	N3-C2-N1	-9.89	121.33	128.89
4	A	201	DDS	C2'-C1'-N9	-3.40	105.60	112.49
4	A	201	DDS	C1'-N9-C4	-3.28	121.59	127.16
4	A	201	DDS	C4-C5-N7	-3.10	106.62	109.48
4	A	201	DDS	PB-O3A-PA	-2.72	125.10	132.73
4	A	201	DDS	C2-N1-C6	2.11	122.53	118.77
4	A	201	DDS	O2B-PB-O3B	2.82	117.91	105.09
4	D	202	DDS	O4'-C1'-N9	3.53	113.83	107.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	201	DDS	O4'-C1'-N9	6.19	118.44	107.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	202	DDS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	579/592 (97%)	0.14	15 (2%) 59 56	14, 29, 49, 59	0
1	D	579/592 (97%)	-0.15	1 (0%) 95 94	7, 17, 33, 45	0
2	B	8/9 (88%)	-0.52	0 100 100	15, 17, 29, 34	0
2	E	8/9 (88%)	-0.43	0 100 100	10, 15, 29, 33	0
3	C	12/13 (92%)	-0.33	0 100 100	14, 19, 39, 55	0
3	F	12/13 (92%)	-0.19	0 100 100	9, 13, 32, 49	0
All	All	1198/1228 (97%)	-0.02	16 (1%) 79 78	7, 22, 45, 59	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	677	ARG	4.6
1	A	819	ARG	4.0
1	A	689	ILE	3.6
1	A	671	LEU	3.5
1	A	846	ARG	3.4
1	A	738	ARG	3.2
1	A	298	LYS	2.8
1	A	866	TYR	2.4
1	A	847	LEU	2.4
1	A	679	LEU	2.4
1	A	816	LYS	2.1
1	A	672	MET	2.1
1	A	814	ARG	2.0
1	D	779	ARG	2.0
1	A	735	PHE	2.0
1	A	857	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	2DA	E	29	20/21	0.99	0.09	-	7,9,12,13	0
2	2DA	B	29	20/21	0.97	0.08	-	13,19,24,28	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	MG	D	2	1/1	0.99	0.13	3.72	13,13,13,13	0
6	SO4	D	3	5/5	0.81	0.12	2.44	77,80,83,89	0
4	DDS	D	202	29/29	0.98	0.09	0.55	8,11,14,17	0
4	DDS	A	201	29/29	0.94	0.10	0.07	17,21,27,32	0
5	MG	A	1	1/1	0.93	0.07	-2.53	26,26,26,26	0
6	SO4	D	1	5/5	0.97	0.12	-	26,33,42,51	0
6	SO4	A	2	5/5	0.94	0.12	-	35,51,51,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.