



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:26 PM GMT

PDB ID : 1TMO
Title : TRIMETHYLAMINE N-OXIDE REDUCTASE FROM SHEWANELLA MASSILIA
Authors : Czjzek, M.; Dos Santos, J.P.; Giordano, G.; Mejean, V.
Deposited on : 1998-08-03
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	829	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey); position: relative;"> % 84% 10% ... </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	2MO	A	801	-	-	X	X

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6857 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRIMETHYLAMINE N-OXIDE REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	794	Total	C	N	O	S	0	0	0
			6255	3972	1080	1167	36			

- Molecule 2 is GUANYLATE-O'-PHOSPHORIC ACID MONO-(2-AMINO-5,6-DIMERCAPTO-4-OXO-3,5,6,7,8A,9,10,10A-OCTAHYDRO-4H-8-OXA-1,3,9,10-TETRAAZA-ANTHRACEN-7-YLMETHYL) ESTER (three-letter code: 2MD) (formula: $C_{20}H_{28}N_{10}O_{13}P_2S_2$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Mo	O	0	0
			3	1	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	505	Total	O	0	0
			505	505		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	123.32Å 140.70Å 59.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 19.87 – 2.40	Depositor EDS
% Data completeness (in resolution range)	90.5 (20.00-2.50) 82.6 (19.87-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.41Å)	Xtriage
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.182 , 0.247 0.209 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	14.5	Xtriage
Anisotropy	0.533	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 61.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 34209 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6857	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2MO, 2MD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/6422	0.66	0/8702

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6255	0	6069	53	0
2	A	94	0	44	4	0
3	A	3	0	0	5	0
4	A	505	0	0	6	0
All	All	6857	0	6113	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:392:PRO:HG2	1:A:395:ASP:HB2	1.57	0.87
1:A:5:ASN:N	1:A:24:LYS:HG3	1.92	0.84
3:A:801:2MO:MO	3:A:801:2MO:OT2	1.57	0.75
1:A:657:HIS:CE1	2:A:800:2MD:S13	2.81	0.73
3:A:801:2MO:OT2	3:A:801:2MO:OT1	2.07	0.72
1:A:663:SER:HB3	4:A:826:HOH:O	1.91	0.70
3:A:801:2MO:MO	3:A:801:2MO:OT1	1.64	0.69
1:A:422:ILE:HG22	1:A:429:VAL:HB	1.74	0.68
1:A:58:ARG:HH11	1:A:58:ARG:HG3	1.58	0.68
1:A:664:ARG:HH11	1:A:664:ARG:HB3	1.60	0.66
1:A:204:GLU:HG3	1:A:502:ARG:HH12	1.64	0.62
1:A:190:PRO:O	1:A:194:LEU:HB2	2.06	0.56
1:A:550:LEU:O	1:A:553:GLU:HG3	2.05	0.56
1:A:112:HIS:HB3	1:A:437:MET:HG2	1.88	0.55
1:A:15:HIS:HE1	1:A:119:ARG:O	1.89	0.54
1:A:657:HIS:NE2	2:A:800:2MD:S13	2.81	0.54
1:A:633:ARG:HB3	1:A:637:GLY:HA3	1.90	0.54
1:A:187:SER:O	1:A:327:GLY:HA3	2.08	0.54
1:A:678:VAL:HG22	1:A:707:ALA:CB	2.39	0.53
1:A:114:GLY:HA3	1:A:143:LYS:HE3	1.88	0.53
1:A:657:HIS:HE1	2:A:800:2MD:S13	2.32	0.53
1:A:57:VAL:HG12	1:A:483:PRO:HB3	1.89	0.52
1:A:678:VAL:HG22	1:A:707:ALA:HB1	1.94	0.50
1:A:757:LEU:HD12	1:A:757:LEU:O	2.11	0.50
2:A:800:2MD:S13	3:A:801:2MO:OT2	2.70	0.49
1:A:328:TRP:O	1:A:331:GLN:HG3	2.12	0.49
1:A:723:HIS:HD2	4:A:1216:HOH:O	1.95	0.49
1:A:334:GLN:HE22	1:A:631:THR:H	1.61	0.48
1:A:124:LEU:O	1:A:130:HIS:HE1	1.96	0.48
1:A:665:GLU:H	1:A:665:GLU:CD	2.17	0.48
1:A:113:ALA:HB1	1:A:131:MET:HE1	1.95	0.47
1:A:152:ALA:HB1	1:A:337:GLU:HG3	1.97	0.47
1:A:648:GLN:NE2	1:A:721:ARG:HE	2.12	0.47
1:A:503:GLY:HA2	1:A:578:PHE:CD2	2.50	0.46
1:A:257:ALA:O	1:A:261:ILE:HG12	2.16	0.46
1:A:282:VAL:HB	1:A:283:PRO:HD3	1.97	0.46
1:A:759:ILE:HD13	1:A:759:ILE:HA	1.84	0.46
1:A:716:PRO:HG2	1:A:719:ILE:HD12	1.97	0.45
1:A:113:ALA:HB1	1:A:131:MET:CE	2.46	0.45
1:A:133:ARG:O	1:A:137:MET:HG3	2.16	0.45
1:A:149:SER:OG	3:A:801:2MO:OT1	2.36	0.44
1:A:204:GLU:HG3	1:A:502:ARG:HH22	1.82	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:502:ARG:NE	4:A:955:HOH:O	2.51	0.43
1:A:312:ASP:OD1	1:A:315:LYS:HE3	2.18	0.43
1:A:678:VAL:HG13	1:A:722:ILE:HG12	1.99	0.43
1:A:622:LYS:HG2	4:A:948:HOH:O	2.19	0.42
1:A:258:HIS:CE1	1:A:295:THR:HG22	2.54	0.42
1:A:58:ARG:NH1	4:A:833:HOH:O	2.52	0.42
1:A:202:THR:O	1:A:203:HIS:HB2	2.19	0.42
1:A:680:ILE:O	1:A:711:VAL:HA	2.20	0.42
1:A:143:LYS:HE2	1:A:144:LYS:H	1.84	0.42
1:A:58:ARG:CG	1:A:58:ARG:HH11	2.28	0.42
1:A:269:LYS:HB2	1:A:269:LYS:NZ	2.36	0.41
1:A:675:ARG:HD2	1:A:675:ARG:N	2.35	0.41
1:A:99:LEU:HB3	1:A:138:HIS:CE1	2.56	0.41
1:A:641:ASP:HB2	4:A:1029:HOH:O	2.21	0.40
1:A:53:ASN:HD21	1:A:470:ASN:HD21	1.70	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	792/829 (96%)	749 (95%)	40 (5%)	3 (0%)	39 61

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	664	ARG
1	A	660	MET
1	A	388	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	665/692 (96%)	635 (96%)	30 (4%)	34 59

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ARG
1	A	51	VAL
1	A	59	TYR
1	A	64	LEU
1	A	68	LEU
1	A	81	PHE
1	A	115	GLN
1	A	119	ARG
1	A	126	SER
1	A	131	MET
1	A	143	LYS
1	A	144	LYS
1	A	194	LEU
1	A	204	GLU
1	A	217	LYS
1	A	252	LEU
1	A	411	TRP
1	A	416	LEU
1	A	425	ASN
1	A	443	ASN
1	A	496	TYR
1	A	553	GLU
1	A	591	ARG
1	A	595	GLU
1	A	631	THR
1	A	664	ARG
1	A	665	GLU
1	A	675	ARG
1	A	678	VAL
1	A	761	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	HIS
1	A	71	HIS
1	A	115	GLN
1	A	130	HIS
1	A	245	ASN
1	A	334	GLN
1	A	425	ASN
1	A	443	ASN
1	A	470	ASN
1	A	573	GLN
1	A	635	HIS
1	A	643	HIS
1	A	648	GLN
1	A	651	HIS
1	A	723	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	2MD	A	799	3	40,52,52	2.07	10 (25%)	50,81,81	5.18	25 (50%)
2	2MD	A	800	3	40,52,52	2.14	11 (27%)	50,81,81	4.02	20 (40%)
3	2MO	A	801	1,2	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2MD	A	799	3	-	0/18/66/66	0/6/6/6
2	2MD	A	800	3	-	0/18/66/66	0/6/6/6
3	2MO	A	801	1,2	-	0/0/0/0	0/0/0/0

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	799	2MD	C13-C12	-3.23	1.34	1.51
2	A	799	2MD	O11-C7	-2.54	1.40	1.43
2	A	800	2MD	O11-C7	-2.18	1.40	1.43
2	A	800	2MD	C13-C12	-2.08	1.40	1.51
2	A	799	2MD	O4'-C1'	2.10	1.43	1.41
2	A	800	2MD	C15-N17	2.21	1.37	1.33
2	A	800	2MD	C16-C20	2.34	1.46	1.41
2	A	799	2MD	C4-N3	2.42	1.39	1.35
2	A	799	2MD	C2-N1	2.61	1.40	1.35
2	A	799	2MD	C16-C20	2.66	1.47	1.41
2	A	800	2MD	C2-N1	2.76	1.40	1.35
2	A	799	2MD	C6-C5	2.99	1.47	1.41
2	A	800	2MD	C6-C5	3.00	1.47	1.41
2	A	800	2MD	C14-N15	3.49	1.48	1.45
2	A	800	2MD	O4'-C1'	4.31	1.46	1.41
2	A	800	2MD	C6-N1	4.84	1.42	1.33
2	A	800	2MD	C7-C14	5.10	1.57	1.53
2	A	799	2MD	C6-N1	5.18	1.42	1.33
2	A	799	2MD	C7-C14	5.84	1.58	1.53
2	A	799	2MD	O14-C15	5.94	1.38	1.24
2	A	800	2MD	O14-C15	6.07	1.39	1.24

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	800	2MD	C5-C6-N1	-8.59	111.84	123.59
2	A	799	2MD	C5-C6-N1	-8.48	112.00	123.59
2	A	800	2MD	C4'-O4'-C1'	-4.25	105.05	109.72
2	A	800	2MD	O11-C11-C10	-4.11	98.20	106.61
2	A	800	2MD	N17-C17-N18	-3.63	119.59	125.53
2	A	799	2MD	N3-C2-N1	-3.55	122.03	127.44
2	A	799	2MD	C20-N8-C7	-3.46	116.89	123.67
2	A	799	2MD	PA-O3B-PB	-3.46	123.01	132.73
2	A	800	2MD	O11-C7-C14	-3.43	106.56	108.97
2	A	800	2MD	PA-O3B-PB	-3.34	123.34	132.73
2	A	800	2MD	N3-C2-N1	-3.01	122.86	127.44
2	A	799	2MD	O11-C11-C10	-2.77	100.96	106.61
2	A	799	2MD	N17-C17-N18	-2.43	121.55	125.53
2	A	800	2MD	C20-N8-C7	-2.32	119.14	123.67
2	A	799	2MD	C6-C5-C4	-2.05	118.45	120.90
2	A	799	2MD	C2'-C1'-N9	-2.01	111.23	114.29
2	A	799	2MD	N2-C2-N3	2.01	121.66	117.80
2	A	800	2MD	O2B-PB-O3B	2.05	114.42	105.09
2	A	799	2MD	N16-C17-N17	2.10	120.67	117.20
2	A	799	2MD	O3B-PA-O5'	2.20	108.78	102.94
2	A	800	2MD	C16-C20-N8	2.47	120.93	118.34
2	A	800	2MD	O3B-PA-O5'	2.47	109.49	102.94
2	A	799	2MD	C15-N17-C17	2.61	119.57	115.94
2	A	800	2MD	O3A-C10-C11	2.65	118.88	109.12
2	A	799	2MD	O4'-C1'-N9	2.79	113.94	108.10
2	A	800	2MD	O4'-C1'-N9	2.80	113.96	108.10
2	A	800	2MD	N16-C17-N17	2.91	122.02	117.20
2	A	799	2MD	C7-C14-N15	3.01	115.17	108.89
2	A	799	2MD	C15-C16-C20	3.04	117.32	114.56
2	A	799	2MD	O11-C7-C14	3.38	111.33	108.97
2	A	799	2MD	C17-N18-C20	3.54	122.49	114.54
2	A	800	2MD	C17-N18-C20	3.63	122.71	114.54
2	A	799	2MD	O3A-C10-C11	3.65	122.57	109.12
2	A	800	2MD	C15-N17-C17	3.77	121.17	115.94
2	A	799	2MD	C20-C16-N15	3.86	123.71	118.85
2	A	800	2MD	C10-C11-C12	4.16	121.56	112.09
2	A	799	2MD	C16-C20-N8	4.32	122.86	118.34
2	A	799	2MD	C10-C11-C12	4.89	123.23	112.09
2	A	799	2MD	O3B-PB-O3A	5.83	118.40	102.94
2	A	800	2MD	C6-N1-C2	6.13	124.45	115.94
2	A	799	2MD	O11-C11-C12	6.31	121.49	109.68
2	A	799	2MD	C6-N1-C2	6.72	125.26	115.94
2	A	800	2MD	C11-O11-C7	9.57	126.41	112.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	800	2MD	C13-C14-N15	20.11	123.21	109.73
2	A	799	2MD	C13-C14-N15	30.52	130.19	109.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	800	2MD	4	0
3	A	801	2MO	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	794/829 (95%)	-0.29	5 (0%) 90 91	2, 11, 29, 64	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	ASN	5.1
1	A	7	ASP	3.0
1	A	6	GLU	2.8
1	A	395	ASP	2.6
1	A	559	ALA	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	2MO	A	801	3/3	0.98	0.22	4.97	16,16,18,20	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	2MD	A	800	47/47	0.94	0.12	-0.04	3,11,16,24	0
2	2MD	A	799	47/47	0.94	0.11	-0.97	1,9,13,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.