



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:33 PM GMT

PDB ID : 4TMR
Title : Crystal structure of ternary complex of Plasmodium vivax SHMT with glycine and a novel pyrazolopyran 99S: methyl 5-{3-[(4S)-6-amino-5-cyano-3-methyl-4-(propan-2-yl)-2,4-dihydropyrano[2,3-c]pyrazol-4-yl]-5-cyanophenyl}thiophene-2-carboxylate .
Authors : Chitnumsub, P.; Jaruwat, A.; Leartsakulpanich, U.; Witschel, M.C.
Deposited on : 2014-06-02
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	442	<div> <div>2%</div> <div>76%</div> <div>21%</div> <div>.</div> </div>
1	B	442	<div> <div>3%</div> <div>75%</div> <div>24%</div> <div>.</div> </div>
1	C	442	<div> <div>5%</div> <div>74%</div> <div>23%</div> <div>.</div> </div>

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 10752 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine hydroxymethyltransferase, putative.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	442	Total	C	N	O	S	0	0	0
			3458	2186	600	655	17			
1	B	442	Total	C	N	O	S	0	0	0
			3458	2186	600	655	17			
1	C	442	Total	C	N	O	S	0	0	0
			3458	2186	600	655	17			

- Molecule 2 is N-GLYCINE-[3-HYDROXY-2-METHYL-5-PHOSPHONOOXYMETHYL-PYRIDIN-4-YL-METHANE] (three-letter code: PLG) (formula: $C_{10}H_{15}N_2O_7P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			20	10	2	7	1		
2	B	1	Total	C	N	O	P	0	0
			20	10	2	7	1		
2	C	1	Total	C	N	O	P	0	0
			20	10	2	7	1		

- Molecule 3 is methyl 5-{3-[(4S)-6-amino-5-cyano-3-methyl-4-(propan-2-yl)-2,4-dihydropyrano[2,3-c]pyrazol-4-yl]-5-cyanophenyl} thiophene-2-carboxylate (three-letter code: 99S) (formula: C₂₄H₂₁N₅O₃S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			33	24	5	3	1		
3	B	1	Total	C	N	O	S	0	0
			33	24	5	3	1		
3	C	1	Total	C	N	O	S	0	0
			33	24	5	3	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		
4	C	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	83	Total	O	0	0
			83	83		
5	B	69	Total	O	0	0
			69	69		
5	C	65	Total	O	0	0
			65	65		



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	100.81Å 58.26Å 234.96Å 90.00° 90.01° 90.00°	Depositor
Resolution (Å)	30.00 – 2.70 29.37 – 2.70	Depositor EDS
% Data completeness (in resolution range)	91.9 (30.00-2.70) 92.0 (29.37-2.70)	Depositor EDS
R_{merge}	0.02	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	23.58 (at 2.72Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.210 , 0.278 0.203 , 0.271	Depositor DCC
R_{free} test set	3462 reflections (11.04%)	DCC
Wilson B-factor (Å ²)	37.7	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 7.1	EDS
Estimated twinning fraction	0.024 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.023 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.470 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.476 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.018 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 34824 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10752	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLG, 99S, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	1/3521 (0.0%)	0.62	1/4754 (0.0%)
1	B	0.51	0/3521	0.64	0/4754
1	C	0.51	1/3521 (0.0%)	0.66	2/4754 (0.0%)
All	All	0.51	2/10563 (0.0%)	0.64	3/14262 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	435	TRP	CD2-CE2	5.13	1.47	1.41
1	A	435	TRP	CD2-CE2	5.09	1.47	1.41

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	327	LEU	CA-CB-CG	5.83	128.71	115.30
1	C	327	LEU	CA-CB-CG	5.66	128.31	115.30
1	C	226	LEU	CA-CB-CG	5.03	126.87	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3458	0	3477	77	0
1	B	3458	0	3477	85	0
1	C	3458	0	3477	67	0
2	A	20	0	11	3	0
2	B	20	0	10	0	0
2	C	20	0	11	2	0
3	A	33	0	21	4	0
3	B	33	0	21	4	0
3	C	33	0	21	5	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	83	0	0	0	0
5	B	69	0	0	2	0
5	C	65	0	0	3	0
All	All	10752	0	10526	234	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 234 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:385:LYS:H	1:A:385:LYS:HD3	1.14	1.07
3:B:502:99S:H19	3:B:502:99S:H12	1.45	0.98
1:A:385:LYS:N	1:A:385:LYS:HD3	1.79	0.97
3:A:502:99S:H19	3:A:502:99S:H12	1.55	0.87
1:B:91:GLU:HG3	1:B:252:LYS:HD2	1.62	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	440/442 (100%)	423 (96%)	16 (4%)	1 (0%)	52	80
1	B	440/442 (100%)	421 (96%)	19 (4%)	0	100	100
1	C	440/442 (100%)	413 (94%)	25 (6%)	2 (0%)	34	63
All	All	1320/1326 (100%)	1257 (95%)	60 (4%)	3 (0%)	52	80

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	366	SER
1	C	222	ASN
1	C	174	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	381/381 (100%)	363 (95%)	18 (5%)	32	63
1	B	381/381 (100%)	364 (96%)	17 (4%)	34	65
1	C	381/381 (100%)	356 (93%)	25 (7%)	21	45
All	All	1143/1143 (100%)	1083 (95%)	60 (5%)	29	58

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	219	ASN
1	B	417	LYS
1	C	411	LYS
1	B	357	THR
1	C	21	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	424	GLN
1	C	321	ASN
1	C	86	ASN
1	B	86	ASN
1	C	227	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLG	A	501	-	17,20,20	1.46	3 (17%)	23,28,28	2.98	9 (39%)
3	99S	A	502	-	27,36,36	2.98	4 (14%)	25,54,54	2.62	5 (20%)
2	PLG	B	501	-	17,20,20	1.51	3 (17%)	23,28,28	2.91	10 (43%)
3	99S	B	502	-	27,36,36	2.94	4 (14%)	25,54,54	2.75	9 (36%)
2	PLG	C	501	-	17,20,20	1.36	1 (5%)	23,28,28	3.00	11 (47%)
3	99S	C	502	-	27,36,36	3.10	5 (18%)	25,54,54	2.41	8 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLG	A	501	-	-	0/10/12/12	0/1/1/1
3	99S	A	502	-	-	0/22/46/46	0/4/4/4
2	PLG	B	501	-	-	0/10/12/12	0/1/1/1
3	99S	B	502	-	-	0/22/46/46	0/4/4/4
2	PLG	C	501	-	-	0/10/12/12	0/1/1/1
3	99S	C	502	-	-	0/22/46/46	0/4/4/4

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	502	99S	C8-C3	-11.53	1.27	1.42
3	A	502	99S	C8-C3	-11.45	1.27	1.42
3	B	502	99S	C8-C3	-10.48	1.28	1.42
3	B	502	99S	C19-C23	-6.85	1.27	1.44
3	A	502	99S	C19-C23	-6.55	1.28	1.44

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	99S	C3-C2-N7	-8.49	120.83	127.72
3	B	502	99S	C3-C2-N7	-8.08	121.17	127.72
3	C	502	99S	C3-C2-N7	-7.44	121.68	127.72
3	A	502	99S	C33-O32-C30	-3.37	108.91	115.84
3	B	502	99S	C18-C19-C23	-3.04	115.66	119.51

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	PLG	3	0
3	A	502	99S	4	0
3	B	502	99S	4	0
2	C	501	PLG	2	0
3	C	502	99S	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/442 (100%)	0.30	10 (2%) 64 64	22, 31, 41, 51	0
1	B	442/442 (100%)	0.38	12 (2%) 58 58	23, 32, 42, 49	0
1	C	442/442 (100%)	0.39	21 (4%) 34 33	19, 32, 48, 64	0
All	All	1326/1326 (100%)	0.36	43 (3%) 51 51	19, 32, 44, 64	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	364	CYS	4.9
1	B	421	GLY	4.6
1	A	421	GLY	4.2
1	B	416	LYS	3.5
1	C	259	GLN	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	99S	A	502	33/33	0.88	0.24	1.29	37,39,43,46	0
3	99S	C	502	33/33	0.88	0.24	0.51	34,39,46,47	0
3	99S	B	502	33/33	0.90	0.23	0.24	33,38,43,44	0
2	PLG	C	501	20/20	0.93	0.19	-0.24	22,23,24,25	0
2	PLG	A	501	20/20	0.95	0.16	-0.89	25,27,29,30	0
2	PLG	B	501	20/20	0.95	0.16	-1.16	22,24,25,26	0
4	CL	B	503	1/1	0.84	0.14	-3.78	33,33,33,33	0
4	CL	C	503	1/1	0.81	0.16	-	33,33,33,33	1

6.5 Other polymers [i](#)

There are no such residues in this entry.