



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:21 PM GMT

PDB ID : 3TOR
Title : Crystal structure of Escherichia coli NrfA with Europium bound
Authors : Lockwood, C.W.J.; Clarke, T.A.; Butt, J.N.; Hemmings, A.M.; Richardson, D.J.
Deposited on : 2011-09-06
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	452	 82% 13% ..
1	B	452	 82% 14% ..
1	C	452	 82% 13% ..
1	D	452	 81% 14% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 16846 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c nitrite reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	441	Total	C	N	O	S	0	1	0
			3481	2181	619	659	22			
1	B	441	Total	C	N	O	S	0	1	0
			3481	2181	619	659	22			
1	C	441	Total	C	N	O	S	0	1	0
			3481	2181	619	659	22			
1	D	441	Total	C	N	O	S	0	1	0
			3480	2180	619	659	22			


- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		
2	D	1	Total	Ca	0	0
			1	1		
2	C	1	Total	Ca	0	0
			1	1		

- Molecule 3 is EUROPIUM ION (three-letter code: EU) (formula: Eu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Eu	0	0
			3	3		
3	A	3	Total	Eu	0	0
			3	3		
3	D	1	Total	Eu	0	0
			1	1		
3	C	2	Total	Eu	0	0
			2	2		

- Molecule 4 is HEME C (three-letter code: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	524	Total	O	0	0
			524	524		
5	B	533	Total	O	0	0
			533	533		
5	C	522	Total	O	0	0
			522	522		
5	D	471	Total	O	0	0
			471	471		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cytochrome c nitrite reductase




- Molecule 1: Cytochrome c nitrite reductase




- Molecule 1: Cytochrome c nitrite reductase





● Molecule 1: Cytochrome c nitrite reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.88Å 90.49Å 292.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	146.10 – 2.00 19.97 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.2 (146.10-2.00) 98.3 (19.97-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.01Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.164 , 0.222 0.166 , 0.225	Depositor DCC
R_{free} test set	6985 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	18.1	Xtrriage
Anisotropy	0.102	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 59.4	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Outliers	6 of 139181 reflections (0.004%)	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16846	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.66 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.3308e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EU, CA, HEC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.05	2/3569 (0.1%)	0.88	3/4827 (0.1%)
1	B	1.10	4/3569 (0.1%)	0.90	4/4827 (0.1%)
1	C	1.07	4/3569 (0.1%)	0.88	1/4827 (0.0%)
1	D	5.90	8/3569 (0.2%)	2.17	9/4827 (0.2%)
All	All	3.09	18/14276 (0.1%)	1.33	17/19308 (0.1%)

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	396[A]	GLU	CD-OE2	213.94	3.60	1.25
1	D	396[B]	GLU	CD-OE2	213.94	3.60	1.25
1	D	396[A]	GLU	CD-OE1	107.80	2.44	1.25
1	D	396[B]	GLU	CD-OE1	107.80	2.44	1.25
1	D	396[A]	GLU	CG-CD	53.70	2.32	1.51

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	D	396[A]	GLU	OE1-CD-OE2	-80.74	26.41	123.30
1	D	396[B]	GLU	OE1-CD-OE2	-80.74	26.41	123.30
1	D	396[A]	GLU	CG-CD-OE2	-42.46	33.38	118.30
1	D	396[B]	GLU	CG-CD-OE2	-42.46	33.38	118.30
1	D	396[A]	GLU	CG-CD-OE1	-33.02	52.25	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3481	0	3371	80	0
1	B	3481	0	3371	87	0
1	C	3481	0	3371	82	0
1	D	3480	0	3371	88	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
3	C	2	0	0	0	0
3	D	1	0	0	0	0
4	A	215	0	155	32	0
4	B	215	0	155	35	0
4	C	215	0	155	31	0
4	D	215	0	155	33	0
5	A	524	0	0	20	1
5	B	533	0	0	16	1
5	C	522	0	0	13	0
5	D	471	0	0	14	0
All	All	16846	0	14104	351	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 351 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:190:GLU:CB	1:C:190:GLU:CG	1.75	1.55
1:B:37:THR:HA	5:B:1763:HOH:O	1.26	1.31
1:D:396[A]:GLU:OE2	1:D:396[A]:GLU:HG2	1.23	1.31
1:C:125:CYS:SG	4:C:3:HEC:CAC	2.21	1.28
1:B:285:CYS:SG	4:B:6:HEC:CAC	2.23	1.27

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:498:HOH:O	5:B:1700:HOH:O[4_445]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	440/452 (97%)	431 (98%)	9 (2%)	0	100	100
1	B	440/452 (97%)	428 (97%)	12 (3%)	0	100	100
1	C	440/452 (97%)	428 (97%)	11 (2%)	1 (0%)	52	48
1	D	440/452 (97%)	429 (98%)	11 (2%)	0	100	100
All	All	1760/1808 (97%)	1716 (98%)	43 (2%)	1 (0%)	56	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	264	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/370 (98%)	350 (96%)	13 (4%)	42	39
1	B	363/370 (98%)	354 (98%)	9 (2%)	55	55
1	C	363/370 (98%)	347 (96%)	16 (4%)	35	30

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	363/370 (98%)	348 (96%)	15 (4%)	37	32
All	All	1452/1480 (98%)	1399 (96%)	53 (4%)	42	38

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	52	GLN
1	C	274	ILE
1	D	396[A]	GLU
1	C	190	GLU
1	C	240	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 78 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	371	GLN
1	C	62	GLN
1	D	346	GLN
1	B	388	HIS
1	B	450	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 33 ligands modelled in this entry, 13 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	HEC	A	3	1,5	24,50,50	2.29	7 (29%)	19,82,82	3.27	10 (52%)
4	HEC	A	4	1	24,50,50	2.00	6 (25%)	19,82,82	2.94	11 (57%)
4	HEC	A	5	1,3	24,50,50	2.15	8 (33%)	19,82,82	3.08	8 (42%)
4	HEC	A	6	1,3	24,50,50	2.09	7 (29%)	19,82,82	2.83	6 (31%)
4	HEC	A	7	1,3	24,50,50	1.96	6 (25%)	19,82,82	2.74	8 (42%)
4	HEC	B	3	1,5	24,50,50	2.70	12 (50%)	19,82,82	3.01	6 (31%)
4	HEC	B	4	1	24,50,50	2.10	10 (41%)	19,82,82	2.69	4 (21%)
4	HEC	B	5	1,3	24,50,50	1.97	6 (25%)	19,82,82	3.64	9 (47%)
4	HEC	B	6	1,3	24,50,50	2.44	8 (33%)	19,82,82	2.57	7 (36%)
4	HEC	B	7	1,3	24,50,50	2.38	8 (33%)	19,82,82	2.79	7 (36%)
4	HEC	C	3	1,5	24,50,50	2.37	8 (33%)	19,82,82	2.59	8 (42%)
4	HEC	C	4	1	24,50,50	2.33	5 (20%)	19,82,82	2.73	7 (36%)
4	HEC	C	5	1,3	24,50,50	2.15	9 (37%)	19,82,82	3.10	11 (57%)
4	HEC	C	6	1,3	24,50,50	2.40	5 (20%)	19,82,82	3.41	10 (52%)
4	HEC	C	7	1,3	24,50,50	2.60	6 (25%)	19,82,82	2.77	8 (42%)
4	HEC	D	3	1,5	24,50,50	2.33	7 (29%)	19,82,82	2.70	4 (21%)
4	HEC	D	4	1	24,50,50	1.96	8 (33%)	19,82,82	3.36	8 (42%)
4	HEC	D	5	1,3	24,50,50	2.22	6 (25%)	19,82,82	3.25	12 (63%)
4	HEC	D	6	1,3	24,50,50	2.46	6 (25%)	19,82,82	3.18	7 (36%)
4	HEC	D	7	1,3	24,50,50	2.33	8 (33%)	19,82,82	2.74	7 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEC	A	3	1,5	-	0/6/54/54	0/0/8/8
4	HEC	A	4	1	-	0/6/54/54	0/0/8/8
4	HEC	A	5	1,3	-	0/6/54/54	0/0/8/8
4	HEC	A	6	1,3	-	0/6/54/54	0/0/8/8
4	HEC	A	7	1,3	-	0/6/54/54	0/0/8/8

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEC	B	3	1,5	-	0/6/54/54	0/0/8/8
4	HEC	B	4	1	-	0/6/54/54	0/0/8/8
4	HEC	B	5	1,3	-	0/6/54/54	0/0/8/8
4	HEC	B	6	1,3	-	0/6/54/54	0/0/8/8
4	HEC	B	7	1,3	-	0/6/54/54	0/0/8/8
4	HEC	C	3	1,5	-	0/6/54/54	0/0/8/8
4	HEC	C	4	1	-	0/6/54/54	0/0/8/8
4	HEC	C	5	1,3	-	0/6/54/54	0/0/8/8
4	HEC	C	6	1,3	-	0/6/54/54	0/0/8/8
4	HEC	C	7	1,3	-	0/6/54/54	0/0/8/8
4	HEC	D	3	1,5	-	0/6/54/54	0/0/8/8
4	HEC	D	4	1	-	0/6/54/54	0/0/8/8
4	HEC	D	5	1,3	-	0/6/54/54	0/0/8/8
4	HEC	D	6	1,3	-	0/6/54/54	0/0/8/8
4	HEC	D	7	1,3	-	0/6/54/54	0/0/8/8

The worst 5 of 146 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	7	HEC	C3B-C2B	-8.85	1.31	1.40
4	B	3	HEC	C3C-C2C	-7.82	1.32	1.40
4	D	6	HEC	C3C-C2C	-7.53	1.32	1.40
4	D	7	HEC	C3B-C2B	-7.07	1.33	1.40
4	C	6	HEC	C3C-C2C	-6.95	1.33	1.40

The worst 5 of 158 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	5	HEC	CBB-CAB-C3B	-9.71	105.78	127.35
4	D	4	HEC	CBB-CAB-C3B	-9.54	106.14	127.35
4	D	3	HEC	CBB-CAB-C3B	-8.92	107.53	127.35
4	D	6	HEC	CBB-CAB-C3B	-8.55	108.35	127.35
4	A	6	HEC	CBB-CAB-C3B	-8.35	108.78	127.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

20 monomers are involved in 127 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	3	HEC	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	4	HEC	5	0
4	A	5	HEC	5	0
4	A	6	HEC	9	0
4	A	7	HEC	7	0
4	B	3	HEC	5	0
4	B	4	HEC	8	0
4	B	5	HEC	5	0
4	B	6	HEC	7	0
4	B	7	HEC	12	0
4	C	3	HEC	5	0
4	C	4	HEC	6	0
4	C	5	HEC	3	0
4	C	6	HEC	7	0
4	C	7	HEC	10	0
4	D	3	HEC	6	0
4	D	4	HEC	5	0
4	D	5	HEC	5	0
4	D	6	HEC	8	0
4	D	7	HEC	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	441/452 (97%)	-0.50	3 (0%) 89 89	8, 16, 28, 39	0
1	B	441/452 (97%)	-0.46	5 (1%) 82 83	9, 17, 29, 40	0
1	C	441/452 (97%)	-0.45	3 (0%) 89 89	9, 18, 31, 42	0
1	D	441/452 (97%)	-0.33	4 (0%) 85 86	11, 20, 34, 43	0
All	All	1764/1808 (97%)	-0.43	15 (0%) 85 86	8, 18, 31, 43	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	37	THR	3.6
1	B	293	ALA	3.4
1	D	221	LYS	3.2
1	C	293	ALA	3.1
1	D	37	THR	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	HEC	C	6	43/43	0.97	0.11	1.66	5,13,17,25	0
4	HEC	D	6	43/43	0.98	0.10	1.53	9,14,20,31	0
4	HEC	B	7	43/43	0.98	0.11	0.77	10,15,31,40	0
4	HEC	C	7	43/43	0.98	0.11	0.65	10,14,31,39	0
4	HEC	A	6	43/43	0.98	0.09	0.59	6,10,16,24	0
4	HEC	A	7	43/43	0.98	0.11	0.55	12,16,33,44	0
4	HEC	B	6	43/43	0.98	0.09	0.51	5,10,18,25	0
4	HEC	D	7	43/43	0.98	0.11	0.34	11,15,35,41	0
4	HEC	A	4	43/43	0.98	0.10	0.18	8,14,16,21	0
4	HEC	B	3	43/43	0.98	0.08	0.16	4,10,14,15	0
4	HEC	B	4	43/43	0.99	0.09	-0.02	9,12,15,19	0
4	HEC	A	3	43/43	0.99	0.08	-0.04	3,8,12,14	0
4	HEC	C	4	43/43	0.98	0.10	-0.09	11,15,19,23	0
4	HEC	A	5	43/43	0.99	0.07	-0.11	5,10,12,13	0
4	HEC	B	5	43/43	0.99	0.08	-0.14	4,10,12,15	0
4	HEC	C	5	43/43	0.99	0.08	-0.23	4,11,14,16	0
4	HEC	D	4	43/43	0.97	0.10	-0.35	15,20,23,25	0
4	HEC	D	3	43/43	0.98	0.08	-0.43	9,15,18,19	0
4	HEC	D	5	43/43	0.98	0.08	-0.44	8,14,17,25	0
4	HEC	C	3	43/43	0.99	0.07	-0.88	8,13,16,17	0
3	EU	A	480	1/1	0.99	0.07	-1.02	24,24,24,24	1
2	CA	B	1	1/1	1.00	0.07	-1.29	13,13,13,13	0
3	EU	B	480	1/1	0.99	0.07	-1.29	25,25,25,25	1
2	CA	A	1	1/1	1.00	0.05	-2.39	12,12,12,12	0
2	CA	C	1	1/1	1.00	0.04	-2.99	13,13,13,13	0
2	CA	D	1	1/1	0.99	0.03	-3.14	19,19,19,19	0
3	EU	D	2	1/1	1.00	0.04	-	19,19,19,19	0
3	EU	A	479	1/1	1.00	0.07	-	16,16,16,16	1
3	EU	C	479	1/1	1.00	0.02	-	19,19,19,19	0
3	EU	A	2	1/1	1.00	0.03	-	14,14,14,14	0
3	EU	B	479	1/1	1.00	0.11	-	13,13,13,13	1
3	EU	B	2	1/1	1.00	0.03	-	15,15,15,15	0
3	EU	C	2	1/1	1.00	0.03	-	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.