



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:27 PM GMT

PDB ID : 1TRB
Title : CONVERGENT EVOLUTION OF SIMILAR FUNCTION IN TWO STRUCTURALLY DIVERGENT ENZYMES
Authors : Kuriyan, J.; Krishna, T.S.R.
Deposited on : 1991-09-07
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2611 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THIOREDOXIN REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	0	0
			2394	1491	424	469	10			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	138	SER	CYS	CONFLICT	UNP P0A9P4

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	164	Total 164	O 164	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: THIOREDOXIN REDUCTASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	123.10 Å 123.10 Å 81.40 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.00 32.35 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.00) 87.5 (32.35-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	15.18 (at 2.00 Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.177 , (Not available) 0.176 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	20.4	Xtriage
Anisotropy	0.282	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 83.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 21977 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2611	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.94	0/2433	1.68	43/3295 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	67	ARG	NE-CZ-NH2	-23.93	108.34	120.30
1	A	67	ARG	NE-CZ-NH1	16.02	128.31	120.30
1	A	311	ARG	NE-CZ-NH2	-14.12	113.24	120.30
1	A	311	ARG	NE-CZ-NH1	13.91	127.25	120.30
1	A	67	ARG	CB-CG-CD	-7.85	91.20	111.60
1	A	205	ARG	NE-CZ-NH2	-7.70	116.45	120.30
1	A	87	VAL	CG1-CB-CG2	7.25	122.49	110.90
1	A	189	ARG	NE-CZ-NH2	7.23	123.91	120.30
1	A	251	ILE	CA-CB-CG2	7.21	125.31	110.90
1	A	197	GLY	CA-C-N	-7.00	101.79	117.20
1	A	95	ARG	NE-CZ-NH1	7.00	123.80	120.30
1	A	117	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	A	251	ILE	CA-CB-CG1	-6.73	98.22	111.00
1	A	52	TRP	CD1-CG-CD2	6.71	111.67	106.30
1	A	52	TRP	CE2-CD2-CG	-6.64	101.99	107.30
1	A	293	ARG	NE-CZ-NH2	-6.63	116.98	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	65	MET	CG-SD-CE	6.35	110.36	100.20
1	A	291	ILE	CG1-CB-CG2	-6.26	97.62	111.40
1	A	198	ASN	N-CA-C	6.17	127.65	111.00
1	A	148	VAL	CA-CB-CG2	-6.15	101.68	110.90
1	A	50	GLU	CA-CB-CG	6.11	126.83	113.40
1	A	221	ARG	NE-CZ-NH2	-6.03	117.28	120.30
1	A	314	ASP	CB-CG-OD1	5.94	123.64	118.30
1	A	312	TYR	CB-CG-CD2	-5.87	117.48	121.00
1	A	189	ARG	NH1-CZ-NH2	-5.86	112.96	119.40
1	A	220	VAL	CG1-CB-CG2	5.80	120.19	110.90
1	A	148	VAL	CG1-CB-CG2	5.75	120.11	110.90
1	A	232	GLU	CA-CB-CG	5.71	125.95	113.40
1	A	55	ASP	CB-CG-OD2	5.70	123.43	118.30
1	A	22	VAL	CG1-CB-CG2	5.64	119.93	110.90
1	A	293	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	A	130	ARG	NE-CZ-NH1	5.57	123.08	120.30
1	A	229	ASP	CA-C-N	-5.47	105.15	117.20
1	A	224	ASP	CA-CB-CG	5.42	125.31	113.40
1	A	197	GLY	O-C-N	5.40	131.35	122.70
1	A	286	ASP	CB-CG-OD1	5.37	123.14	118.30
1	A	312	TYR	CD1-CG-CD2	5.34	123.78	117.90
1	A	265	VAL	CA-CB-CG1	-5.32	102.92	110.90
1	A	251	ILE	N-CA-CB	-5.25	98.72	110.80
1	A	282	PHE	CB-CG-CD2	-5.22	117.14	120.80
1	A	304	MET	CG-SD-CE	5.22	108.56	100.20
1	A	251	ILE	CB-CG1-CD1	-5.22	99.29	113.90
1	A	43	LEU	CA-CB-CG	5.00	126.81	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	311	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2394	0	2356	41	0
2	A	53	0	31	0	0
3	A	164	0	0	6	0
All	All	2611	0	2387	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:184:LYS:HD2	1:A:185:ILE:H	1.61	0.66
1:A:87:VAL:HG13	1:A:89:LEU:HD13	1.79	0.64
1:A:184:LYS:HD2	1:A:185:ILE:N	2.13	0.63
1:A:272:ASN:H	1:A:311:ARG:HH22	1.46	0.62
1:A:183:GLU:HB3	1:A:184:LYS:HZ2	1.65	0.62
1:A:2:THR:HG22	3:A:644:HOH:O	1.99	0.61
1:A:85:ASN:HA	1:A:251:ILE:HD11	1.82	0.61
1:A:162:LEU:HG	1:A:189:ARG:NH1	2.15	0.61
1:A:155:ASN:HD22	1:A:186:LEU:HD22	1.66	0.61
1:A:183:GLU:HB3	1:A:184:LYS:NZ	2.15	0.60
1:A:153:GLY:O	1:A:181:ARG:HG3	2.02	0.59
1:A:184:LYS:HG2	3:A:724:HOH:O	2.04	0.58
1:A:162:LEU:HG	1:A:189:ARG:CZ	2.33	0.58
1:A:155:ASN:ND2	1:A:186:LEU:HD22	2.19	0.57
1:A:87:VAL:HG22	1:A:96:LEU:HG	1.87	0.57
1:A:207:LEU:HD11	1:A:220:VAL:HG22	1.88	0.55
1:A:245:HIS:HE1	3:A:653:HOH:O	1.90	0.55
1:A:42:GLN:HG2	3:A:712:HOH:O	2.06	0.54
1:A:3:THR:HG23	1:A:102:GLU:HG2	1.89	0.54
1:A:223:ARG:NH1	1:A:224:ASP:HB2	2.24	0.53
1:A:204:ASN:HB3	1:A:225:THR:HA	1.90	0.53
1:A:189:ARG:HH11	1:A:193:LYS:HE3	1.74	0.52
1:A:6:SER:O	1:A:105:CYS:HA	2.09	0.52
1:A:223:ARG:O	1:A:224:ASP:HB3	2.09	0.52
1:A:208:GLU:HG3	1:A:223:ARG:HG2	1.93	0.51
1:A:158:VAL:O	1:A:162:LEU:HB2	2.10	0.51
1:A:174:ILE:HD13	1:A:222:LEU:HD11	1.95	0.48
1:A:158:VAL:HG12	1:A:162:LEU:HD22	1.97	0.46
1:A:59:LEU:HD13	1:A:67:ARG:NH2	2.29	0.46
1:A:220:VAL:HG13	1:A:222:LEU:HD13	1.96	0.46
1:A:220:VAL:HG12	1:A:234:LEU:HB2	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:PHE:O	1:A:104:THR:HA	2.16	0.46
1:A:125:GLU:HG2	3:A:674:HOH:O	2.17	0.45
1:A:39:LYS:NZ	1:A:81:PHE:HB3	2.34	0.43
1:A:162:LEU:CG	1:A:189:ARG:NH1	2.81	0.42
1:A:159:GLU:HG3	1:A:186:LEU:HD21	2.01	0.42
1:A:224:ASP:O	1:A:225:THR:HB	2.19	0.42
1:A:43:LEU:HB2	3:A:615:HOH:O	2.21	0.41
1:A:156:THR:O	1:A:160:GLU:HG2	2.21	0.40
1:A:162:LEU:HG	1:A:189:ARG:HD2	2.03	0.40
1:A:39:LYS:HZ1	1:A:81:PHE:HB3	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	314/320 (98%)	296 (94%)	16 (5%)	2 (1%)	30 22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	224	ASP
1	A	231	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	251/253 (99%)	223 (89%)	28 (11%)	7 4

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	11	LEU
1	A	22	VAL
1	A	43	LEU
1	A	64	LEU
1	A	87	VAL
1	A	89	LEU
1	A	148	VAL
1	A	162	LEU
1	A	164	LEU
1	A	173	LEU
1	A	181	ARG
1	A	184	LYS
1	A	189	ARG
1	A	196	ASN
1	A	198	ASN
1	A	201	LEU
1	A	207	LEU
1	A	215	MET
1	A	220	VAL
1	A	222	LEU
1	A	225	THR
1	A	230	ASN
1	A	234	LEU
1	A	236	VAL
1	A	241	VAL
1	A	251	ILE
1	A	258	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	155	ASN
1	A	166	ASN
1	A	245	HIS
1	A	248	ASN
1	A	266	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	275	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	500	-	48,58,58	1.47	6 (12%)	54,89,89	2.50	9 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	500	-	-	0/30/50/50	0/6/6/6

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	FAD	C5A-C4A	-2.71	1.34	1.40
2	A	500	FAD	C10-N10	-2.68	1.36	1.39
2	A	500	FAD	C10-N1	-2.30	1.31	1.35
2	A	500	FAD	C4X-C10	2.08	1.44	1.41
2	A	500	FAD	O4B-C1B	3.37	1.45	1.41
2	A	500	FAD	C4-C4X	5.30	1.51	1.41

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	FAD	C4X-C10-N10	-8.36	115.59	120.52
2	A	500	FAD	C4X-C4-N3	-7.34	113.55	123.59
2	A	500	FAD	C4-C4X-C10	-3.86	117.47	119.94
2	A	500	FAD	C1B-N9A-C4A	-3.22	122.08	126.94
2	A	500	FAD	N3A-C2A-N1A	-2.81	126.74	128.89
2	A	500	FAD	C4B-O4B-C1B	-2.75	106.70	109.72
2	A	500	FAD	C4A-C5A-N7A	2.18	111.48	109.48
2	A	500	FAD	C4X-N5-C5X	2.73	119.90	116.76
2	A	500	FAD	C4-N3-C2	11.16	124.89	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/320 (98%)	-0.26	14 (4%) 38 39	8, 21, 55, 96	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	227	ASN	11.8
1	A	226	GLN	6.5
1	A	225	THR	6.4
1	A	228	SER	4.7
1	A	189	ARG	4.0
1	A	231	ILE	3.6
1	A	196	ASN	3.5
1	A	223	ARG	3.2
1	A	178	ASP	3.1
1	A	214	GLN	2.6
1	A	215	MET	2.5
1	A	230	ASN	2.4
1	A	100	ASN	2.4
1	A	181	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FAD	A	500	53/53	0.98	0.07	-0.82	8,12,20,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.