



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:22 PM GMT

PDB ID : 3TSA  
Title : Spinosyn Rhamnosyltransferase SpnG  
Authors : Isiorho, E.A.; Liu, H.-W.; Keatinge-Clay, A.T.  
Deposited on : 2011-09-12  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	391	 81% 12% • 5%
1	B	391	 82% 11% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GLC	B	391	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NDP-rhamnosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	372	Total	C	N	O	S	0	0	0
			2772	1760	487	510	15			
1	B	372	Total	C	N	O	S	0	0	0
			2770	1758	488	509	15			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	EXPRESSION TAG	UNP Q9ALM8
A	360	ALA	SER	CONFLICT	UNP Q9ALM8
B	0	HIS	-	EXPRESSION TAG	UNP Q9ALM8
B	360	ALA	SER	CONFLICT	UNP Q9ALM8

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is SUGAR (ALPHA-D-GLUCOSE) (three-letter code: GLC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			12	6	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	291	Total	O	0	0
			291	291		
4	B	287	Total	O	0	0
			287	287		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: NDP-rhamnosyltransferase



#### • Molecule 1: NDP-rhamnosyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.16Å 57.06Å 68.11Å 81.46° 73.84° 85.95°	Depositor
Resolution (Å)	19.37 – 1.70 19.37 – 1.70	Depositor EDS
% Data completeness (in resolution range)	89.5 (19.37-1.70) 87.0 (19.37-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.86 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.181 , 0.231 0.180 , 0.229	Depositor DCC
$R_{free}$ test set	3801 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.6	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 44.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 76104 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6133	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.15	5/2842 (0.2%)	1.10	10/3901 (0.3%)
1	B	1.14	8/2839 (0.3%)	1.08	11/3895 (0.3%)
All	All	1.15	13/5681 (0.2%)	1.09	21/7796 (0.3%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	149	GLY	N-CA	10.95	1.62	1.46
1	A	31	VAL	CB-CG2	6.03	1.65	1.52
1	B	3	VAL	CB-CG1	5.42	1.64	1.52
1	A	41	ALA	CA-CB	5.32	1.63	1.52
1	B	162	VAL	CB-CG2	5.27	1.64	1.52
1	B	314	TYR	CD1-CE1	5.26	1.47	1.39
1	B	322	ARG	CG-CD	5.25	1.65	1.51
1	A	383	GLU	CB-CG	5.19	1.62	1.52
1	A	89	GLN	CG-CD	5.18	1.62	1.51
1	B	151	PHE	CD1-CE1	5.17	1.49	1.39
1	B	198	TYR	CD1-CE1	5.12	1.47	1.39
1	B	224	MET	CB-CG	5.07	1.67	1.51
1	B	320	TYR	CG-CD2	5.06	1.45	1.39

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	148	ALA	C-N-CA	-14.87	91.08	122.30
1	A	372	MET	CG-SD-CE	-9.06	85.71	100.20
1	A	120	ASP	CB-CG-OD1	8.97	126.37	118.30
1	B	224	MET	CA-CB-CG	7.53	126.10	113.30
1	B	141	ARG	NE-CZ-NH2	-7.30	116.65	120.30
1	A	148	ALA	O-C-N	-6.67	111.86	123.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	17	MET	CG-SD-CE	-6.25	90.19	100.20
1	B	372	MET	CG-SD-CE	-5.85	90.84	100.20
1	B	4	LEU	CB-CG-CD1	5.82	120.90	111.00
1	B	333	LEU	CB-CG-CD2	-5.62	101.45	111.00
1	B	319	ASP	CB-CG-OD2	-5.49	113.36	118.30
1	B	256	VAL	CB-CA-C	-5.48	100.99	111.40
1	A	314	TYR	CB-CG-CD2	-5.43	117.74	121.00
1	A	154	ARG	NE-CZ-NH2	-5.42	117.59	120.30
1	A	367	ASP	CB-CG-OD1	5.19	122.97	118.30
1	A	314	TYR	CB-CG-CD1	5.14	124.09	121.00
1	A	160	ASP	CB-CG-OD1	5.14	122.93	118.30
1	B	284	ARG	NE-CZ-NH2	-5.11	117.74	120.30
1	A	120	ASP	CB-CG-OD2	-5.08	113.73	118.30
1	B	160	ASP	CB-CG-OD1	5.07	122.86	118.30
1	B	198	TYR	CB-CG-CD1	-5.02	117.99	121.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2772	0	2767	42	0
1	B	2770	0	2769	25	0
2	A	1	0	0	0	0
3	B	12	0	11	0	0
4	A	291	0	0	15	0
4	B	287	0	0	8	0
All	All	6133	0	5547	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:ARG:HD2	4:A:599:HOH:O	1.46	1.16
1:A:107:ARG:HD3	4:A:485:HOH:O	1.66	0.95
1:A:309:LEU:HG	4:A:471:HOH:O	1.72	0.87
1:A:107:ARG:CD	4:A:485:HOH:O	2.27	0.81
1:A:69:ARG:CG	1:A:69:ARG:HH11	1.94	0.80
1:A:79:ASP:HB2	4:A:653:HOH:O	1.81	0.80
1:A:69:ARG:HG2	1:A:69:ARG:HH11	1.48	0.76
1:A:374:HIS:HD2	4:A:568:HOH:O	1.72	0.73
1:B:256:VAL:HG23	4:B:513:HOH:O	1.90	0.71
1:A:374:HIS:HE1	4:A:575:HOH:O	1.72	0.71
1:B:10:TYR:CD1	1:B:227:MET:HE1	2.27	0.69
1:B:101:GLN:HG3	1:B:105:TYR:CE1	2.28	0.68
1:A:69:ARG:HG2	1:A:69:ARG:NH1	2.04	0.67
1:B:244:ALA:HB1	4:B:692:HOH:O	1.95	0.67
1:B:160:ASP:N	1:B:161:PRO:HD2	2.11	0.66
1:B:220:VAL:HB	4:B:692:HOH:O	1.95	0.65
1:A:374:HIS:CD2	4:A:568:HOH:O	2.50	0.62
1:B:275:GLU:HG3	4:B:513:HOH:O	2.00	0.60
1:B:125:ILE:HG22	4:B:620:HOH:O	2.01	0.60
1:A:113:ARG:CD	4:A:599:HOH:O	2.23	0.59
1:A:0:HIS:HE1	1:A:30:GLU:OE1	1.85	0.59
1:B:283:LEU:HD11	1:B:289:VAL:HG22	1.85	0.58
1:B:284:ARG:HG2	4:B:685:HOH:O	2.04	0.57
1:B:243:ALA:HA	1:B:346:THR:HG22	1.86	0.57
1:A:69:ARG:HG3	4:A:664:HOH:O	2.04	0.56
1:A:269:ASP:N	1:A:269:ASP:OD2	2.39	0.55
1:A:283:LEU:HD11	1:A:289:VAL:HG22	1.89	0.55
1:B:101:GLN:HG3	1:B:105:TYR:HE1	1.69	0.55
1:B:342:HIS:O	1:B:346:THR:HG23	2.06	0.54
1:A:8:LEU:HD13	1:A:10:TYR:CE2	2.42	0.54
1:B:10:TYR:HD1	1:B:227:MET:HE1	1.70	0.53
1:A:247:LEU:HD12	1:A:248:PRO:HD2	1.90	0.53
1:A:7:PRO:HG2	1:A:39:LEU:CD1	2.38	0.53
1:A:267:LEU:HD13	1:A:271:ALA:HB3	1.91	0.53
1:A:22:TRP:HB3	1:B:22:TRP:HB3	1.92	0.52
1:A:288:LEU:HD13	1:A:352:VAL:HG11	1.93	0.50
1:A:374:HIS:CE1	1:B:28:GLY:HA3	2.47	0.49
1:A:380:ARG:HH21	1:A:380:ARG:HG2	1.77	0.49
1:A:246:GLU:HB3	4:A:421:HOH:O	2.13	0.48
1:B:10:TYR:CD1	1:B:227:MET:CE	2.96	0.48
1:B:160:ASP:N	1:B:161:PRO:CD	2.76	0.48
1:B:104:GLU:HG2	4:B:686:HOH:O	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:7:PRO:HG2	1:B:39:LEU:CD1	2.45	0.47
1:A:102:LEU:HB3	1:A:103:PRO:HD3	1.97	0.46
1:B:10:TYR:HD1	1:B:227:MET:CE	2.29	0.45
1:A:8:LEU:HD13	1:A:10:TYR:HE2	1.81	0.45
1:A:251:GLU:CG	4:A:571:HOH:O	2.64	0.45
1:A:162:VAL:O	1:A:166:HIS:HD2	1.99	0.45
1:A:251:GLU:CD	4:A:535:HOH:O	2.55	0.44
1:A:25:GLN:HE21	1:A:31:VAL:HB	1.82	0.44
1:A:68:LEU:O	1:A:69:ARG:HB2	2.18	0.44
1:A:215:THR:HG21	4:A:666:HOH:O	2.18	0.43
1:A:160:ASP:N	1:A:161:PRO:CD	2.81	0.43
1:A:160:ASP:HB2	1:A:161:PRO:HD3	2.01	0.43
1:A:246:GLU:CB	4:A:421:HOH:O	2.66	0.42
1:A:8:LEU:HD13	1:A:10:TYR:CD2	2.55	0.42
1:A:279:LEU:HA	1:A:279:LEU:HD12	1.85	0.41
1:A:380:ARG:NH2	1:A:380:ARG:HG2	2.35	0.41
1:A:175:GLU:HG2	1:A:385:THR:HG23	2.03	0.41
1:B:99:LEU:HD12	1:B:99:LEU:HA	1.92	0.41
1:B:200:PRO:HB3	1:B:300:THR:OG1	2.21	0.41
1:B:7:PRO:HD2	4:B:606:HOH:O	2.19	0.41
1:A:369:ILE:HA	1:A:372:MET:HE3	2.03	0.40
1:B:164:ARG:HA	1:B:168:LEU:O	2.21	0.40
1:A:298:ALA:HB1	1:A:320:TYR:CZ	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	368/391 (94%)	361 (98%)	6 (2%)	1 (0%)	46	26
1	B	368/391 (94%)	362 (98%)	6 (2%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	736/782 (94%)	723 (98%)	12 (2%)	1 (0%)	56 35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	69	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	290/302 (96%)	279 (96%)	11 (4%)	40 17
1	B	289/302 (96%)	276 (96%)	13 (4%)	34 13
All	All	579/604 (96%)	555 (96%)	24 (4%)	37 15

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	LEU
1	A	69	ARG
1	A	89	GLN
1	A	175	GLU
1	A	191	PRO
1	A	267	LEU
1	A	269	ASP
1	A	278	PRO
1	A	308	GLN
1	A	314	TYR
1	A	345	PHE
1	B	69	ARG
1	B	89	GLN
1	B	99	LEU
1	B	101	GLN
1	B	134	ASP
1	B	146	PRO

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	184	SER
1	B	227	MET
1	B	256	VAL
1	B	276	SER
1	B	345	PHE
1	B	380	ARG
1	B	382	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	0	HIS
1	A	25	GLN
1	A	85	GLN
1	A	96	GLN
1	A	166	HIS
1	A	197	GLN
1	A	280	ASN
1	A	308	GLN
1	A	313	GLN
1	A	374	HIS
1	B	25	GLN
1	B	85	GLN
1	B	89	GLN
1	B	140	HIS
1	B	166	HIS
1	B	280	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	GLC	B	391	-	12,12,12	1.56	5 (41%)	17,17,17	3.36	11 (64%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	B	391	-	-	0/2/22/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	391	GLC	O5-C5	-2.27	1.38	1.44
3	B	391	GLC	O3-C3	-2.24	1.37	1.43
3	B	391	GLC	C4-C5	2.08	1.57	1.53
3	B	391	GLC	O6-C6	2.17	1.51	1.42
3	B	391	GLC	C6-C5	2.59	1.61	1.51

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	391	GLC	O5-C5-C6	-4.30	95.48	106.36
3	B	391	GLC	O3-C3-C2	-3.09	103.37	110.34
3	B	391	GLC	O4-C4-C3	-2.94	103.71	110.34
3	B	391	GLC	O4-C4-C5	2.42	115.66	109.24
3	B	391	GLC	C1-C2-C3	3.44	115.55	110.43
3	B	391	GLC	O6-C6-C5	3.63	123.34	111.33
3	B	391	GLC	O2-C2-C1	3.65	117.85	109.82
3	B	391	GLC	O1-C1-C2	3.71	119.15	109.21
3	B	391	GLC	O5-C5-C4	3.71	116.65	109.68

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	391	GLC	C3-C4-C5	4.44	117.93	110.20
3	B	391	GLC	O5-C1-C2	7.59	121.91	109.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	372/391 (95%)	-0.27	0 100 100	12, 18, 33, 40	0
1	B	372/391 (95%)	-0.24	7 (1%) 70 74	12, 19, 32, 46	0
All	All	744/782 (95%)	-0.26	7 (0%) 85 88	12, 19, 33, 46	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	55	GLY	4.5
1	B	68	LEU	3.8
1	B	214	ARG	2.9
1	B	269	ASP	2.8
1	B	380	ARG	2.5
1	B	227	MET	2.3
1	B	173	THR	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GLC	B	391	12/12	0.88	0.14	4.48	17,24,36,39	0
2	MG	A	391	1/1	0.99	0.02	-5.62	17,17,17,17	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.