



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 31, 2016 – 11:04 AM EDT

PDB ID : 5TS2
Title : Crystal structure of a phosphopantetheine adenylyltransferase (CoaD, PPAT) from *Pseudomonas aeruginosa* bound to dephospho coenzyme A
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2016-10-27
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028320
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028320

i

X-RAY DIFFRACTION

A.

 R_{free}

1	A	167	<div><div></div></div> 87%12%
---	---	-----	-------------------------------

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	COD	A	201	-	-	-	X
2	COD	B	201	-	-	-	X
2	COD	E	201	-	-	-	X
2	COD	F	201	-	-	-	X
3	CA	A	202	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7991 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphopantetheine adenylyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	165	Total	C	N	O	S	0	1	0
			1288	827	232	226	3			
1	B	158	Total	C	N	O	S	0	2	0
			1227	790	213	221	3			
1	C	165	Total	C	N	O	S	0	0	0
			1283	823	231	226	3			
1	D	159	Total	C	N	O	S	0	1	0
			1232	793	212	224	3			
1	E	165	Total	C	N	O	S	0	0	0
			1274	818	228	225	3			
1	F	159	Total	C	N	O	S	0	0	0
			1221	786	214	219	2			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP B7V2S6
A	-6	ALA	-	expression tag	UNP B7V2S6
A	-5	HIS	-	expression tag	UNP B7V2S6
A	-4	HIS	-	expression tag	UNP B7V2S6
A	-3	HIS	-	expression tag	UNP B7V2S6
A	-2	HIS	-	expression tag	UNP B7V2S6
A	-1	HIS	-	expression tag	UNP B7V2S6
A	0	HIS	-	expression tag	UNP B7V2S6
B	-7	MET	-	initiating methionine	UNP B7V2S6
B	-6	ALA	-	expression tag	UNP B7V2S6
B	-5	HIS	-	expression tag	UNP B7V2S6
B	-4	HIS	-	expression tag	UNP B7V2S6
B	-3	HIS	-	expression tag	UNP B7V2S6
B	-2	HIS	-	expression tag	UNP B7V2S6
B	-1	HIS	-	expression tag	UNP B7V2S6
B	0	HIS	-	expression tag	UNP B7V2S6
C	-7	MET	-	initiating methionine	UNP B7V2S6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-6	ALA	-	expression tag	UNP B7V2S6
C	-5	HIS	-	expression tag	UNP B7V2S6
C	-4	HIS	-	expression tag	UNP B7V2S6
C	-3	HIS	-	expression tag	UNP B7V2S6
C	-2	HIS	-	expression tag	UNP B7V2S6
C	-1	HIS	-	expression tag	UNP B7V2S6
C	0	HIS	-	expression tag	UNP B7V2S6
D	-7	MET	-	initiating methionine	UNP B7V2S6
D	-6	ALA	-	expression tag	UNP B7V2S6
D	-5	HIS	-	expression tag	UNP B7V2S6
D	-4	HIS	-	expression tag	UNP B7V2S6
D	-3	HIS	-	expression tag	UNP B7V2S6
D	-2	HIS	-	expression tag	UNP B7V2S6
D	-1	HIS	-	expression tag	UNP B7V2S6
D	0	HIS	-	expression tag	UNP B7V2S6
E	-7	MET	-	initiating methionine	UNP B7V2S6
E	-6	ALA	-	expression tag	UNP B7V2S6
E	-5	HIS	-	expression tag	UNP B7V2S6
E	-4	HIS	-	expression tag	UNP B7V2S6
E	-3	HIS	-	expression tag	UNP B7V2S6
E	-2	HIS	-	expression tag	UNP B7V2S6
E	-1	HIS	-	expression tag	UNP B7V2S6
E	0	HIS	-	expression tag	UNP B7V2S6
F	-7	MET	-	initiating methionine	UNP B7V2S6
F	-6	ALA	-	expression tag	UNP B7V2S6
F	-5	HIS	-	expression tag	UNP B7V2S6
F	-4	HIS	-	expression tag	UNP B7V2S6
F	-3	HIS	-	expression tag	UNP B7V2S6
F	-2	HIS	-	expression tag	UNP B7V2S6
F	-1	HIS	-	expression tag	UNP B7V2S6
F	0	HIS	-	expression tag	UNP B7V2S6

- Molecule 2 is DEPHOSPHO COENZYME A (three-letter code: COD) (formula: $C_{21}H_{35}N_7O_{13}P_2S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 44	C 21	N 7	O 13	P 2	S 1	0	0
2	B	1	Total 44	C 21	N 7	O 13	P 2	S 1	0	0
2	C	1	Total 44	C 21	N 7	O 13	P 2	S 1	0	0
2	D	1	Total 44	C 21	N 7	O 13	P 2	S 1	0	0
2	E	1	Total 44	C 21	N 7	O 13	P 2	S 1	0	0
2	F	1	Total 44	C 21	N 7	O 13	P 2	S 1	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	E	1	Total	Ca	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	2	Total	Cl	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	1	Total 1	Cl 1	0	0
4	B	2	Total 2	Cl 2	0	0
4	C	3	Total 3	Cl 3	0	0
4	A	1	Total 1	Cl 1	0	0
4	F	2	Total 2	Cl 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	36	Total 36	O 36	0	0
5	B	21	Total 21	O 21	0	0
5	C	28	Total 28	O 28	0	0
5	D	44	Total 44	O 44	0	0
5	E	30	Total 30	O 30	0	0
5	F	30	Total 30	O 30	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphopantetheine adenylyltransferase

Chain A: 




- Molecule 1: Phosphopantetheine adenylyltransferase

Chain B: 




- Molecule 1: Phosphopantetheine adenylyltransferase

Chain C: 




- Molecule 1: Phosphopantetheine adenylyltransferase

Chain D: 




- Molecule 1: Phosphopantetheine adenylyltransferase

Chain E: 



- Molecule 1: Phosphopantetheine adenylyltransferase

Chain F: 

MET	ALA	HIS	HIS	HIS	HIS	HIS	HIS	HO	R3	Y4	F28	P62	L73	K81	A82	L89	R90	M103	R106	F116	L117	S124	F125	I126	L130	S142	K158	ARG
-----	-----	-----	-----	-----	-----	-----	-----	----	----	----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	98.38Å 101.35Å 105.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.87 – 2.30 46.86 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.5 (46.87-2.30) 97.5 (46.86-2.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.78 (at 2.29Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.180 , 0.244 0.179 , 0.244	Depositor DCC
R_{free} test set	2159 reflections (4.65%)	DCC
Wilson B-factor (Å ²)	43.2	Xtriage
Anisotropy	0.277	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7991	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, COD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1325	0.55	0/1801
1	B	0.40	0/1261	0.54	0/1716
1	C	0.41	0/1316	0.54	0/1787
1	D	0.44	0/1262	0.60	0/1714
1	E	0.38	0/1307	0.55	0/1778
1	F	0.38	0/1249	0.55	0/1697
All	All	0.40	0/7720	0.55	0/10493

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1288	0	1259	12	0
1	B	1227	0	1208	8	0
1	C	1283	0	1256	15	0
1	D	1232	0	1219	12	0
1	E	1274	0	1237	9	0
1	F	1221	0	1203	16	0
2	A	44	0	33	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	44	0	33	5	0
2	C	44	0	32	4	0
2	D	44	0	32	3	0
2	E	44	0	32	3	0
2	F	44	0	32	2	0
3	A	1	0	0	0	0
3	E	1	0	0	0	0
4	A	1	0	0	0	0
4	B	2	0	0	0	0
4	C	3	0	0	0	0
4	D	2	0	0	0	0
4	E	1	0	0	0	0
4	F	2	0	0	0	0
5	A	36	0	0	0	0
5	B	21	0	0	0	0
5	C	28	0	0	0	0
5	D	44	0	0	0	0
5	E	30	0	0	0	0
5	F	30	0	0	2	0
All	All	7991	0	7576	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 67 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:ARG:NH2	1:A:80:GLN:O	2.19	0.76
1:C:-5:HIS:HB2	1:F:62:PRO:HG3	1.71	0.72
1:D:90:ARG:HD2	1:D:124:SER:O	1.99	0.61
1:E:90:ARG:HD2	1:E:124:SER:O	2.02	0.58
1:E:3:ARG:NH2	1:E:80:GLN:O	2.38	0.57

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	164/167 (98%)	164 (100%)	0	0	100	100
1	B	158/167 (95%)	157 (99%)	1 (1%)	0	100	100
1	C	163/167 (98%)	163 (100%)	0	0	100	100
1	D	158/167 (95%)	157 (99%)	1 (1%)	0	100	100
1	E	163/167 (98%)	162 (99%)	1 (1%)	0	100	100
1	F	157/167 (94%)	157 (100%)	0	0	100	100
All	All	963/1002 (96%)	960 (100%)	3 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/142 (94%)	132 (98%)	2 (2%)	72	85
1	B	128/142 (90%)	126 (98%)	2 (2%)	70	84
1	C	133/142 (94%)	129 (97%)	4 (3%)	48	65
1	D	129/142 (91%)	127 (98%)	2 (2%)	70	84
1	E	131/142 (92%)	129 (98%)	2 (2%)	72	85
1	F	126/142 (89%)	126 (100%)	0	100	100
All	All	781/852 (92%)	769 (98%)	12 (2%)	72	85

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	78	LYS
1	C	101	LEU
1	D	128	SER
1	C	41	LYS
1	D	121	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 13 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	COD	A	201	-	38,46,46	3.00	12 (31%)	41,68,68	5.33	6 (14%)
2	COD	B	201	-	38,46,46	2.92	14 (36%)	41,68,68	5.18	5 (12%)
2	COD	C	201	-	38,46,46	2.98	14 (36%)	41,68,68	5.32	5 (12%)
2	COD	D	201	-	38,46,46	2.97	12 (31%)	41,68,68	5.41	6 (14%)
2	COD	E	201	-	38,46,46	2.96	13 (34%)	41,68,68	5.30	6 (14%)
2	COD	F	201	-	38,46,46	2.97	14 (36%)	41,68,68	5.30	8 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COD	A	201	-	-	0/39/59/59	0/3/3/3
2	COD	B	201	-	-	0/39/59/59	0/3/3/3
2	COD	C	201	-	-	0/39/59/59	0/3/3/3
2	COD	D	201	-	-	0/39/59/59	0/3/3/3
2	COD	E	201	-	-	0/39/59/59	0/3/3/3
2	COD	F	201	-	-	0/39/59/59	0/3/3/3

The worst 5 of 79 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	201	COD	O17-C11	-8.10	1.29	1.41
2	A	201	COD	O17-C11	-8.08	1.29	1.41
2	C	201	COD	O17-C11	-7.97	1.29	1.41
2	F	201	COD	O17-C11	-7.68	1.30	1.41
2	E	201	COD	O17-C11	-7.51	1.30	1.41

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	201	COD	N7-C6-N1	-15.55	92.43	118.52
2	C	201	COD	N7-C6-N1	-15.22	92.98	118.52
2	D	201	COD	N7-C6-N1	-15.11	93.16	118.52
2	F	201	COD	N7-C6-N1	-14.88	93.56	118.52
2	A	201	COD	N7-C6-N1	-14.76	93.74	118.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	COD	2	0
2	B	201	COD	5	0
2	C	201	COD	4	0
2	D	201	COD	3	0
2	E	201	COD	3	0
2	F	201	COD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	165/167 (98%)	-0.43	0 100 100	36, 49, 67, 79	0
1	B	158/167 (94%)	-0.38	0 100 100	34, 49, 69, 94	0
1	C	165/167 (98%)	-0.37	0 100 100	36, 51, 70, 86	0
1	D	159/167 (95%)	-0.39	0 100 100	26, 42, 65, 84	0
1	E	165/167 (98%)	-0.48	0 100 100	36, 51, 65, 92	0
1	F	159/167 (95%)	-0.40	0 100 100	32, 49, 78, 91	0
All	All	971/1002 (96%)	-0.41	0 100 100	26, 49, 70, 94	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CA	A	202	1/1	1.00	0.16	6.76	24,24,24,24	0
2	COD	A	201	44/44	0.85	0.23	3.58	36,62,93,106	44
2	COD	B	201	44/44	0.88	0.17	2.58	46,75,96,105	0
2	COD	E	201	44/44	0.89	0.18	2.43	30,60,93,95	44
2	COD	F	201	44/44	0.86	0.18	2.08	35,74,94,103	44
2	COD	C	201	44/44	0.92	0.15	1.75	33,66,99,103	44
2	COD	D	201	44/44	0.89	0.14	0.76	30,58,86,91	0
4	CL	D	203	1/1	0.98	0.12	0.31	61,61,61,61	0
4	CL	C	204	1/1	0.97	0.10	-0.87	66,66,66,66	0
4	CL	C	203	1/1	0.99	0.05	-	56,56,56,56	0
4	CL	B	203	1/1	0.94	0.18	-	64,64,64,64	0
4	CL	F	203	1/1	0.97	0.09	-	92,92,92,92	0
4	CL	F	202	1/1	0.85	0.19	-	80,80,80,80	0
4	CL	B	202	1/1	0.93	0.12	-	62,62,62,62	0
4	CL	E	203	1/1	0.95	0.08	-	62,62,62,62	0
3	CA	E	202	1/1	0.76	0.12	-	127,127,127,127	0
4	CL	C	202	1/1	0.97	0.17	-	67,67,67,67	0
4	CL	A	203	1/1	0.94	0.14	-	61,61,61,61	0
4	CL	D	202	1/1	0.95	0.15	-	65,65,65,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.