



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:23 PM GMT

PDB ID : 3TU5
Title : Actin complex with Gelsolin Segment 1 fused to Cobl segment
Authors : Kudryashov, D.S.; Sawaya, M.R.; Durer, Z.A.O.
Deposited on : 2011-09-15
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	
2	B	297	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CA	B	326	-	-	-	X
5	MPD	A	378	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4057 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Actin, alpha skeletal muscle.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	370	Total	C	N	O	S	0	0	0
			2890	1829	488	553	20			

- Molecule 2 is a protein called Gelsolin, Protein cordon-bleu, Thymosin beta-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	132	Total	C	N	O	S	0	0	0
			1053	678	178	196	1			


There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-17	MET	-	EXPRESSION TAG	UNP P06396
B	-16	ASN	-	EXPRESSION TAG	UNP P06396
B	-15	HIS	-	EXPRESSION TAG	UNP P06396
B	-14	LYS	-	EXPRESSION TAG	UNP P06396
B	-13	VAL	-	EXPRESSION TAG	UNP P06396
B	-12	HIS	-	EXPRESSION TAG	UNP P06396
B	-11	HIS	-	EXPRESSION TAG	UNP P06396
B	-10	HIS	-	EXPRESSION TAG	UNP P06396
B	-9	HIS	-	EXPRESSION TAG	UNP P06396
B	-8	HIS	-	EXPRESSION TAG	UNP P06396
B	-7	HIS	-	EXPRESSION TAG	UNP P06396
B	-6	ILE	-	EXPRESSION TAG	UNP P06396
B	-5	GLU	-	EXPRESSION TAG	UNP P06396
B	-4	GLY	-	EXPRESSION TAG	UNP P06396
B	-3	ARG	-	EXPRESSION TAG	UNP P06396
B	-2	HIS	-	EXPRESSION TAG	UNP P06396
B	0	GLY	-	EXPRESSION TAG	UNP P06396
B	1	SER	-	EXPRESSION TAG	UNP P06396
B	128	VAL	THR	CONFLICT	UNP Q5NBX1

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Ca 1 1	0	0
3	A	1	Total Ca 1 1	0	0

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $\text{C}_{10}\text{H}_{16}\text{N}_5\text{O}_{13}\text{P}_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $\text{C}_6\text{H}_{14}\text{O}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			8	6	2		
5	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	40	Total	O	0	0
			40	40		
6	B	25	Total	O	0	0
			25	25		

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	71.18Å 71.18Å 222.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.42 – 3.00 25.33 – 3.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (25.42-3.00) 96.6 (25.33-3.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.24 (at 2.99Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.165 , 0.204 0.163 , 0.203	Depositor DCC
R_{free} test set	668 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	65.8	Xtriage
Anisotropy	0.320	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 57.3	EDS
Estimated twinning fraction	0.032 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 13297 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4057	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, CA, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2952	0.67	0/4000
2	B	0.52	0/1078	0.66	0/1457
All	All	0.51	0/4030	0.67	0/5457

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2890	0	2863	39	0
2	B	1053	0	1033	7	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	31	0	12	1	0
5	A	16	0	28	1	0
6	A	40	0	0	1	0
6	B	25	0	0	0	0
All	All	4057	0	3936	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291:LYS:H	1:A:291:LYS:HE2	1.47	0.78
1:A:59:GLN:HG2	1:A:62:ARG:HH12	1.52	0.75
1:A:314:GLN:O	1:A:318:THR:HG23	1.90	0.72
2:B:67:ASN:H	2:B:99:GLN:NE2	1.88	0.72
2:B:72:ASP:HB3	2:B:125:LEU:HG	1.71	0.71
1:A:316:GLU:O	1:A:320:LEU:HD23	1.94	0.66
1:A:291:LYS:HE2	1:A:291:LYS:N	2.10	0.66
1:A:313:MET:O	1:A:317:ILE:HG23	1.95	0.66
1:A:241:GLU:HG3	1:A:247:VAL:HG22	1.77	0.65
1:A:352:PHE:HA	1:A:355:MET:HG3	1.83	0.59
1:A:287:ILE:HA	1:A:290:ARG:HG3	1.85	0.58
1:A:59:GLN:O	1:A:62:ARG:HB3	2.06	0.56
1:A:335:ARG:HA	1:A:338:SER:OG	2.07	0.54
1:A:354:GLN:HE21	2:B:4:GLU:H	1.56	0.54
2:B:54:ASN:OD1	2:B:56:ASN:HB2	2.07	0.54
1:A:205:GLU:HA	1:A:208:ILE:HD12	1.89	0.54
1:A:135:ALA:HB1	5:A:378:MPD:H13	1.90	0.54
1:A:13:GLY:HA3	4:A:377:ATP:O2B	2.08	0.53
1:A:153:LEU:HD22	1:A:313:MET:HE3	1.90	0.53
1:A:73:HIS:HB3	1:A:159:VAL:HG13	1.92	0.52
2:B:24:LYS:HD3	2:B:124:LYS:HE3	1.91	0.52
1:A:147:ARG:HD3	1:A:296:ASN:OD1	2.13	0.49
1:A:12:ASN:HD21	1:A:86:TRP:HE1	1.61	0.49
1:A:50:LYS:HE2	1:A:52:SER:O	2.13	0.47
1:A:259:GLU:OE2	1:A:312:ARG:NH1	2.47	0.47
1:A:183:ARG:HG2	1:A:206:ARG:HH21	1.80	0.47
1:A:105:LEU:HD12	1:A:132:MET:CE	2.45	0.47
1:A:297:ASN:HB2	1:A:328:LYS:O	2.15	0.46
1:A:190:MET:O	1:A:194:THR:HG23	2.17	0.45
1:A:366:GLY:O	1:A:369:ILE:HG22	2.17	0.44
1:A:88:HIS:ND1	1:A:92:ASN:OD1	2.51	0.43
1:A:273:GLY:O	1:A:277:THR:HG23	2.18	0.43
1:A:358:THR:OG1	1:A:361:GLU:HG3	2.19	0.43
1:A:180:LEU:HB2	1:A:269:MET:SD	2.58	0.43
1:A:12:ASN:ND2	6:A:405:HOH:O	2.52	0.42
1:A:38:PRO:HB2	1:A:41:GLN:HG3	2.00	0.42
2:B:24:LYS:HB3	2:B:124:LYS:HG3	2.00	0.42
1:A:205:GLU:HG2	1:A:205:GLU:H	1.66	0.42
1:A:220:ALA:O	1:A:312:ARG:HG3	2.19	0.42
1:A:220:ALA:CB	1:A:226:GLU:HG3	2.49	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:161:HIS:CD2	1:A:177:ARG:HB2	2.55	0.42
2:B:5:HIS:HB3	2:B:8:PHE:CD1	2.55	0.41
1:A:53:TYR:CD2	1:A:57:GLU:HG2	2.55	0.41
1:A:113:LYS:HD2	1:A:371:HIS:CE1	2.56	0.41
1:A:221:LEU:CD1	1:A:315:LYS:HD2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	368/377 (98%)	358 (97%)	9 (2%)	1 (0%)	46	84
2	B	130/297 (44%)	128 (98%)	2 (2%)	0	100	100
All	All	498/674 (74%)	486 (98%)	11 (2%)	1 (0%)	52	88

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	372	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/320 (98%)	275 (88%)	38 (12%)	6	25
2	B	110/244 (45%)	104 (94%)	6 (6%)	27	65
All	All	423/564 (75%)	379 (90%)	44 (10%)	9	32

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	GLU
1	A	47	MET
1	A	49	GLN
1	A	60	SER
1	A	62	ARG
1	A	77	THR
1	A	80	ASP
1	A	113	LYS
1	A	118	LYS
1	A	132	MET
1	A	147	ARG
1	A	159	VAL
1	A	185	LEU
1	A	193	LEU
1	A	199	SER
1	A	201	VAL
1	A	205	GLU
1	A	216	LEU
1	A	224	GLU
1	A	236	LEU
1	A	244	ASP
1	A	250	ILE
1	A	253	GLU
1	A	257	CYS
1	A	284	LYS
1	A	291	LYS
1	A	293	LEU
1	A	299	MET
1	A	309	ILE
1	A	312	ARG
1	A	317	ILE
1	A	318	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	320	LEU
1	A	334	GLU
1	A	359	LYS
1	A	368	SER
1	A	372	ARG
1	A	373	LYS
2	B	3	VAL
2	B	9	LEU
2	B	17	LEU
2	B	27	LEU
2	B	84	LEU
2	B	118	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	ASN
1	A	41	GLN
1	A	49	GLN
1	A	92	ASN
1	A	137	GLN
1	A	225	ASN
1	A	246	GLN
1	A	280	ASN
2	B	99	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	ATP	A	377	3	24,33,33	1.54	3 (12%)	31,52,52	2.28	11 (35%)
5	MPD	A	378	-	6,7,7	0.31	0	7,10,10	0.64	0
5	MPD	A	379	-	6,7,7	0.34	0	7,10,10	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	377	3	-	0/18/38/38	0/3/3/3
5	MPD	A	378	-	-	0/5/5/5	0/0/0/0
5	MPD	A	379	-	-	0/5/5/5	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	377	ATP	C2-N1	2.91	1.39	1.33
4	A	377	ATP	C2-N3	3.18	1.37	1.32
4	A	377	ATP	C4-N3	3.55	1.40	1.35

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	377	ATP	C2'-C1'-N9	-5.74	105.52	114.29
4	A	377	ATP	N3-C2-N1	-4.16	125.71	128.89
4	A	377	ATP	O5'-C5'-C4'	-2.56	99.69	109.12
4	A	377	ATP	C2'-C3'-C4'	-2.03	98.43	102.61
4	A	377	ATP	N6-C6-N1	2.06	123.61	119.20
4	A	377	ATP	O4'-C4'-C5'	2.11	116.88	109.32
4	A	377	ATP	C4'-O4'-C1'	2.24	112.18	109.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	377	ATP	O2B-PB-O3B	2.61	116.92	105.09
4	A	377	ATP	O3G-PG-O2G	3.27	119.82	107.38
4	A	377	ATP	O3A-PA-O5'	3.70	112.75	102.94
4	A	377	ATP	C4-C5-N7	5.49	114.53	109.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	377	ATP	1	0
5	A	378	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	370/377 (98%)	-0.46	8 (2%) 65 35	38, 71, 146, 177	0
2	B	132/297 (44%)	-0.64	2 (1%) 76 49	38, 59, 89, 122	0
All	All	502/674 (74%)	-0.51	10 (1%) 68 39	38, 66, 135, 177	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	48	GLY	4.5
1	A	49	GLN	2.8
2	B	1	SER	2.6
2	B	131	GLN	2.5
1	A	246	GLN	2.4
1	A	236	LEU	2.3
1	A	197	GLY	2.3
1	A	247	VAL	2.3
1	A	47	MET	2.1
1	A	244	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	MPD	A	378	8/8	0.93	0.23	3.38	66,68,70,70	0
3	CA	B	326	1/1	0.96	0.15	2.86	65,65,65,65	0
5	MPD	A	379	8/8	0.85	0.29	1.73	84,87,88,91	0
3	CA	A	376	1/1	0.99	0.20	0.77	53,53,53,53	0
4	ATP	A	377	31/31	0.98	0.12	-0.72	46,54,63,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.