



Full wwPDB X-ray Structure Validation Report i

Jan 31, 2016 – 10:29 PM GMT

PDB ID : 1TW2
Title : Crystal structure of Carminomycin-4-O-methyltransferase (DnrK) in complex with S-adenosyl-L-homocysteine (SAH) and 4-methoxy-e-rhodomycin T (M-ET)
Authors : Jansson, A.; Koskinen, H.; Mantsala, P.; Niemi, J.; Schneider, G.; Structural Proteomics in Europe (SPINE)
Deposited on : 2004-06-30
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ERT	B	1600	-	-	-	X

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Carminomycin 4-O-methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	338	Total	C 2610	N 1650	O 467	S 488	5	30	0
1	B	350	Total	C 2699	N 1705	O 484	S 505	5	3	0

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	CLONING ARTIFACT	UNP Q06528
A	-3	SER	-	CLONING ARTIFACT	UNP Q06528
A	-2	PRO	-	CLONING ARTIFACT	UNP Q06528
A	-1	ASN	-	CLONING ARTIFACT	UNP Q06528
A	0	SER	-	CLONING ARTIFACT	UNP Q06528
B	-4	GLY	-	CLONING ARTIFACT	UNP Q06528
B	-3	SER	-	CLONING ARTIFACT	UNP Q06528
B	-2	PRO	-	CLONING ARTIFACT	UNP Q06528
B	-1	ASN	-	CLONING ARTIFACT	UNP Q06528
B	0	SER	-	CLONING ARTIFACT	UNP Q06528

- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: C₁₄H₂₀N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	26	14	6	5	1	0	0
2	B	1	26	14	6	5	1	0	0

- Molecule 3 is METHYL (4R)-2-ETHYL-2,5,12-TRIHYDROXY-7-METHOXY-6,11-DIOX O-4-{{[2,3,6-TRIDEOXY-3-(DIMETHYLAMINO)-BETA-D-RIBO-HEXOPYRANOSYL] OXY}-1H,2H,3H,4H,6H,11H-TETRACENE-1-CARBOXYLATE (three-letter code: ERT) (formula: C₃₁H₃₇NO₁₁).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 43 31 1 11	0	0
3	B	1	Total C N O 43 31 1 11	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	97	Total O 97 97	0	0
4	B	68	Total O 68 68	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Carminomycin 4-O-methyltransferase



- Molecule 1: Carminomycin 4-O-methyltransferase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.68 Å 102.58 Å 124.26 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	54.00 – 2.50 54.53 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (54.00-2.50) 99.9 (54.53-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$< I/\sigma(I) >$ ¹	4.26 (at 2.51 Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R , R_{free}	0.203 , 0.271 0.213 , 0.272	Depositor DCC
R_{free} test set	1387 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.100	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 28.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 27543 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5612	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.37 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.0530e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ERT, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.62	2/2664 (0.1%)	0.92	15/3631 (0.4%)
1	B	0.53	0/2755	0.90	13/3757 (0.3%)
All	All	0.58	2/5419 (0.0%)	0.91	28/7388 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	283	GLU	CB-CG	-14.36	1.24	1.52
1	A	284	ARG	C-N	-6.63	1.18	1.34

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	283	GLU	N-CA-CB	-9.34	93.79	110.60
1	A	284	ARG	CA-CB-CG	7.67	130.28	113.40
1	A	97	ASP	CB-CG-OD2	6.98	124.58	118.30
1	B	168	ASP	CB-CG-OD2	6.96	124.56	118.30
1	B	107	ASP	CB-CG-OD2	6.91	124.52	118.30
1	B	312	ARG	NE-CZ-NH2	-6.90	116.85	120.30
1	B	318	ASP	CB-CG-OD2	6.53	124.17	118.30
1	B	97	ASP	CB-CG-OD2	6.46	124.12	118.30
1	B	312	ARG	NE-CZ-NH1	6.14	123.37	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	223	ASP	CB-CG-OD2	6.02	123.72	118.30
1	B	156	ASP	CB-CG-OD2	6.01	123.71	118.30
1	A	283	GLU	CA-CB-CG	5.76	126.07	113.40
1	B	14	ASP	CB-CG-OD2	5.55	123.29	118.30
1	A	342	ASP	CB-CG-OD2	5.54	123.29	118.30
1	B	124	ASP	CB-CG-OD2	5.50	123.25	118.30
1	A	107	ASP	CB-CG-OD2	5.46	123.21	118.30
1	B	96	ASP	CB-CG-OD2	5.45	123.21	118.30
1	A	261	ASP	CB-CG-OD2	5.33	123.10	118.30
1	A	246	ASP	CB-CG-OD2	5.33	123.10	118.30
1	A	318	ASP	CB-CG-OD2	5.27	123.05	118.30
1	A	164	ASP	CB-CG-OD2	5.21	122.99	118.30
1	A	228	ASP	CB-CG-OD2	5.21	122.98	118.30
1	B	231	ASP	CB-CG-OD2	5.20	122.98	118.30
1	B	236	ASP	CB-CG-OD2	5.20	122.98	118.30
1	A	162	ASP	CB-CG-OD2	5.19	122.97	118.30
1	A	14	ASP	CB-CG-OD2	5.14	122.92	118.30
1	A	300	ASP	CB-CG-OD2	5.03	122.83	118.30
1	B	332	ARG	NE-CZ-NH1	5.03	122.81	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	282	HIS	Mainchain
1	A	284	ARG	Mainchain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2610	0	2604	55	0
1	B	2699	0	2696	45	0
2	A	26	0	19	7	0
2	B	26	0	19	3	0
3	A	43	0	35	6	0
3	B	43	0	35	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	97	0	0	10	0
4	B	68	0	0	4	0
All	All	5612	0	5408	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:THR:HG21	4:A:725:HOH:O	1.42	1.20
4:A:704:HOH:O	1:B:122:LEU:HG	1.74	0.87
1:A:17:ARG:O	1:A:21:ARG:HG3	1.77	0.84
1:A:256:ASN:HA	1:A:303:MET:HE2	1.64	0.77
2:A:635:SAH:SD	3:A:600:ERT:C40	2.74	0.76
1:A:256:ASN:HA	1:A:303:MET:CE	2.16	0.75
1:B:118:SER:O	1:B:132:THR:HG21	1.87	0.74
1:B:29:MET:O	1:B:33:THR:HG23	1.90	0.71
1:A:184:ASP:OD2	4:A:718:HOH:O	2.09	0.71
1:A:109:THR:CG2	4:A:725:HOH:O	2.15	0.70
1:A:33:THR:HG21	1:B:122:LEU:HD12	1.74	0.69
1:A:14:ASP:OD2	1:A:17:ARG:HG3	1.93	0.68
1:A:159:LEU:HD11	3:A:600:ERT:H402	1.77	0.66
1:A:152:ARG:HH12	1:A:210:MET:CE	2.08	0.66
3:B:1600:ERT:O4'	3:B:1600:ERT:H7'3	1.96	0.65
1:A:29:MET:O	1:A:33:THR:HG23	1.96	0.65
1:A:152:ARG:HH12	1:A:210:MET:HE3	1.63	0.64
2:B:1635:SAH:SD	3:B:1600:ERT:C40	2.86	0.64
1:B:96:ASP:OD1	1:B:103:ARG:NH1	2.32	0.62
1:B:169:ALA:HB3	1:B:170:PRO:HD3	1.83	0.61
1:A:122:LEU:HB3	1:B:33:THR:HG21	1.82	0.61
1:A:118:SER:O	1:A:132:THR:HG21	2.01	0.61
1:B:330:GLU:OE2	4:B:1671:HOH:O	2.15	0.60
1:A:300:ASP:OD2	1:B:66:ARG:HD2	2.02	0.58
3:A:600:ERT:H142	3:A:600:ERT:C21	2.34	0.58
1:A:163:GLN:NE2	1:A:189:LYS:HB2	2.20	0.57
1:A:210:MET:HE3	2:A:635:SAH:O2'	2.03	0.57
1:B:127:ARG:HD2	4:B:1676:HOH:O	2.05	0.56
1:B:122:LEU:N	1:B:123:PRO:CD	2.69	0.56
1:B:142:TYR:CD2	1:B:210:MET:CE	2.90	0.55
1:A:107:ASP:OD2	1:A:109:THR:HB	2.07	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:189:LYS:NZ	4:B:1652:HOH:O	2.39	0.54
1:A:118:SER:HB3	1:A:305:VAL:HG13	1.90	0.54
1:B:303:MET:CE	1:B:307:LEU:CD1	2.85	0.54
1:A:66:ARG:HD2	1:B:300:ASP:OD2	2.08	0.54
1:A:79:GLU:HG3	1:A:85:PHE:CE1	2.43	0.53
1:A:159:LEU:CD1	3:A:600:ERT:H402	2.39	0.53
1:B:142:TYR:CD2	1:B:210:MET:HE3	2.43	0.53
1:A:15:ALA:HB3	4:A:695:HOH:O	2.08	0.53
1:A:313:THR:OG1	1:A:316:LYS:HG3	2.09	0.53
1:A:163:GLN:HG3	4:A:703:HOH:O	2.09	0.53
1:B:170:PRO:HG3	1:B:282:HIS:CE1	2.44	0.53
1:B:303:MET:CE	1:B:307:LEU:HD12	2.39	0.53
1:A:210:MET:CE	2:A:635:SAH:H2'	2.40	0.51
1:A:68:ILE:HG23	1:A:77:LEU:HD12	1.91	0.51
1:A:102:GLN:O	1:A:106:HIS:HD2	1.93	0.51
2:A:635:SAH:SD	3:A:600:ERT:H401	2.49	0.51
1:B:159:LEU:O	3:B:1600:ERT:H8'2	2.10	0.51
1:A:142:TYR:CD2	1:A:210:MET:HE1	2.47	0.50
1:A:162:ASP:OD1	1:A:338:THR:OG1	2.27	0.50
1:A:210:MET:HE2	2:A:635:SAH:H2'	1.93	0.50
1:A:118:SER:CB	1:A:305:VAL:HG13	2.42	0.50
1:A:65:LEU:HD21	1:B:288:HIS:HE1	1.77	0.49
2:B:1635:SAH:SD	3:B:1600:ERT:H403	2.53	0.49
2:A:635:SAH:SD	3:A:600:ERT:H402	2.55	0.47
1:B:152:ARG:HH12	1:B:210:MET:CE	2.28	0.47
1:A:300:ASP:OD2	1:B:66:ARG:CD	2.63	0.47
1:B:93:LEU:HD22	1:B:98:HIS:NE2	2.30	0.46
1:B:148:ARG:HG3	1:B:150:ASP:OD1	2.16	0.46
4:A:732:HOH:O	1:B:21:ARG:HD2	2.15	0.46
1:B:343:LEU:HD11	3:B:1600:ERT:H6'1	1.98	0.46
1:A:39:LEU:HD12	1:A:57:THR:HG21	1.98	0.46
1:B:303:MET:HE3	1:B:307:LEU:HD12	1.98	0.46
1:B:303:MET:CE	1:B:307:LEU:HD13	2.45	0.46
1:A:66:ARG:HD3	4:B:1687:HOH:O	2.16	0.46
1:A:98:HIS:CE1	1:A:99:PRO:HD2	2.51	0.45
1:A:283:GLU:HB3	1:A:284:ARG:H	1.51	0.45
1:B:199:ARG:HH11	1:B:199:ARG:HG2	1.81	0.45
1:A:27:THR:HA	1:A:70:HIS:HE1	1.81	0.45
1:A:148:ARG:NH1	1:A:150:ASP:OD2	2.50	0.44
1:A:210:MET:HE2	2:A:635:SAH:C8	2.48	0.44
1:B:14:ASP:O	1:B:17:ARG:HG2	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:PRO:HG3	1:B:33:THR:HB	1.99	0.44
1:A:189:LYS:HE2	1:A:216:THR:HG21	1.99	0.44
1:B:118:SER:HB3	1:B:305:VAL:HG13	2.00	0.43
1:B:26:HIS:O	1:B:30:VAL:HG23	2.18	0.43
1:B:298:GLU:O	1:B:302:ARG:HG3	2.18	0.43
1:A:33:THR:HB	1:B:123:PRO:HG3	2.01	0.42
1:A:122:LEU:HD13	1:B:33:THR:HG21	2.00	0.42
1:B:184:ASP:HA	1:B:249:ILE:HB	2.02	0.42
1:A:69:ARG:HD3	1:B:291:SER:OG	2.19	0.42
1:B:118:SER:CB	1:B:305:VAL:HG13	2.48	0.42
1:A:29:MET:O	1:A:33:THR:CG2	2.67	0.42
1:A:289:GLU:HB2	4:A:702:HOH:O	2.20	0.42
3:B:1600:ERT:H2'2	3:B:1600:ERT:H6'3	2.02	0.42
1:A:120:THR:HG21	1:B:120:THR:HG21	2.01	0.42
1:A:30:VAL:HG22	4:A:704:HOH:O	2.18	0.41
4:A:704:HOH:O	1:B:122:LEU:CG	2.50	0.41
1:A:297:THR:HG21	1:B:70:HIS:HA	2.02	0.41
1:B:210:MET:HE1	2:B:1635:SAH:C8	2.50	0.41
1:A:256:ASN:HD22	1:A:303:MET:HE2	1.86	0.41
1:B:130:ARG:CB	1:B:130:ARG:HH11	2.34	0.41
1:B:152:ARG:HH12	1:B:210:MET:HE3	1.86	0.41
1:A:52:ALA:O	1:A:56:ARG:HG3	2.21	0.41
1:A:98:HIS:ND1	1:A:99:PRO:HD2	2.37	0.40
1:A:284:ARG:HA	1:A:344:SER:OG	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	336/360 (93%)	324 (96%)	11 (3%)	1 (0%)	46 68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	348/360 (97%)	336 (97%)	10 (3%)	2 (1%)	30 50
All	All	684/720 (95%)	660 (96%)	21 (3%)	3 (0%)	39 61

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	161	CYS
1	A	163	GLN
1	B	283	GLU

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	270/285 (95%)	240 (89%)	30 (11%)	8 14
1	B	279/285 (98%)	243 (87%)	36 (13%)	5 10
All	All	549/570 (96%)	483 (88%)	66 (12%)	6 12

All (66) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	LEU
1	A	33	THR
1	A	53	LEU
1	A	58	ASP
1	A	80	ASP
1	A	92	GLU
1	A	103	ARG
1	A	109	THR
1	A	121	ARG
1	A	122	LEU
1	A	132	THR
1	A	148	ARG
1	A	154	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	159	LEU
1	A	183	LEU
1	A	192	PHE
1	A	216	THR
1	A	229	ARG
1	A	237	PHE
1	A	243	ARG
1	A	244	LYS
1	A	274	GLU
1	A	280	LEU
1	A	305	VAL
1	A	307	LEU
1	A	327	VAL
1	A	334	LEU
1	A	339	ILE
1	A	344	SER
1	A	345	LEU
1	B	9	ARG
1	B	12	GLN
1	B	24	SER
1	B	33	THR
1	B	48	ARG
1	B	51	LYS
1	B	60	ARG
1	B	92	GLU
1	B	103	ARG
1	B	109	THR
1	B	130	ARG
1	B	132	THR
1	B	148	ARG
1	B	161	CYS
1	B	162	ASP
1	B	183	LEU
1	B	192	PHE
1	B	199	ARG
1	B	216	THR
1	B	237	PHE
1	B	239	GLU
1	B	243	ARG
1	B	268	ARG
1	B	284	ARG
1	B	286	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	302	ARG
1	B	303	MET
1	B	305	VAL
1	B	314	ARG
1	B	315	GLU
1	B	316	LYS
1	B	334	LEU
1	B	336	SER
1	B	339	ILE
1	B	344	SER
1	B	345	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	HIS
1	A	70	HIS
1	A	102	GLN
1	A	106	HIS
1	A	163	GLN
1	A	181	HIS
1	A	256	ASN
1	A	282	HIS
1	B	12	GLN
1	B	70	HIS
1	B	102	GLN
1	B	106	HIS
1	B	282	HIS
1	B	288	HIS
1	B	290	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ERT	A	600	-	45,47,47	2.26	10 (22%)	61,73,73	2.84	15 (24%)
2	SAH	A	635	-	20,28,28	3.07	4 (20%)	19,40,40	5.55	8 (42%)
3	ERT	B	1600	-	45,47,47	2.24	8 (17%)	61,73,73	3.49	13 (21%)
2	SAH	B	1635	-	20,28,28	3.09	5 (25%)	19,40,40	5.42	11 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ERT	A	600	-	-	0/19/70/70	0/5/5/5
2	SAH	A	635	-	-	0/7/31/31	0/3/3/3
3	ERT	B	1600	-	-	0/19/70/70	0/5/5/5
2	SAH	B	1635	-	-	0/7/31/31	0/3/3/3

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	635	SAH	C5'-SD	-6.60	1.68	1.81
2	B	1635	SAH	C5'-SD	-6.19	1.68	1.81
3	B	1600	ERT	C8'-N3'	-5.25	1.26	1.46
3	A	600	ERT	O11-C11	-5.10	1.25	1.37
3	B	1600	ERT	C7'-N3'	-5.08	1.27	1.46
3	A	600	ERT	C8'-N3'	-5.05	1.27	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	600	ERT	C7'-N3'	-4.95	1.27	1.46
3	B	1600	ERT	O11-C11	-4.73	1.25	1.37
3	B	1600	ERT	C15-C12	-4.37	1.39	1.48
3	A	600	ERT	C15-C12	-4.13	1.39	1.48
2	A	635	SAH	CG-SD	-3.88	1.66	1.81
2	B	1635	SAH	CG-SD	-3.72	1.67	1.81
3	B	1600	ERT	C18-C12	-2.90	1.40	1.47
3	A	600	ERT	C18-C12	-2.85	1.40	1.47
3	A	600	ERT	O4-C4	-2.42	1.33	1.37
2	B	1635	SAH	C5-C4	-2.01	1.36	1.40
3	A	600	ERT	O12-C12	2.08	1.26	1.22
3	A	600	ERT	C17-C18	2.81	1.47	1.41
3	B	1600	ERT	C17-C18	2.82	1.47	1.41
3	A	600	ERT	O10-C21	3.97	1.43	1.33
3	B	1600	ERT	O10-C21	4.51	1.44	1.33
2	A	635	SAH	C2-N1	6.96	1.47	1.33
2	B	1635	SAH	C2-N1	7.47	1.48	1.33
2	B	1635	SAH	C2-N3	8.04	1.46	1.32
2	A	635	SAH	C2-N3	8.26	1.46	1.32
3	B	1600	ERT	O13-C21	8.62	1.43	1.21
3	A	600	ERT	O13-C21	8.66	1.43	1.21

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	635	SAH	N3-C2-N1	-21.40	112.51	128.89
2	B	1635	SAH	N3-C2-N1	-20.77	112.99	128.89
3	A	600	ERT	O13-C21-C10	-5.79	115.15	124.82
2	A	635	SAH	C4-C5-N7	-4.15	105.66	109.48
3	A	600	ERT	C2'-C3'-N3'	-4.10	103.56	115.70
2	B	1635	SAH	CB-CG-SD	-3.93	105.99	113.57
2	A	635	SAH	CB-CG-SD	-3.83	106.19	113.57
3	B	1600	ERT	O4-C4-C3	-3.50	118.47	124.35
3	A	600	ERT	O12-C12-C15	-3.38	115.92	120.89
2	B	1635	SAH	C4-C5-N7	-3.24	106.50	109.48
3	B	1600	ERT	O12-C12-C15	-3.15	116.25	120.89
3	B	1600	ERT	O13-C21-C10	-2.93	119.92	124.82
3	A	600	ERT	O4-C4-C3	-2.92	119.44	124.35
3	A	600	ERT	O10-C21-O13	-2.84	117.93	123.79
3	B	1600	ERT	C2'-C3'-N3'	-2.68	107.78	115.70
3	A	600	ERT	C13-C9-C8	-2.56	104.59	110.93
2	B	1635	SAH	C1'-N9-C4	-2.07	123.83	126.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1635	SAH	O3'-C3'-C4'	-2.02	104.99	111.05
2	B	1635	SAH	O3'-C3'-C2'	-2.02	105.26	111.83
2	A	635	SAH	C2'-C3'-C4'	2.17	107.07	102.61
3	B	1600	ERT	O5'-C5'-C4'	2.26	113.45	109.53
3	A	600	ERT	O4'-C4'-C3'	2.32	113.99	109.82
3	A	600	ERT	C15-C12-C18	2.40	122.09	117.99
3	B	1600	ERT	C1'-O5'-C5'	2.51	120.19	114.29
3	A	600	ERT	C4'-C3'-N3'	2.52	117.66	110.62
2	B	1635	SAH	C2'-C3'-C4'	2.62	108.00	102.61
2	A	635	SAH	O4'-C4'-C5'	2.62	116.00	108.85
3	A	600	ERT	C8'-N3'-C7'	2.70	118.97	110.43
3	A	600	ERT	O4-C4-C16	2.74	120.04	115.78
2	B	1635	SAH	O4'-C4'-C5'	2.88	116.70	108.85
3	B	1600	ERT	C15-C12-C18	3.14	123.36	117.99
3	B	1600	ERT	C8'-N3'-C7'	3.16	120.42	110.43
3	B	1600	ERT	C7'-N3'-C3'	3.19	122.31	113.09
3	A	600	ERT	C8'-N3'-C3'	3.81	124.12	113.09
2	B	1635	SAH	C2-N1-C6	3.85	125.64	118.77
3	B	1600	ERT	O4-C4-C16	3.94	121.91	115.78
3	B	1600	ERT	C40-O4-C4	4.29	124.04	117.54
2	A	635	SAH	C5'-SD-CG	4.36	115.48	102.41
2	A	635	SAH	O4'-C1'-N9	4.63	117.79	108.10
2	B	1635	SAH	C5'-SD-CG	4.64	116.33	102.41
2	A	635	SAH	C2-N1-C6	4.77	127.29	118.77
2	B	1635	SAH	O4'-C1'-N9	5.01	118.58	108.10
3	B	1600	ERT	O10-C21-C10	5.10	119.88	111.17
3	A	600	ERT	C40-O4-C4	6.79	127.84	117.54
3	A	600	ERT	O10-C21-C10	9.21	126.91	111.17
3	A	600	ERT	C20-C19-C10	14.18	119.12	113.15
3	B	1600	ERT	C20-C19-C10	23.81	123.18	113.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	600	ERT	6	0
2	A	635	SAH	7	0
3	B	1600	ERT	6	0
2	B	1635	SAH	3	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	338/360 (93%)	-0.28	1 (0%)	94 95	18, 26, 37, 49	7 (2%)
1	B	350/360 (97%)	-0.22	3 (0%)	85 88	16, 27, 40, 52	1 (0%)
All	All	688/720 (95%)	-0.25	4 (0%)	90 91	16, 26, 38, 52	8 (1%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	17	ARG	2.6
1	B	352	ALA	2.4
1	A	14	ASP	2.4
1	B	83	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i

There are no carbohydrates in this entry.

6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å²)	Q<0.9
3	ERT	B	1600	43/43	0.79	0.32	8.58	61,74,79,79	0
3	ERT	A	600	43/43	0.94	0.14	-0.04	20,24,31,36	0
2	SAH	A	635	26/26	0.97	0.12	-0.55	15,20,21,22	0
2	SAH	B	1635	26/26	0.97	0.11	-0.84	14,17,19,20	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.