wwPDB X-ray Structure Validation Summary Report (i)

Jul 21, 2016 – 09:24 PM EDT

PDB ID	:	4TW5
Title	:	Structure Of the First Two Thioredoxin Domains of Saccharomyces cerevisiae Eps1p
Authors	:	Fass, D.; Biran, S.
Deposited on	:	2014-06-30
Resolution	:	2.37 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027790
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027790

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4019 (2.40-2.36)
Clashscore	102246	4595 (2.40-2.36)
Ramachandran outliers	100387	4520 (2.40-2.36)
Sidechain outliers	100360	4522 (2.40-2.36)
RSRZ outliers	91569	4034 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	2% 79% 18% •
1	B	272	.% 86% 11% • •
1	C	272	.% 85% 13% •
1	D	272	2% 81% 15% • •

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 8819 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Eps1p.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	264	2151	1363	358	419	11	0	2	0
1	B	264	2145	1358	357	419	11	0	1	0
1	C	264	2145	1358	357	419	11	0	1	0
1	D	264	2138	1353	355	419	11	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	HIS	-	expression tag	UNP C7GJH6
A	25	MET	-	expression tag	UNP C7GJH6
A	26	LEU	-	expression tag	UNP C7GJH6
A	27	ASP	-	expression tag	UNP C7GJH6
B	24	HIS	-	expression tag	UNP C7GJH6
B	25	MET	-	expression tag	UNP C7GJH6
B	26	LEU	-	expression tag	UNP C7GJH6
B	27	ASP	-	expression tag	UNP C7GJH6
C	24	HIS	-	expression tag	UNP C7GJH6
C	25	MET	-	expression tag	UNP C7GJH6
C	26	LEU	-	expression tag	UNP C7GJH6
C	27	ASP	-	expression tag	UNP C7GJH6
D	24	HIS	-	expression tag	UNP C7GJH6
D	25	MET	-	expression tag	UNP C7GJH6
D	26	LEU	-	expression tag	UNP C7GJH6
D	27	ASP	-	expression tag	UNP C7GJH6

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	72	72	72	0	0
			Total	O		
2	B	52	52	52	0	0
			Total	O		
2	C	61	61	61	0	0
			Total	O		
2	D	55	55	55	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Eps1p

Chain A:



- Molecule 1: Eps1p

Chain B:



- Molecule 1: Eps1p

Chain C:



- Molecule 1: Eps1p

Chain D:




I170
A171
G172
L179
K198
R208
R215
Q216
L217
T224
C236
L239
I247
T248
R254
E255
P256
K257
L270
R289
R290
T291
F292
THR
ASN
SER

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	40.03Å 133.03Å 124.41Å 90.00° 92.00° 90.00°	Depositor
Resolution (Å)	41.77 – 2.37 41.77 – 2.37	Depositor EDS
% Data completeness (in resolution range)	99.4 (41.77-2.37) 95.4 (41.77-2.37)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.62 (at 2.37Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.211 , 0.272 0.211 , 0.272	Depositor DCC
R_{free} test set	3135 reflections (6.23%)	DCC
Wilson B-factor (Å^2)	33.1	Xtriage
Anisotropy	0.423	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 24.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.46$, $< L^2 > = 0.28$	Xtriage
Estimated twinning fraction	0.107 for h,-k,-l	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	8819	wwPDB-VP
Average B, all atoms (Å^2)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.03 % of the origin peak, indicating pseudo translational symmetry.The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.4022e-03.The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.49	0/2216	0.62	0/3004
1	B	0.45	0/2207	0.62	0/2993
1	C	0.52	0/2207	0.67	1/2993 (0.0%)
1	D	0.43	0/2196	0.62	1/2978 (0.0%)
All	All	0.47	0/8826	0.63	2/11968 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	208	ARG	NE-CZ-NH1	-5.93	117.33	120.30
1	C	60	CYS	CA-CB-SG	5.05	123.09	114.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2151	0	2042	40	0
1	B	2145	0	2029	24	0
1	C	2145	0	2029	21	1
1	D	2138	0	2022	35	0
2	A	72	0	0	12	0
2	B	52	0	0	7	0
2	C	61	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	55	0	0	10	0
All	All	8819	0	8122	116	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 116 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:237:GLU:OE2	2:B:301:HOH:O	1.90	0.89
1:C:249:ASN:ND2	2:C:301:HOH:O	2.14	0.78
1:B:275:ASN:O	2:B:322:HOH:O	2.08	0.72
1:C:57:SER:OG	2:C:333:HOH:O	2.07	0.71
1:C:220:ASP:OD2	1:C:289:ARG:HD2	1.91	0.71

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:198:LYS:NZ	1:C:264:ASN:OD1[1_655]	2.19	0.01

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/272 (97%)	258 (98%)	6 (2%)	0	100	100
1	B	263/272 (97%)	257 (98%)	6 (2%)	0	100	100
1	C	263/272 (97%)	260 (99%)	3 (1%)	0	100	100
1	D	262/272 (96%)	257 (98%)	5 (2%)	0	100	100
All	All	1052/1088 (97%)	1032 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/249 (98%)	243 (100%)	0	100	100
1	B	242/249 (97%)	240 (99%)	2 (1%)	86	94
1	C	242/249 (97%)	242 (100%)	0	100	100
1	D	241/249 (97%)	239 (99%)	2 (1%)	86	94
All	All	968/996 (97%)	964 (100%)	4 (0%)	93	98

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	43	LYS
1	B	62	HIS
1	D	128	ARG
1	D	170	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	51	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	264/272 (97%)	0.03	5 (1%)	70	72	19, 35, 67, 87	0
1	B	264/272 (97%)	0.07	4 (1%)	76	78	25, 41, 75, 91	0
1	C	264/272 (97%)	0.08	4 (1%)	76	78	17, 36, 75, 96	0
1	D	264/272 (97%)	0.07	5 (1%)	70	72	25, 41, 65, 81	0
All	All	1056/1088 (97%)	0.06	18 (1%)	73	75	17, 38, 72, 96	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	59	TYR	6.8
1	D	292	PHE	4.1
1	C	59	TYR	3.5
1	D	59	TYR	3.5
1	A	292	PHE	3.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.