



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 8, 2016 – 10:50 AM EST

PDB ID : 5U4Q
Title : 1.5 Angstrom Resolution Crystal Structure of NAD-Dependent Epimerase from *Klebsiella pneumoniae* in Complex with NAD.
Authors : Minasov, G.; Shuvalova, L.; Kiryukhina, O.; Dubrovskaya, I.; Bishop, B.; Kwon, K.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2016-12-05
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	
1	B	337	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5905 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called dTDP-glucose 4,6-dehydratase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	321	Total	C	N	O	S	Se	0	21	0
			2736	1750	460	509	1	16			
1	B	304	Total	C	N	O	S	Se	0	8	0
			2477	1584	416	465	1	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP A0A169YGX7
A	-1	ASN	-	expression tag	UNP A0A169YGX7
A	0	ALA	-	expression tag	UNP A0A169YGX7
B	-2	SER	-	expression tag	UNP A0A169YGX7
B	-1	ASN	-	expression tag	UNP A0A169YGX7
B	0	ALA	-	expression tag	UNP A0A169YGX7

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	358	Total	O	0	20
			370	370		
4	B	225	Total	O	0	12
			233	233		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.24Å 55.84Å 97.32Å 90.00° 93.21° 90.00°	Depositor
Resolution (Å)	29.21 – 1.50 29.21 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.4 (29.21-1.50) 99.5 (29.21-1.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.75 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.180 , 0.211 0.181 , 0.212	Depositor DCC
R_{free} test set	5033 reflections (5.19%)	DCC
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5905	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/2788	0.70	0/3750
1	B	0.48	0/2526	0.67	0/3410
All	All	0.51	0/5314	0.69	0/7160

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2736	0	2658	5	0
1	B	2477	0	2395	11	0
2	A	44	0	26	1	0
2	B	44	0	26	2	0
3	A	1	0	0	0	0
4	A	370	0	0	0	0
4	B	233	0	0	5	0
All	All	5905	0	5105	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:303:GLN:NE2	1:A:307:ASP:OD2	2.29	0.60
1:A:312[B]:LYS:O	1:A:312[B]:LYS:HG3	2.03	0.58
2:B:401:NAD:C5N	4:B:517:HOH:O	2.50	0.58
2:B:401:NAD:H5N	4:B:517:HOH:O	2.03	0.57
1:B:248:PRO:HB2	4:B:630:HOH:O	2.09	0.52
1:B:58[A]:GLN:NE2	4:B:501[A]:HOH:O	2.34	0.51
1:B:195:PHE:O	1:B:199:MSE:HB2	2.12	0.49
1:B:271:TYR:CE2	1:B:317:VAL:HG13	2.51	0.46
1:B:5:VAL:HG22	1:B:80:ILE:CG2	2.47	0.45
1:B:140:THR:HB	1:B:249:ALA:HB2	1.99	0.45
1:B:188:PRO:HA	1:B:193:PHE:CG	2.53	0.44
1:B:248:PRO:CB	4:B:630:HOH:O	2.65	0.44
1:B:67:MSE:HE1	1:B:109:ILE:HD13	2.01	0.43
1:A:5:VAL:HG22	1:A:80:ILE:CG2	2.48	0.43
1:A:98:TYR:O	1:A:102:ASN:HB2	2.18	0.43
1:B:67:MSE:HE1	1:B:109:ILE:CD1	2.49	0.42
1:B:46[B]:LEU:HA	1:B:46[B]:LEU:HD23	1.86	0.42
1:A:179:PHE:O	2:A:401:NAD:H4N	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	336/337 (100%)	320 (95%)	16 (5%)	0	100	100
1	B	304/337 (90%)	291 (96%)	13 (4%)	0	100	100
All	All	640/674 (95%)	611 (96%)	29 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/270 (107%)	288 (100%)	1 (0%)	94	86
1	B	261/270 (97%)	259 (99%)	2 (1%)	86	70
All	All	550/540 (102%)	547 (100%)	3 (0%)	90	81

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	193	PHE
1	B	247	SER
1	B	281	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	A	401	-	42,48,48	0.79	1 (2%)	46,73,73	1.73	5 (10%)
2	NAD	B	401	-	42,48,48	0.85	1 (2%)	46,73,73	1.41	5 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	401	-	-	0/22/62/62	0/5/5/5
2	NAD	B	401	-	-	0/22/62/62	0/5/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NAD	C3N-C7N	2.14	1.53	1.50
2	B	401	NAD	C5A-C4A	2.62	1.46	1.40

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NAD	N3A-C2A-N1A	-7.22	123.20	128.87
2	B	401	NAD	N3A-C2A-N1A	-5.95	124.20	128.87
2	A	401	NAD	O7N-C7N-N7N	-3.68	117.34	122.58
2	A	401	NAD	C1B-N9A-C4A	-2.51	124.01	126.81
2	B	401	NAD	C2A-N1A-C6A	2.15	122.60	118.77
2	B	401	NAD	C2N-C3N-C7N	2.17	125.48	119.24
2	B	401	NAD	N6A-C6A-N1A	2.23	122.26	118.52
2	A	401	NAD	C2A-N1A-C6A	2.77	123.72	118.77
2	B	401	NAD	C3N-C7N-N7N	3.30	121.56	117.82
2	A	401	NAD	C3N-C7N-N7N	4.76	123.21	117.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	NAD	1	0
2	B	401	NAD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	308/337 (91%)	0.36	25 (8%) 15 14	12, 19, 40, 50	0
1	B	294/337 (87%)	1.05	53 (18%) 2 2	14, 28, 69, 95	0
All	All	602/674 (89%)	0.69	78 (12%) 5 4	12, 23, 59, 95	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	265	PRO	9.8
1	B	88	VAL	8.2
1	B	306	TYR	7.6
1	B	89	ARG	6.9
1	B	271	TYR	6.6
1	B	92	LEU	6.2
1	B	268	LEU	6.1
1	B	280	GLY	5.9
1	B	266	VAL	5.6
1	B	314	GLN	5.6
1	B	333	GLN	5.6
1	B	273	THR	5.4
1	A	266	VAL	5.3
1	A	94	ASN	5.2
1	B	198	ALA	4.9
1	B	278	ALA	4.8
1	B	12	ILE	4.4
1	B	51	TYR	4.4
1	B	94	ASN	4.4
1	A	290	ILE	4.3
1	B	275	LEU	4.3
1	B	215	ARG	4.2
1	B	87	GLY	4.2
1	B	93	GLU	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	202	GLY	4.2
1	B	279	LEU	4.0
1	B	272	ILE	3.8
1	B	298	THR	3.7
1	A	36[A]	TYR	3.7
1	B	322	LYS	3.6
1	B	277	GLU	3.6
1	B	307	ASP	3.5
1	A	334	ILE	3.5
1	A	311[A]	PHE	3.3
1	A	289	PRO	3.3
1	B	200	LEU	3.3
1	A	298	THR	3.2
1	B	303[A]	GLN	3.1
1	B	270	ASP	3.1
1	A	86	ALA	3.0
1	B	267	GLU	2.9
1	B	262	ASN	2.9
1	A	205	ILE	2.9
1	A	95	PRO	2.9
1	B	0	ALA	2.8
1	A	204	SER	2.8
1	B	318	LYS	2.8
1	A	314	GLN	2.7
1	A	280	GLY	2.7
1	B	123	TYR	2.7
1	A	109	ILE	2.6
1	B	82	LEU	2.6
1	A	134[A]	ARG	2.6
1	B	263	SER	2.6
1	A	0	ALA	2.6
1	A	208	TYR	2.6
1	A	284	GLN	2.6
1	A	245	SER	2.4
1	B	312	LYS	2.4
1	B	239	ALA	2.4
1	B	311	PHE	2.4
1	A	82	LEU	2.4
1	B	79	VAL	2.3
1	B	330	ASP	2.3
1	B	261	GLY	2.3
1	B	319	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	195	PHE	2.3
1	A	265	PRO	2.3
1	A	96	TYR	2.2
1	B	50	ALA	2.1
1	A	176	LEU	2.1
1	B	247	SER	2.1
1	A	286	ASN	2.1
1	B	240	ASP	2.1
1	B	52	PRO	2.0
1	B	248	PRO	2.0
1	B	181	VAL	2.0
1	B	134	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CL	A	402	1/1	1.00	0.10	0.08	16,16,16,16	0
2	NAD	A	401	44/44	0.96	0.08	-0.78	15,17,22,24	0
2	NAD	B	401	44/44	0.96	0.08	-0.84	18,20,27,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.