



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 10:33 PM GMT

PDB ID : 1U5R
Title : Crystal Structure of the TAO2 Kinase Domain: Activation and Specifity of a Ste20p MAP3K
Authors : Zhou, T.; Raman, M.; Gao, Y.; Earnest, S.; Chen, Z.; Machius, M.; Cobb, M.H.; Goldsmith, E.J.
Deposited on : 2004-07-28
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	348	
1	B	348	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	SEP	B	181	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5431 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called serine/threonine protein kinase TAO2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	309	Total	C	N	O	P	S	0	0	0
			2490	1586	432	457	1	14			
1	B	309	Total	C	N	O	P	S	0	0	0
			2490	1586	432	457	1	14			

There are 58 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-27	MET	-	CLONING ARTIFACT	UNP Q9JLS3
A	-26	SER	-	CLONING ARTIFACT	UNP Q9JLS3
A	-25	TYR	-	CLONING ARTIFACT	UNP Q9JLS3
A	-24	TYR	-	CLONING ARTIFACT	UNP Q9JLS3
A	-23	HIS	-	EXPRESSION TAG	UNP Q9JLS3
A	-22	HIS	-	EXPRESSION TAG	UNP Q9JLS3
A	-21	HIS	-	EXPRESSION TAG	UNP Q9JLS3
A	-20	HIS	-	EXPRESSION TAG	UNP Q9JLS3
A	-19	HIS	-	EXPRESSION TAG	UNP Q9JLS3
A	-18	HIS	-	EXPRESSION TAG	UNP Q9JLS3
A	-17	ASP	-	CLONING ARTIFACT	UNP Q9JLS3
A	-16	TYR	-	CLONING ARTIFACT	UNP Q9JLS3
A	-15	ASP	-	CLONING ARTIFACT	UNP Q9JLS3
A	-14	ILE	-	CLONING ARTIFACT	UNP Q9JLS3
A	-13	PRO	-	CLONING ARTIFACT	UNP Q9JLS3
A	-12	THR	-	CLONING ARTIFACT	UNP Q9JLS3
A	-11	THR	-	CLONING ARTIFACT	UNP Q9JLS3
A	-10	GLU	-	CLONING ARTIFACT	UNP Q9JLS3
A	-9	ASN	-	CLONING ARTIFACT	UNP Q9JLS3
A	-8	LEU	-	CLONING ARTIFACT	UNP Q9JLS3
A	-7	TYR	-	CLONING ARTIFACT	UNP Q9JLS3
A	-6	PHE	-	CLONING ARTIFACT	UNP Q9JLS3
A	-5	GLN	-	CLONING ARTIFACT	UNP Q9JLS3
A	-4	GLY	-	CLONING ARTIFACT	UNP Q9JLS3
A	-3	ALA	-	CLONING ARTIFACT	UNP Q9JLS3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	CLONING ARTIFACT	UNP Q9JLS3
A	-1	ASP	-	CLONING ARTIFACT	UNP Q9JLS3
A	0	PRO	-	CLONING ARTIFACT	UNP Q9JLS3
A	181	SEP	SER	MODIFIED RESIDUE	UNP Q9JLS3
B	-27	MET	-	CLONING ARTIFACT	UNP Q9JLS3
B	-26	SER	-	CLONING ARTIFACT	UNP Q9JLS3
B	-25	TYR	-	CLONING ARTIFACT	UNP Q9JLS3
B	-24	TYR	-	CLONING ARTIFACT	UNP Q9JLS3
B	-23	HIS	-	EXPRESSION TAG	UNP Q9JLS3
B	-22	HIS	-	EXPRESSION TAG	UNP Q9JLS3
B	-21	HIS	-	EXPRESSION TAG	UNP Q9JLS3
B	-20	HIS	-	EXPRESSION TAG	UNP Q9JLS3
B	-19	HIS	-	EXPRESSION TAG	UNP Q9JLS3
B	-18	HIS	-	EXPRESSION TAG	UNP Q9JLS3
B	-17	ASP	-	CLONING ARTIFACT	UNP Q9JLS3
B	-16	TYR	-	CLONING ARTIFACT	UNP Q9JLS3
B	-15	ASP	-	CLONING ARTIFACT	UNP Q9JLS3
B	-14	ILE	-	CLONING ARTIFACT	UNP Q9JLS3
B	-13	PRO	-	CLONING ARTIFACT	UNP Q9JLS3
B	-12	THR	-	CLONING ARTIFACT	UNP Q9JLS3
B	-11	THR	-	CLONING ARTIFACT	UNP Q9JLS3
B	-10	GLU	-	CLONING ARTIFACT	UNP Q9JLS3
B	-9	ASN	-	CLONING ARTIFACT	UNP Q9JLS3
B	-8	LEU	-	CLONING ARTIFACT	UNP Q9JLS3
B	-7	TYR	-	CLONING ARTIFACT	UNP Q9JLS3
B	-6	PHE	-	CLONING ARTIFACT	UNP Q9JLS3
B	-5	GLN	-	CLONING ARTIFACT	UNP Q9JLS3
B	-4	GLY	-	CLONING ARTIFACT	UNP Q9JLS3
B	-3	ALA	-	CLONING ARTIFACT	UNP Q9JLS3
B	-2	MET	-	CLONING ARTIFACT	UNP Q9JLS3
B	-1	ASP	-	CLONING ARTIFACT	UNP Q9JLS3
B	0	PRO	-	CLONING ARTIFACT	UNP Q9JLS3
B	181	SEP	SER	MODIFIED RESIDUE	UNP Q9JLS3

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Mg 2 2	0	0
2	A	2	Total Mg 2 2	0	0

- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 3 | B | 1 | Total Ca
1 1 | 0 | 0 |
| 3 | A | 1 | Total Ca
1 1 | 0 | 0 |

-
- The diagram illustrates the chemical structure of Adenosine Triphosphate (ATP). It consists of an adenine base (a purine ring system with an amino group at C6) linked to a ribose sugar (a five-membered ring with hydroxyl groups at C2' and C3'). The ribose is further linked to a chain of three phosphate groups (labeled alpha, beta, and gamma). The phosphate groups are connected by phosphoanhydride bonds, and each phosphate group has a terminal phosphate group (labeled alpha, beta, and gamma) and a terminal phosphate group (labeled alpha, beta, and gamma). The structure is shown in a 3D representation with wedge and dash bonds indicating stereochemistry.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	B	1	Total 31	C 10	N 5	O 13	P 3	0	0


- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|--------------------|---------|---------|
| 5 | A | 209 | Total O
209 209 | 0 | 0 |
| 5 | B | 174 | Total O
174 174 | 0 | 0 |

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: serine/threonine protein kinase TAO2



- Molecule 1: serine/threonine protein kinase TAO2



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	186.47Å 186.47Å 91.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.10	Depositor
% Data completeness (in resolution range)	96.9 (50.00-2.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.211 , 0.244	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5431	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CA, ATP, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/2538	0.70	0/3428
1	B	0.46	0/2538	0.68	0/3428
All	All	0.48	0/5076	0.69	0/6856

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2490	0	2468	75	0
1	B	2490	0	2467	85	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	31	0	10	0	0
4	B	31	0	10	0	0
5	A	209	0	0	8	0
5	B	174	0	0	9	0
All	All	5431	0	4955	159	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 159 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:181:SEP:C	1:B:182:PHE:N	2.30	0.94
1:A:126:VAL:HG23	5:A:637:HOH:O	1.68	0.93
1:B:64:LYS:HA	1:B:64:LYS:NZ	1.85	0.92
1:A:256:ASN:HB2	5:A:668:HOH:O	1.77	0.84
1:A:117:GLU:HG2	5:A:704:HOH:O	1.76	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/348 (88%)	291 (95%)	12 (4%)	3 (1%)	19	13
1	B	305/348 (88%)	286 (94%)	14 (5%)	5 (2%)	12	6
All	All	611/696 (88%)	577 (94%)	26 (4%)	8 (1%)	15	9

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	306	ASN
1	B	163	GLY
1	B	98	GLU
1	B	306	ASN
1	B	62	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/300 (89%)	255 (95%)	13 (5%)	31	28
1	B	268/300 (89%)	255 (95%)	13 (5%)	31	28
All	All	536/600 (89%)	510 (95%)	26 (5%)	31	28

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	267	GLN
1	B	31	LEU
1	B	201	GLN
1	A	306	ASN
1	B	12	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	308	GLN
1	B	85	HIS
1	B	240	ASN
1	A	306	ASN
1	B	246	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	A	181	1,3	8,9,10	1.09	0	8,12,14	2.24	5 (62%)
1	SEP	B	181	1,3	8,9,10	1.09	0	8,12,14	3.84	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	181	1,3	-	0/6/8/10	0/0/0/0
1	SEP	B	181	1,3	-	0/6/8/10	0/0/0/0

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	181	SEP	OG-CB-CA	-8.85	100.73	108.27
1	B	181	SEP	O2P-P-OG	-4.39	93.92	106.56
1	A	181	SEP	O2P-P-OG	-3.10	97.64	106.56
1	A	181	SEP	OG-P-O1P	-3.05	99.39	107.14
1	A	181	SEP	O3P-P-O1P	2.10	117.33	110.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	181	SEP	2	0
1	B	181	SEP	4	0

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ATP	A	411	2	24,33,33	2.35	10 (41%)	31,52,52	3.73	13 (41%)
4	ATP	B	412	2	24,33,33	2.29	10 (41%)	31,52,52	3.63	13 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	411	2	-	0/18/38/38	0/3/3/3
4	ATP	B	412	2	-	0/18/38/38	0/3/3/3

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	412	ATP	O5'-C5'	-5.15	1.23	1.44
4	A	411	ATP	O5'-C5'	-4.84	1.25	1.44
4	B	412	ATP	PA-O5'	-3.58	1.42	1.59
4	A	411	ATP	PA-O5'	-3.40	1.43	1.59
4	A	411	ATP	C8-N7	-2.93	1.29	1.34

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	411	ATP	C5'-C4'-C3'	-7.36	85.99	115.21
4	B	412	ATP	O5'-PA-O1A	-7.30	81.30	109.62
4	B	412	ATP	C5'-C4'-C3'	-7.18	86.70	115.21
4	A	411	ATP	O5'-PA-O1A	-6.83	83.12	109.62
4	A	411	ATP	N3-C2-N1	-2.83	126.73	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.