



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:35 PM GMT

PDB ID : 3U52  
Title : X-ray Crystal Structure of Xenon-Pressurized Phenol Hydroxylase from Pseudomonas sp. OX1  
Authors : McCormick, M.S.; Lippard, S.J.  
Deposited on : 2011-10-10  
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	511	<div> <div>6%</div> <div>82%</div> <div>14%</div> <div>• •</div> </div>
1	B	511	<div> <div>9%</div> <div>79%</div> <div>15%</div> <div>• •</div> </div>
2	C	333	<div> <div>%</div> <div>77%</div> <div>14%</div> <div>• • 5%</div> </div>
2	D	333	<div> <div>8%</div> <div>81%</div> <div>14%</div> <div>• •</div> </div>
3	E	119	<div> <div>8%</div> <div>75%</div> <div>20%</div> <div>• •</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	119	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	EPE	B	517	-	-	X	X
7	GOL	A	516	-	-	-	X
7	GOL	A	519	-	-	-	X
7	GOL	B	518	-	-	-	X
7	GOL	D	334	-	-	X	X
8	XE	A	520	-	-	-	X
8	XE	A	521	-	-	X	-
8	XE	A	522	-	-	X	-
8	XE	B	520	-	-	-	X
8	XE	B	521	-	-	-	X
9	MPO	B	516	-	-	X	-

## 2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 16039 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenol hydroxylase component pHN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	494	Total	C	N	O	S	0	0	0
			4123	2652	693	754	24			
1	B	493	Total	C	N	O	S	0	0	0
			4110	2643	690	753	24			

- Molecule 2 is a protein called Phenol hydroxylase component pHL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	318	Total	C	N	O	S	0	0	0
			2589	1630	449	492	18			
2	D	328	Total	C	N	O	S	0	0	0
			2659	1673	461	507	18			

- Molecule 3 is a protein called Phenol hydroxylase component pHO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	118	Total	C	N	O	S	0	0	0
			925	602	145	173	5			
3	F	118	Total	C	N	O	S	0	0	0
			925	602	145	173	5			

- Molecule 4 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Fe	0	0
			2	2		
4	A	2	Total	Fe	0	0
			2	2		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Zn	0	0
			1	1		
5	A	1	Total	Zn	0	0
			1	1		

- Molecule 6 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Cu	0	0
			1	1		
6	A	1	Total	Cu	0	0
			1	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		
7	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is XENON (three-letter code: XE) (formula: Xe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	11	Total	Xe	0	0
			11	11		
8	A	11	Total	Xe	0	0
			11	11		
8	D	1	Total	Xe	0	0
			1	1		
8	C	1	Total	Xe	0	0
			1	1		

- Molecule 9 is 3[N-MORPHOLINO]PROPANE SULFONIC ACID (three-letter code: MPO) (formula: C<sub>7</sub>H<sub>15</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	B	1	Total	C	N	O	S	0	0
			13	7	1	4	1		

- Molecule 10 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C<sub>8</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	160	Total	O	0	0
			160	160		
11	B	139	Total	O	0	0
			139	139		
11	C	118	Total	O	0	0
			118	118		
11	D	99	Total	O	0	0
			99	99		
11	E	40	Total	O	0	0
			40	40		
11	F	32	Total	O	0	0
			32	32		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phenol hydroxylase component phN




- Molecule 1: Phenol hydroxylase component phN




- Molecule 2: Phenol hydroxylase component phL









- Molecule 2: Phenol hydroxylase component phL



- Molecule 3: Phenol hydroxylase component phO



- Molecule 3: Phenol hydroxylase component phO



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.94Å 141.76Å 181.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.00 – 1.95 39.87 – 1.95	Depositor EDS
% Data completeness (in resolution range)	92.5 (38.00-1.95) 92.5 (39.87-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.57 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.183 , 0.228 0.188 , 0.184	Depositor DCC
$R_{free}$ test set	7399 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.9	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 55.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 145806 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	16039	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, MPO, XE, FE, EPE, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.89	0/4257	0.85	8/5766 (0.1%)
1	B	0.85	3/4244 (0.1%)	0.79	2/5751 (0.0%)
2	C	0.93	6/2648 (0.2%)	1.12	20/3580 (0.6%)
2	D	0.84	0/2718	0.95	13/3676 (0.4%)
3	E	0.83	0/953	0.87	2/1297 (0.2%)
3	F	0.79	0/953	0.76	0/1297
All	All	0.87	9/15773 (0.1%)	0.90	45/21367 (0.2%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	316	GLU	CB-CG	5.94	1.63	1.52
2	C	238	PHE	CG-CD2	5.80	1.47	1.38
2	C	199	ARG	CD-NE	-5.48	1.37	1.46
1	B	445	GLU	CG-CD	5.48	1.60	1.51
2	C	238	PHE	CE1-CZ	5.41	1.47	1.37

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	199	ARG	NE-CZ-NH2	-22.56	109.02	120.30
2	C	22	ARG	NE-CZ-NH2	-15.90	112.35	120.30
2	D	199	ARG	NE-CZ-NH2	-15.68	112.46	120.30
2	C	199	ARG	NE-CZ-NH1	14.56	127.58	120.30
2	D	22	ARG	NE-CZ-NH2	-11.65	114.48	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4123	0	3866	65	0
1	B	4110	0	3842	82	0
2	C	2589	0	2501	54	0
2	D	2659	0	2571	44	0
3	E	925	0	898	22	0
3	F	925	0	898	13	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	24	0	32	5	0
7	B	12	0	16	1	0
7	C	12	0	16	0	0
7	D	6	0	8	4	0
7	F	6	0	8	0	0
8	A	11	0	0	6	0
8	B	11	0	0	3	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
9	B	13	0	11	9	0
10	B	15	0	17	7	0
11	A	160	0	0	2	0
11	B	139	0	0	6	0
11	C	118	0	0	3	0
11	D	99	0	0	3	0
11	E	40	0	0	1	0
11	F	32	0	0	0	0
All	All	16039	0	14684	248	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 248 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:73:PHE:O	3:E:74:LEU:HB2	1.43	1.17
1:A:198:PHE:CE1	8:A:521:XE:XE	2.83	1.09
1:B:118:VAL:HG23	9:B:516:MPO:H61	1.19	1.08
1:A:337:THR:HG21	1:A:376:ARG:HH11	1.14	1.08
2:C:189:TRP:CD1	2:C:190:MET:CE	2.36	1.08

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	492/511 (96%)	477 (97%)	15 (3%)	0	100	100
1	B	491/511 (96%)	473 (96%)	17 (4%)	1 (0%)	52	43
2	C	316/333 (95%)	313 (99%)	3 (1%)	0	100	100
2	D	326/333 (98%)	319 (98%)	7 (2%)	0	100	100
3	E	116/119 (98%)	113 (97%)	2 (2%)	1 (1%)	21	9
3	F	116/119 (98%)	111 (96%)	5 (4%)	0	100	100
All	All	1857/1926 (96%)	1806 (97%)	49 (3%)	2 (0%)	56	48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	E	74	LEU
1	B	467	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	434/447 (97%)	412 (95%)	22 (5%)	29	13
1	B	432/447 (97%)	414 (96%)	18 (4%)	36	21
2	C	272/285 (95%)	254 (93%)	18 (7%)	21	7
2	D	280/285 (98%)	263 (94%)	17 (6%)	23	9
3	E	98/99 (99%)	88 (90%)	10 (10%)	9	2
3	F	98/99 (99%)	89 (91%)	9 (9%)	11	3
All	All	1614/1662 (97%)	1520 (94%)	94 (6%)	25	10

5 of 94 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	62	ARG
2	C	253	MET
3	F	59	LYS
2	C	68	GLU
2	C	156	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	153	ASN
1	B	344	GLN
2	D	165	GLN
1	B	160	HIS
1	B	330	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 44 ligands modelled in this entry, 32 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	GOL	A	516	4	5,5,5	0.36	0	5,5,5	1.20	0
7	GOL	A	517	-	5,5,5	0.40	0	5,5,5	1.39	1 (20%)
7	GOL	A	518	-	5,5,5	0.30	0	5,5,5	0.39	0
7	GOL	A	519	-	5,5,5	0.58	0	5,5,5	1.08	0
9	MPO	B	516	-	12,13,13	4.39	3 (25%)	15,17,17	6.88	8 (53%)
10	EPE	B	517	-	14,15,15	0.47	0	18,20,20	2.21	6 (33%)
7	GOL	B	518	-	5,5,5	0.35	0	5,5,5	0.37	0
7	GOL	B	519	4	5,5,5	0.27	0	5,5,5	0.82	0
7	GOL	C	334	-	5,5,5	0.39	0	5,5,5	0.56	0
7	GOL	C	335	-	5,5,5	0.48	0	5,5,5	0.77	0
7	GOL	D	334	-	5,5,5	0.58	0	5,5,5	1.21	0
7	GOL	F	120	-	5,5,5	0.27	0	5,5,5	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	516	4	-	0/4/4/4	0/0/0/0
7	GOL	A	517	-	-	0/4/4/4	0/0/0/0
7	GOL	A	518	-	-	0/4/4/4	0/0/0/0
7	GOL	A	519	-	-	0/4/4/4	0/0/0/0
9	MPO	B	516	-	-	0/7/15/15	0/1/1/1
10	EPE	B	517	-	-	0/9/19/19	0/1/1/1
7	GOL	B	518	-	-	0/4/4/4	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	B	519	4	-	0/4/4/4	0/0/0/0
7	GOL	C	334	-	-	0/4/4/4	0/0/0/0
7	GOL	C	335	-	-	0/4/4/4	0/0/0/0
7	GOL	D	334	-	-	0/4/4/4	0/0/0/0
7	GOL	F	120	-	-	0/4/4/4	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	B	516	MPO	C3-N1	-11.34	1.20	1.47
9	B	516	MPO	C7-N1	-7.64	1.26	1.46
9	B	516	MPO	C2-C3	-6.06	1.25	1.51

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	517	GOL	C3-C2-C1	-2.72	100.44	111.12
9	B	516	MPO	C5-C4-N1	-2.17	106.84	110.12
10	B	517	EPE	O8-C8-C7	-2.10	101.63	111.28
10	B	517	EPE	C3-C2-N1	-2.07	106.93	110.63
10	B	517	EPE	C7-N4-C5	2.33	117.24	111.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	517	GOL	2	0
7	A	519	GOL	3	0
9	B	516	MPO	9	0
10	B	517	EPE	7	0
7	B	519	GOL	1	0
7	D	334	GOL	4	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	494/511 (96%)	0.24	31 (6%) 23 33	15, 24, 37, 51	0
1	B	493/511 (96%)	0.44	47 (9%) 10 17	18, 28, 41, 54	0
2	C	318/333 (95%)	0.09	2 (0%) 90 94	16, 24, 35, 58	0
2	D	328/333 (98%)	0.41	25 (7%) 17 26	18, 28, 46, 71	0
3	E	118/119 (99%)	0.33	9 (7%) 17 26	19, 32, 42, 46	0
3	F	118/119 (99%)	0.49	9 (7%) 17 26	22, 35, 47, 53	0
All	All	1869/1926 (97%)	0.32	123 (6%) 22 31	15, 27, 42, 71	0

The worst 5 of 123 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	331	VAL	7.3
2	D	10	VAL	7.2
1	B	386	ALA	6.1
2	D	9	SER	5.7
2	D	8	ASN	5.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
7	GOL	D	334	6/6	0.93	0.30	7.28	22,35,41,44	0
10	EPE	B	517	15/15	0.62	0.28	4.52	47,68,82,82	0
7	GOL	A	519	6/6	0.93	0.21	3.60	30,36,37,38	0
7	GOL	A	516	6/6	0.89	0.31	3.36	37,40,43,44	0
8	XE	A	520	1/1	0.88	0.33	2.84	49,49,49,49	1
8	XE	B	520	1/1	0.94	0.30	2.80	56,56,56,56	1
7	GOL	B	518	6/6	0.63	0.26	2.45	59,65,68,70	0
8	XE	B	521	1/1	0.95	0.33	2.37	53,53,53,53	1
7	GOL	A	517	6/6	0.90	0.20	1.88	32,40,42,42	0
7	GOL	B	519	6/6	0.93	0.25	1.61	43,45,46,50	0
9	MPO	B	516	13/13	0.84	0.22	1.38	50,54,67,69	0
7	GOL	A	518	6/6	0.94	0.11	0.62	30,33,34,36	0
8	XE	A	530	1/1	0.88	0.12	0.59	53,53,53,53	1
8	XE	B	530	1/1	0.92	0.12	0.11	50,50,50,50	1
4	FE	A	513	1/1	0.96	0.14	-0.01	34,34,34,34	0
7	GOL	C	335	6/6	0.96	0.11	-0.02	25,32,36,39	0
4	FE	B	513	1/1	0.97	0.13	-0.31	36,36,36,36	0
4	FE	B	512	1/1	1.00	0.13	-0.36	31,31,31,31	0
4	FE	A	512	1/1	0.99	0.12	-0.82	33,33,33,33	0
7	GOL	F	120	6/6	0.96	0.07	-1.01	31,35,36,38	0
5	ZN	A	514	1/1	1.00	0.09	-1.10	19,19,19,19	0
5	ZN	B	514	1/1	0.99	0.09	-1.17	23,23,23,23	0
6	CU	B	515	1/1	0.98	0.04	-1.26	33,33,33,33	0
8	XE	B	524	1/1	0.96	0.05	-1.52	45,45,45,45	0
8	XE	D	335	1/1	0.99	0.03	-1.57	49,49,49,49	0
8	XE	B	527	1/1	0.97	0.06	-1.71	46,46,46,46	1
8	XE	B	525	1/1	0.99	0.03	-1.74	44,44,44,44	0
8	XE	B	526	1/1	0.99	0.02	-1.75	49,49,49,49	0
8	XE	A	526	1/1	0.98	0.03	-1.78	48,48,48,48	0
8	XE	A	525	1/1	1.00	0.03	-1.78	40,40,40,40	0
8	XE	B	529	1/1	0.98	0.05	-1.80	43,43,43,43	0
6	CU	A	515	1/1	0.99	0.04	-1.87	32,32,32,32	0
8	XE	A	527	1/1	0.99	0.03	-1.89	44,44,44,44	1
8	XE	B	522	1/1	0.96	0.15	-2.00	46,46,46,46	1
8	XE	A	524	1/1	0.95	0.05	-2.05	42,42,42,42	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
8	XE	A	522	1/1	0.95	0.10	-2.07	42,42,42,42	1
8	XE	A	521	1/1	0.98	0.10	-2.27	47,47,47,47	1
8	XE	A	528	1/1	0.99	0.04	-2.42	36,36,36,36	0
8	XE	A	529	1/1	0.99	0.03	-2.54	41,41,41,41	0
8	XE	B	528	1/1	1.00	0.03	-3.39	40,40,40,40	0
8	XE	C	336	1/1	0.99	0.02	-5.34	41,41,41,41	0
8	XE	B	523	1/1	0.99	0.04	-	37,37,37,37	0
8	XE	A	523	1/1	0.99	0.05	-	32,32,32,32	0
7	GOL	C	334	6/6	0.74	0.19	-	48,55,56,58	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.