



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:35 PM GMT

PDB ID : 3U5R  
Title : Crystal structure of a hypothetical protein SMc02350 from *Sinorhizobium meliloti* 1021  
Authors : Agarwal, R.; Almo, S.C.; Swaminathan, S.; New York Structural Genomics Research Consortium (NYSGRC)  
Deposited on : 2011-10-11  
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry. We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org). A user guide is available at <http://wwpdb.org/validation/2016/XrayValidationReportHelp> with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1192 (2.04-2.04)
Clashscore	102246	1269 (2.04-2.04)
Ramachandran outliers	100387	1258 (2.04-2.04)
Sidechain outliers	100360	1258 (2.04-2.04)
RSRZ outliers	91569	1194 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	218	 72% 11% • 16%
1	F	218	 77% 9% • 13%
1	G	218	 74% 11% • 13%
1	H	218	 72% 10% • 16%

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 5981 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	E	183	1374	872	236	263	3	0	0	0
1	F	189	1419	900	243	273	3	0	0	0
1	G	189	1420	901	243	273	3	0	0	0
1	H	184	1373	871	234	265	3	0	0	0

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	1	MET	-	EXPRESSION TAG	UNP Q92MN6
E	2	HIS	-	EXPRESSION TAG	UNP Q92MN6
E	3	HIS	-	EXPRESSION TAG	UNP Q92MN6
E	4	HIS	-	EXPRESSION TAG	UNP Q92MN6
E	5	HIS	-	EXPRESSION TAG	UNP Q92MN6
E	6	HIS	-	EXPRESSION TAG	UNP Q92MN6
E	7	HIS	-	EXPRESSION TAG	UNP Q92MN6
E	8	SER	-	EXPRESSION TAG	UNP Q92MN6
E	9	SER	-	EXPRESSION TAG	UNP Q92MN6
E	10	GLY	-	EXPRESSION TAG	UNP Q92MN6
E	11	VAL	-	EXPRESSION TAG	UNP Q92MN6
E	12	ASP	-	EXPRESSION TAG	UNP Q92MN6
E	13	LEU	-	EXPRESSION TAG	UNP Q92MN6
E	14	GLY	-	EXPRESSION TAG	UNP Q92MN6
E	15	THR	-	EXPRESSION TAG	UNP Q92MN6
E	16	GLU	-	EXPRESSION TAG	UNP Q92MN6
E	17	ASN	-	EXPRESSION TAG	UNP Q92MN6
E	18	LEU	-	EXPRESSION TAG	UNP Q92MN6
E	19	TYR	-	EXPRESSION TAG	UNP Q92MN6
E	20	PHE	-	EXPRESSION TAG	UNP Q92MN6
E	21	GLN	-	EXPRESSION TAG	UNP Q92MN6

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	22	SER	-	EXPRESSION TAG	UNP Q92MN6
E	23	MET	-	EXPRESSION TAG	UNP Q92MN6
F	1	MET	-	EXPRESSION TAG	UNP Q92MN6
F	2	HIS	-	EXPRESSION TAG	UNP Q92MN6
F	3	HIS	-	EXPRESSION TAG	UNP Q92MN6
F	4	HIS	-	EXPRESSION TAG	UNP Q92MN6
F	5	HIS	-	EXPRESSION TAG	UNP Q92MN6
F	6	HIS	-	EXPRESSION TAG	UNP Q92MN6
F	7	HIS	-	EXPRESSION TAG	UNP Q92MN6
F	8	SER	-	EXPRESSION TAG	UNP Q92MN6
F	9	SER	-	EXPRESSION TAG	UNP Q92MN6
F	10	GLY	-	EXPRESSION TAG	UNP Q92MN6
F	11	VAL	-	EXPRESSION TAG	UNP Q92MN6
F	12	ASP	-	EXPRESSION TAG	UNP Q92MN6
F	13	LEU	-	EXPRESSION TAG	UNP Q92MN6
F	14	GLY	-	EXPRESSION TAG	UNP Q92MN6
F	15	THR	-	EXPRESSION TAG	UNP Q92MN6
F	16	GLU	-	EXPRESSION TAG	UNP Q92MN6
F	17	ASN	-	EXPRESSION TAG	UNP Q92MN6
F	18	LEU	-	EXPRESSION TAG	UNP Q92MN6
F	19	TYR	-	EXPRESSION TAG	UNP Q92MN6
F	20	PHE	-	EXPRESSION TAG	UNP Q92MN6
F	21	GLN	-	EXPRESSION TAG	UNP Q92MN6
F	22	SER	-	EXPRESSION TAG	UNP Q92MN6
F	23	MET	-	EXPRESSION TAG	UNP Q92MN6
G	1	MET	-	EXPRESSION TAG	UNP Q92MN6
G	2	HIS	-	EXPRESSION TAG	UNP Q92MN6
G	3	HIS	-	EXPRESSION TAG	UNP Q92MN6
G	4	HIS	-	EXPRESSION TAG	UNP Q92MN6
G	5	HIS	-	EXPRESSION TAG	UNP Q92MN6
G	6	HIS	-	EXPRESSION TAG	UNP Q92MN6
G	7	HIS	-	EXPRESSION TAG	UNP Q92MN6
G	8	SER	-	EXPRESSION TAG	UNP Q92MN6
G	9	SER	-	EXPRESSION TAG	UNP Q92MN6
G	10	GLY	-	EXPRESSION TAG	UNP Q92MN6
G	11	VAL	-	EXPRESSION TAG	UNP Q92MN6
G	12	ASP	-	EXPRESSION TAG	UNP Q92MN6
G	13	LEU	-	EXPRESSION TAG	UNP Q92MN6
G	14	GLY	-	EXPRESSION TAG	UNP Q92MN6
G	15	THR	-	EXPRESSION TAG	UNP Q92MN6
G	16	GLU	-	EXPRESSION TAG	UNP Q92MN6
G	17	ASN	-	EXPRESSION TAG	UNP Q92MN6

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
G	18	LEU	-	EXPRESSION TAG	UNP Q92MN6
G	19	TYR	-	EXPRESSION TAG	UNP Q92MN6
G	20	PHE	-	EXPRESSION TAG	UNP Q92MN6
G	21	GLN	-	EXPRESSION TAG	UNP Q92MN6
G	22	SER	-	EXPRESSION TAG	UNP Q92MN6
G	23	MET	-	EXPRESSION TAG	UNP Q92MN6
H	1	MET	-	EXPRESSION TAG	UNP Q92MN6
H	2	HIS	-	EXPRESSION TAG	UNP Q92MN6
H	3	HIS	-	EXPRESSION TAG	UNP Q92MN6
H	4	HIS	-	EXPRESSION TAG	UNP Q92MN6
H	5	HIS	-	EXPRESSION TAG	UNP Q92MN6
H	6	HIS	-	EXPRESSION TAG	UNP Q92MN6
H	7	HIS	-	EXPRESSION TAG	UNP Q92MN6
H	8	SER	-	EXPRESSION TAG	UNP Q92MN6
H	9	SER	-	EXPRESSION TAG	UNP Q92MN6
H	10	GLY	-	EXPRESSION TAG	UNP Q92MN6
H	11	VAL	-	EXPRESSION TAG	UNP Q92MN6
H	12	ASP	-	EXPRESSION TAG	UNP Q92MN6
H	13	LEU	-	EXPRESSION TAG	UNP Q92MN6
H	14	GLY	-	EXPRESSION TAG	UNP Q92MN6
H	15	THR	-	EXPRESSION TAG	UNP Q92MN6
H	16	GLU	-	EXPRESSION TAG	UNP Q92MN6
H	17	ASN	-	EXPRESSION TAG	UNP Q92MN6
H	18	LEU	-	EXPRESSION TAG	UNP Q92MN6
H	19	TYR	-	EXPRESSION TAG	UNP Q92MN6
H	20	PHE	-	EXPRESSION TAG	UNP Q92MN6
H	21	GLN	-	EXPRESSION TAG	UNP Q92MN6
H	22	SER	-	EXPRESSION TAG	UNP Q92MN6
H	23	MET	-	EXPRESSION TAG	UNP Q92MN6


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	E	114	Total O 114 114	0	0
2	F	75	Total O 75 75	0	0
2	G	62	Total O 62 62	0	0
2	H	144	Total O 144 144	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: uncharacterized protein




- Molecule 1: uncharacterized protein



- Molecule 1: uncharacterized protein



- Molecule 1: uncharacterized protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.66Å 58.57Å 85.48Å 82.65° 83.58° 66.29°	Depositor
Resolution (Å)	34.02 – 2.05 34.02 – 2.05	Depositor EDS
% Data completeness (in resolution range)	97.8 (34.02-2.05) 94.6 (34.02-2.05)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.17 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.182 , 0.232 0.183 , 0.231	Depositor DCC
$R_{free}$ test set	3026 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.4	Xtriage
Anisotropy	0.033	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 59763 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5981	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	E	1.23	0/1402	0.97	1/1902 (0.1%)
1	F	1.05	2/1450 (0.1%)	0.87	3/1971 (0.2%)
1	G	1.02	0/1451	0.93	2/1973 (0.1%)
1	H	1.30	6/1401 (0.4%)	1.03	5/1904 (0.3%)
All	All	1.15	8/5704 (0.1%)	0.95	11/7750 (0.1%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	182	VAL	CB-CG2	5.46	1.64	1.52
1	F	152	ARG	CG-CD	5.41	1.65	1.51
1	H	166	ARG	CZ-NH2	5.34	1.40	1.33
1	H	135	VAL	CB-CG2	-5.12	1.42	1.52
1	H	118	VAL	CB-CG2	5.11	1.63	1.52

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	130	ASP	CB-CG-OD1	5.91	123.62	118.30
1	F	45	ASP	CB-CG-OD1	5.68	123.42	118.30
1	H	166	ARG	NE-CZ-NH2	5.49	123.04	120.30
1	H	78	ARG	NE-CZ-NH2	-5.39	117.61	120.30
1	H	130	ASP	CB-CG-OD2	-5.36	113.48	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1374	0	1345	18	0
1	F	1419	0	1373	12	0
1	G	1420	0	1374	17	0
1	H	1373	0	1333	18	0
2	E	114	0	0	2	0
2	F	75	0	0	0	0
2	G	62	0	0	1	0
2	H	144	0	0	4	0
All	All	5981	0	5425	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 65 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:161:GLN:HE21	1:H:166:ARG:HH22	1.09	0.99
1:F:174:THR:HG22	1:F:176:ALA:H	1.35	0.92
1:F:174:THR:CG2	1:F:176:ALA:HB2	2.12	0.79
1:F:174:THR:HG21	1:F:176:ALA:HB2	1.63	0.79
1:H:161:GLN:NE2	1:H:166:ARG:HH22	1.80	0.78

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	E	181/218 (83%)	175 (97%)	6 (3%)	0	100 100
1	F	187/218 (86%)	179 (96%)	8 (4%)	0	100 100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	187/218 (86%)	180 (96%)	7 (4%)	0	100	100
1	H	182/218 (84%)	175 (96%)	7 (4%)	0	100	100
All	All	737/872 (84%)	709 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	138/169 (82%)	134 (97%)	4 (3%)	50	42
1	F	142/169 (84%)	137 (96%)	5 (4%)	43	35
1	G	142/169 (84%)	136 (96%)	6 (4%)	36	28
1	H	137/169 (81%)	130 (95%)	7 (5%)	29	19
All	All	559/676 (83%)	537 (96%)	22 (4%)	39	30

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	31	SER
1	G	79	GLU
1	H	177	ASP
1	G	57	LYS
1	G	67	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	159	HIS
1	G	161	GLN
1	H	159	HIS
1	F	161	GLN
1	H	101	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	E	183/218 (83%)	-0.45	1 (0%) 91 93	11, 19, 36, 48	0
1	F	189/218 (86%)	-0.23	5 (2%) 59 65	17, 29, 47, 59	0
1	G	189/218 (86%)	-0.25	3 (1%) 74 79	17, 32, 49, 56	0
1	H	184/218 (84%)	-0.37	1 (0%) 91 93	11, 18, 34, 47	0
All	All	745/872 (85%)	-0.32	10 (1%) 79 83	11, 23, 45, 59	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	168	GLY	4.1
1	G	168	GLY	3.4
1	H	207	ALA	2.9
1	F	170	GLY	2.3
1	E	70	ARG	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.