



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:51 PM GMT

PDB ID : 4U5N  
Title : IMPORTIN-ALPHA MINOR NLS SITE INHIBITOR  
Authors : Stewart, M.; Valkov, E.; Holvey, R.S.  
Deposited on : 2014-07-25  
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4425 (2.34-2.30)
Clashscore	102246	5057 (2.34-2.30)
Ramachandran outliers	100387	5008 (2.34-2.30)
Sidechain outliers	100360	5007 (2.34-2.30)
RSRZ outliers	91569	4432 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	426	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	3D4	A	501	-	-	-	X




In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Importin subunit alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	426	Total	C	N	O	S	0	2	0
			3254	2072	551	621	10			

- Molecule 2 is N 2-[4-(pyridin-3-yl)benzyl]-L-lysyl-N-[(1R,2S,3R)-1-{(2R)-1-amino-1-oxo-3-phenylpropan-2-yl}amino]-1,3-dihydroxybutan-2-yl]glycinamide (three-letter code: 3D4) (formula: C<sub>33</sub>H<sub>45</sub>N<sub>7</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			45	33	7	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	113	Total O 113 113	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Importin subunit alpha-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.50Å 91.08Å 98.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.54 – 2.31 19.82 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.54-2.31) 97.3 (19.82-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.61 (at 2.50Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1702)	Depositor
R, $R_{free}$	0.187 , 0.221 0.303 , 0.287	Depositor DCC
$R_{free}$ test set	1256 reflections (5.39%)	DCC
Wilson B-factor (Å <sup>2</sup> )	38.8	Xtriage
Anisotropy	0.461	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 30.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 24567 reflections	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	3412	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 3D4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.43	0/3318	0.57	0/4522

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3254	0	3330	22	0
2	A	45	0	45	1	0
3	A	113	0	0	0	0
All	All	3412	0	3375	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108:LYS:HG3	1:A:109:GLN:HG3	1.79	0.63

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:THR:HG21	1:A:184:TRP:HD1	1.68	0.57
1:A:284:GLU:H	1:A:284:GLU:CD	2.09	0.56
1:A:207:ASP:OD1	1:A:251:GLN:NE2	2.39	0.55
1:A:429:LEU:HD22	1:A:476:LEU:HD11	1.87	0.55
1:A:145:THR:HG21	1:A:184:TRP:CD1	2.43	0.53
1:A:470:LEU:O	1:A:474:GLU:HG3	2.11	0.51
1:A:104:LEU:HD21	1:A:115:ILE:HG13	1.94	0.50
2:A:501:3D4:OAG	2:A:501:3D4:H31	2.12	0.49
1:A:282:PRO:HB2	1:A:284:GLU:OE1	2.12	0.49
1:A:386:LEU:HD21	1:A:425:LEU:HD13	1.94	0.49
1:A:360:SER:HA	1:A:400:ALA:HA	1.95	0.49
1:A:215:VAL:O	1:A:258:ARG:NH2	2.44	0.49
1:A:397:ALA:O	1:A:401:ILE:HG12	2.13	0.48
1:A:425:LEU:HG	1:A:440:ILE:HG23	1.98	0.46
1:A:472:LYS:O	1:A:476:LEU:HG	2.16	0.45
1:A:145:THR:CG2	1:A:184:TRP:CD1	3.01	0.43
1:A:272:CYS:HB3	1:A:312:PRO:HB2	2.01	0.42
1:A:238:ARG:O	1:A:239:ASN:HB2	2.20	0.42
1:A:434:THR:O	1:A:438:GLN:HG3	2.19	0.42
1:A:253:LEU:O	1:A:257:VAL:HG23	2.21	0.41
1:A:233:LEU:HA	1:A:233:LEU:HD23	1.85	0.41
1:A:378:LEU:HA	1:A:378:LEU:HD23	1.89	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	426/426 (100%)	414 (97%)	11 (3%)	1 (0%)	52 64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	485	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/357 (101%)	356 (99%)	3 (1%)	86	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	247[A]	ASP
1	A	247[B]	ASP
1	A	485	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	477	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	3D4	A	501	-	45,47,47	2.11	8 (17%)	50,61,61	1.38	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3D4	A	501	-	-	0/46/47/47	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	3D4	CAA-CBP	-5.84	1.37	1.51
2	A	501	3D4	OAD-CBH	-2.42	1.19	1.23
2	A	501	3D4	O-C	-2.17	1.19	1.23
2	A	501	3D4	OAE-CBI	-2.03	1.19	1.23
2	A	501	3D4	OAH-CBP	4.57	1.53	1.43
2	A	501	3D4	CBI-NBF	5.96	1.46	1.34
2	A	501	3D4	C-NBD	6.12	1.46	1.33
2	A	501	3D4	CBH-NAC	6.83	1.46	1.32

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	3D4	OAE-CBI-NBF	-3.86	116.47	123.01
2	A	501	3D4	CAA-CBP-CBS	-2.33	107.81	112.93
2	A	501	3D4	OAD-CBH-NAC	-2.27	119.72	123.08
2	A	501	3D4	CAQ-CAS-CBN	-2.01	118.21	121.14
2	A	501	3D4	CAT-CBN-CAS	2.06	121.45	117.55
2	A	501	3D4	CAM-NBC-CAU	2.42	121.29	116.84

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	3D4	CBR-CBH-NAC	2.69	120.94	116.60
2	A	501	3D4	CAZ-CBI-NBF	3.38	124.63	116.12
2	A	501	3D4	CBS-NBF-CBI	3.58	128.63	123.05

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	3D4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	426/426 (100%)	1.56	109 (25%) ⓘ ⓘ	33, 46, 87, 107	0

All (109) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	74	VAL	7.9
1	A	485	TYR	7.1
1	A	73	THR	6.4
1	A	143	ALA	4.9
1	A	483	SER	4.7
1	A	72	GLY	4.7
1	A	432	LYS	4.6
1	A	107	GLU	4.6
1	A	109	GLN	4.4
1	A	497	SER	4.2
1	A	484	VAL	4.2
1	A	478	ARG	4.1
1	A	184	TRP	4.0
1	A	390	ASP	3.9
1	A	263	ASN	3.9
1	A	100	ALA	3.8
1	A	75	ASN	3.7
1	A	220	THR	3.7
1	A	480	GLU	3.7
1	A	108	LYS	3.5
1	A	225	TYR	3.5
1	A	358	THR	3.4
1	A	477	GLN	3.2
1	A	103	LEU	3.2
1	A	476	LEU	3.2
1	A	491	LEU	3.2
1	A	373	VAL	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	147	ILE	3.1
1	A	331	VAL	3.1
1	A	411	GLU	3.1
1	A	409	THR	3.1
1	A	481	ASN	3.0
1	A	392	LYS	3.0
1	A	466	GLU	2.9
1	A	493	GLU	2.9
1	A	315	ARG	2.9
1	A	142	TRP	2.9
1	A	144	LEU	2.8
1	A	436	ILE	2.8
1	A	335	GLY	2.8
1	A	88	ASN	2.8
1	A	342	SER	2.8
1	A	326	GLU	2.8
1	A	148	ALA	2.7
1	A	368	ASP	2.7
1	A	435	LYS	2.7
1	A	442	ASP	2.7
1	A	414	VAL	2.7
1	A	132	ASP	2.7
1	A	482	GLU	2.7
1	A	465	GLU	2.7
1	A	151	THR	2.7
1	A	241	ASN	2.6
1	A	309	ILE	2.6
1	A	359	MET	2.6
1	A	474	GLU	2.6
1	A	117	ARG	2.6
1	A	272	CYS	2.6
1	A	334	ALA	2.5
1	A	301	LEU	2.5
1	A	362	ILE	2.5
1	A	357	TRP	2.5
1	A	105	SER	2.5
1	A	236	LEU	2.5
1	A	99	ALA	2.5
1	A	277	TYR	2.5
1	A	494	LYS	2.5
1	A	80	ASP	2.5
1	A	488	SER	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	456	GLU	2.4
1	A	256	LEU	2.4
1	A	490	ASN	2.4
1	A	96	ALA	2.4
1	A	259	LEU	2.4
1	A	255	THR	2.4
1	A	274	ALA	2.4
1	A	110	PRO	2.3
1	A	458	GLU	2.3
1	A	448	PHE	2.3
1	A	106	ARG	2.3
1	A	308	PRO	2.3
1	A	356	THR	2.3
1	A	312	PRO	2.2
1	A	412	GLN	2.2
1	A	339	VAL	2.2
1	A	218	LEU	2.2
1	A	223	CYS	2.2
1	A	305	THR	2.2
1	A	141	ALA	2.2
1	A	332	ILE	2.2
1	A	340	PHE	2.2
1	A	149	SER	2.2
1	A	115	ILE	2.2
1	A	296	PRO	2.2
1	A	179	SER	2.1
1	A	190	ALA	2.1
1	A	237	CYS	2.1
1	A	76	TRP	2.1
1	A	102	LYS	2.1
1	A	97	THR	2.1
1	A	288	MET	2.1
1	A	182	ALA	2.1
1	A	214	ALA	2.1
1	A	180	GLU	2.1
1	A	444	ILE	2.1
1	A	138	PHE	2.0
1	A	240	LYS	2.0
1	A	471	ASP	2.0
1	A	441	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	3D4	A	501	45/45	0.52	0.62	10.67	46,64,95,104	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.