



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:36 PM GMT

PDB ID : 3U6U
Title : Crystal structure of the putative acetylglutamate kinase from thermus thermophilus
Authors : Karthe, P.; Kumarevel, T.S.; Kuramitsu, S.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2011-10-13
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5832 (1.94-1.90)
Clashscore	102246	6540 (1.94-1.90)
Ramachandran outliers	100387	6464 (1.94-1.90)
Sidechain outliers	100360	6465 (1.94-1.90)
RSRZ outliers	91569	5846 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	
1	C	269	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	271	-	-	-	X

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4126 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative acetylglutamate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	0	0
			1864	1185	327	347	5			
1	C	266	Total	C	N	O	S	0	0	0
			1992	1264	357	366	5			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	105	Total 105	O 105	0	0
3	C	145	Total 145	O 145	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Putative acetylglutamate kinase



• Molecule 1: Putative acetylglutamate kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	155.60 Å 155.60 Å 79.53 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.90 – 1.92 44.92 – 1.92	Depositor EDS
% Data completeness (in resolution range)	95.1 (19.90-1.92) 98.1 (44.92-1.92)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.94 (at 1.92 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.222 , 0.237 0.228 , 0.244	Depositor DCC
R_{free} test set	2464 reflections (3.00%)	DCC
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.6	EDS
Estimated twinning fraction	0.022 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 163557 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4126	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/1888	0.64	0/2551
1	C	0.31	0/2020	0.59	1/2734 (0.0%)
All	All	0.34	0/3908	0.62	1/5285 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	58	PRO	N-CA-CB	5.26	109.61	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1864	0	1921	36	0
1	C	1992	0	2039	45	0
2	A	10	0	0	0	0
2	C	10	0	0	0	0
3	A	105	0	0	3	0
3	C	145	0	0	3	0
All	All	4126	0	3960	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:LEU:HB3	1:A:228:MET:HE1	1.23	1.13
1:C:35:HIS:HE1	1:C:83:ASN:HD22	1.23	0.84
1:A:196:LEU:HD12	1:A:266:THR:HG21	1.62	0.82
1:A:55:LEU:HD11	1:A:65:LEU:HD13	1.67	0.76
1:A:212:VAL:CG2	1:A:269:ARG:HA	2.19	0.73
1:C:233:MET:O	1:C:237:GLU:HG2	1.89	0.71
1:A:230:ARG:HB3	3:A:359:HOH:O	1.90	0.71
1:C:215:ILE:HD12	1:C:215:ILE:H	1.57	0.69
1:A:197:LEU:HD13	1:A:202:ASP:O	1.92	0.69
1:C:116:VAL:HG13	1:C:127:HIS:HB3	1.74	0.68
1:C:110:GLY:HA3	1:C:134:THR:O	1.94	0.67
1:A:212:VAL:HG23	1:A:269:ARG:HA	1.76	0.66
1:A:110:GLY:HA3	1:A:134:THR:O	1.97	0.64
1:A:55:LEU:CD1	1:A:65:LEU:HD13	2.27	0.64
1:C:208:ARG:NH1	1:C:265:GLY:H	1.97	0.63
1:C:212:VAL:HG23	1:C:269:ARG:C	2.20	0.61
1:A:269:ARG:OXT	1:A:269:ARG:HG2	2.01	0.60
1:A:212:VAL:HG13	1:A:239:VAL:HB	1.84	0.60
1:C:215:ILE:HD12	1:C:215:ILE:N	2.17	0.60
1:C:4:VAL:HB	1:C:33:LEU:HD23	1.83	0.60
1:A:212:VAL:HG23	1:A:269:ARG:CA	2.32	0.59
1:A:212:VAL:HG23	1:A:269:ARG:C	2.23	0.59
1:C:212:VAL:N	1:C:269:ARG:OXT	2.33	0.57
1:A:244:LYS:O	1:A:269:ARG:HB2	2.05	0.56
1:C:227:ARG:HE	1:C:231:LYS:HZ1	1.52	0.56
1:C:212:VAL:HG13	1:C:236:VAL:HG13	1.88	0.56
1:C:212:VAL:HG23	1:C:269:ARG:OXT	2.06	0.55
1:C:65:LEU:HD11	1:C:127:HIS:CD2	2.41	0.55
1:C:2:ILE:HB	1:C:31:LEU:HD23	1.89	0.55
1:A:4:VAL:HB	1:A:33:LEU:HD23	1.90	0.53
1:C:208:ARG:HH12	1:C:265:GLY:H	1.56	0.52
1:C:196:LEU:HD13	1:C:248:PHE:CG	2.45	0.52
1:A:210:ILE:HG21	1:A:215:ILE:HG12	1.93	0.51
1:C:221:LEU:HD12	1:C:233:MET:SD	2.51	0.50
1:A:230:ARG:HD2	3:A:359:HOH:O	2.10	0.50
1:C:237:GLU:OE2	1:C:237:GLU:HA	2.12	0.50
1:A:253:VAL:O	1:A:256:PRO:HD3	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:ASN:N	1:A:256:PRO:CD	2.75	0.49
1:C:212:VAL:HG21	1:C:239:VAL:HG11	1.94	0.49
1:C:13:ASN:HD22	1:C:13:ASN:C	2.16	0.49
1:C:55:LEU:CD1	1:C:65:LEU:HD13	2.42	0.48
1:A:50:HIS:CD2	1:A:69:LYS:HZ3	2.31	0.48
1:A:50:HIS:HD2	1:A:69:LYS:HZ3	1.61	0.48
1:A:148:GLN:NE2	3:A:276:HOH:O	2.46	0.48
1:C:232:VAL:O	1:C:236:VAL:HG23	2.14	0.48
1:A:211:PRO:HA	1:A:269:ARG:O	2.14	0.47
1:A:55:LEU:HD11	1:A:65:LEU:CD1	2.40	0.47
1:A:229:LYS:O	1:A:233:MET:HG2	2.15	0.46
1:C:170:ASP:CG	1:C:173:GLN:HG2	2.35	0.46
1:C:227:ARG:HG3	1:C:228:MET:N	2.30	0.46
1:A:235:ALA:CB	1:A:246:VAL:HG11	2.46	0.46
1:A:198:ALA:O	1:A:225:GLN:NE2	2.37	0.46
1:C:55:LEU:HD11	1:C:65:LEU:HD13	1.99	0.45
1:C:170:ASP:OD1	1:C:173:GLN:HG2	2.17	0.45
1:C:111:ARG:HD3	3:C:315:HOH:O	2.16	0.45
1:C:106:ARG:HH12	1:C:164:ASN:HD21	1.65	0.44
1:C:170:ASP:HB3	1:C:173:GLN:HG2	2.00	0.44
1:C:98:ILE:HG23	1:C:98:ILE:O	2.18	0.43
1:A:50:HIS:HB3	1:A:69:LYS:HZ2	1.83	0.43
1:C:35:HIS:CE1	1:C:83:ASN:HD22	2.15	0.43
1:A:212:VAL:CG1	1:A:239:VAL:HB	2.48	0.43
1:A:53:ARG:HD2	1:A:65:LEU:HD23	2.00	0.43
1:C:111:ARG:NH1	3:C:352:HOH:O	2.51	0.43
1:A:50:HIS:HB3	1:A:69:LYS:NZ	2.34	0.43
1:C:227:ARG:HG3	1:C:228:MET:H	1.84	0.42
1:A:51:PRO:HA	1:A:52:PRO:HD3	1.89	0.42
1:C:176:ALA:O	1:C:180:THR:HG23	2.20	0.42
1:C:41:THR:HA	1:C:77:VAL:HG21	2.01	0.42
1:C:227:ARG:HH21	1:C:231:LYS:NZ	2.18	0.42
1:A:98:ILE:O	1:A:98:ILE:HG23	2.20	0.42
1:C:227:ARG:HE	1:C:231:LYS:NZ	2.18	0.42
1:C:107:LEU:HD21	1:C:155:LEU:HD22	2.02	0.41
1:C:35:HIS:HD2	1:C:36:GLY:O	2.04	0.41
1:C:255:ASN:N	1:C:256:PRO:CD	2.83	0.41
1:A:2:ILE:O	1:A:31:LEU:HD12	2.20	0.41
1:A:109:VAL:HA	1:A:160:LEU:O	2.20	0.41
1:C:173:GLN:CD	3:C:323:HOH:O	2.59	0.41
1:C:196:LEU:HD23	1:C:228:MET:HE3	2.02	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:259:ARG:O	1:C:264:GLU:HB2	2.21	0.40
1:A:268:VAL:O	1:A:269:ARG:HB3	2.22	0.40
1:C:259:ARG:HB3	1:C:264:GLU:OE1	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	240/269 (89%)	232 (97%)	7 (3%)	1 (0%)	39	27
1	C	262/269 (97%)	252 (96%)	9 (3%)	1 (0%)	39	27
All	All	502/538 (93%)	484 (96%)	16 (3%)	2 (0%)	39	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	37	GLY
1	C	37	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/212 (90%)	184 (96%)	7 (4%)	41	28

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	200/212 (94%)	195 (98%)	5 (2%)	55	46
All	All	391/424 (92%)	379 (97%)	12 (3%)	47	36

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	PRO
1	A	69	LYS
1	A	201	PRO
1	A	205	SER
1	A	206	LEU
1	A	215	ILE
1	A	269	ARG
1	C	13	ASN
1	C	114	THR
1	C	196	LEU
1	C	201	PRO
1	C	231	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	HIS
1	A	164	ASN
1	C	13	ASN
1	C	35	HIS
1	C	164	ASN
1	C	255	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	270	-	4,4,4	0.17	0	6,6,6	0.16	0
2	SO4	A	271	-	4,4,4	0.25	0	6,6,6	0.09	0
2	SO4	C	270	-	4,4,4	0.11	0	6,6,6	0.16	0
2	SO4	C	271	-	4,4,4	0.23	0	6,6,6	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	270	-	-	0/0/0/0	0/0/0/0
2	SO4	A	271	-	-	0/0/0/0	0/0/0/0
2	SO4	C	270	-	-	0/0/0/0	0/0/0/0
2	SO4	C	271	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/269 (92%)	0.76	30 (12%) 6 6	20, 37, 59, 65	0
1	C	266/269 (98%)	0.80	35 (13%) 4 5	16, 31, 60, 66	0
All	All	514/538 (95%)	0.78	65 (12%) 5 6	16, 34, 60, 66	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	118	TYR	7.6
1	A	55	LEU	6.9
1	A	215	ILE	6.8
1	C	204	ALA	6.3
1	A	57	HIS	6.3
1	C	55	LEU	6.2
1	A	204	ALA	6.2
1	A	56	THR	5.0
1	C	58	PRO	4.9
1	A	58	PRO	4.7
1	C	124	VAL	4.7
1	C	119	VAL	4.6
1	A	216	GLU	4.5
1	C	123	LYS	4.2
1	A	206	LEU	4.1
1	A	221	LEU	4.1
1	A	212	VAL	4.1
1	C	128	ARG	4.0
1	A	65	LEU	4.0
1	A	202	ASP	3.9
1	C	116	VAL	3.6
1	A	269	ARG	3.5
1	A	217	ASP	3.4
1	A	54	PHE	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	200	TYR	3.3
1	C	217	ASP	3.3
1	A	222	ALA	3.3
1	A	68	ARG	3.3
1	C	236	VAL	3.2
1	A	227	ARG	3.2
1	A	203	GLU	3.2
1	C	126	VAL	3.2
1	C	127	HIS	3.1
1	A	198	ALA	3.1
1	C	56	THR	2.9
1	A	62	VAL	2.9
1	A	220	TYR	2.9
1	C	65	LEU	2.8
1	C	198	ALA	2.8
1	C	114	THR	2.7
1	A	53	ARG	2.7
1	C	53	ARG	2.6
1	C	57	HIS	2.6
1	C	223	LEU	2.5
1	C	215	ILE	2.5
1	A	240	LYS	2.5
1	C	226	GLY	2.5
1	A	219	GLU	2.5
1	C	63	SER	2.5
1	C	227	ARG	2.5
1	C	62	VAL	2.4
1	A	201	PRO	2.4
1	C	113	LYS	2.3
1	C	115	ALA	2.3
1	C	50	HIS	2.3
1	C	54	PHE	2.3
1	C	216	GLU	2.3
1	C	129	GLY	2.3
1	C	214	ARG	2.3
1	C	173	GLN	2.2
1	A	63	SER	2.2
1	A	259	ARG	2.2
1	A	262	SER	2.1
1	C	125	LYS	2.1
1	A	254	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	SO4	A	271	5/5	0.89	0.21	6.60	36,37,38,39	5
2	SO4	A	270	5/5	0.97	0.12	1.93	37,38,38,42	0
2	SO4	C	271	5/5	0.83	0.22	1.71	32,36,37,37	5
2	SO4	C	270	5/5	0.99	0.11	0.93	32,34,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.