



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:36 PM GMT

PDB ID : 3U9T
Title : Crystal structure of *P. aeruginosa* 3-methylcrotonyl-CoA carboxylase (MCC)
750 kD holoenzyme, free enzyme
Authors : Huang, C.S.; Tong, L.
Deposited on : 2011-10-19
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	655	
2	B	555	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 7635 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methylcrotonyl-CoA carboxylase, alpha-subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	558	4323	2693	815	797	18	0	0	0

- Molecule 2 is a protein called Methylcrotonyl-CoA carboxylase, beta-subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	437	3312	2101	600	592	19	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	8	MET	-	EXPRESSION TAG	UNP Q9I297
B	9	GLY	-	EXPRESSION TAG	UNP Q9I297
B	10	SER	-	EXPRESSION TAG	UNP Q9I297
B	11	SER	-	EXPRESSION TAG	UNP Q9I297
B	12	HIS	-	EXPRESSION TAG	UNP Q9I297
B	13	HIS	-	EXPRESSION TAG	UNP Q9I297
B	14	HIS	-	EXPRESSION TAG	UNP Q9I297
B	15	HIS	-	EXPRESSION TAG	UNP Q9I297
B	16	HIS	-	EXPRESSION TAG	UNP Q9I297
B	17	HIS	-	EXPRESSION TAG	UNP Q9I297
B	18	SER	-	EXPRESSION TAG	UNP Q9I297
B	19	SER	-	EXPRESSION TAG	UNP Q9I297
B	20	GLY	-	EXPRESSION TAG	UNP Q9I297
B	21	LEU	-	EXPRESSION TAG	UNP Q9I297
B	22	VAL	-	EXPRESSION TAG	UNP Q9I297
B	23	PRO	-	EXPRESSION TAG	UNP Q9I297
B	24	ARG	-	EXPRESSION TAG	UNP Q9I297
B	25	GLY	-	EXPRESSION TAG	UNP Q9I297
B	26	SER	-	EXPRESSION TAG	UNP Q9I297
B	27	HIS	-	EXPRESSION TAG	UNP Q9I297

4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	158.93Å 158.93Å 311.97Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.23 – 2.90 46.22 – 2.90	Depositor EDS
% Data completeness (in resolution range)	92.9 (46.23-2.90) 93.0 (46.22-2.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.91Å)	Xtrriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.209 , 0.269 0.208 , 0.271	Depositor DCC
R_{free} test set	1583 reflections (5.02%)	DCC
Wilson B-factor (Å ²)	70.2	Xtrriage
Anisotropy	0.205	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 54.4	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Outliers	0 of 33710 reflections	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7635	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/4406	0.76	1/5958 (0.0%)
2	B	0.59	0/3380	0.79	1/4580 (0.0%)
All	All	0.54	0/7786	0.78	2/10538 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	454	GLY	N-CA-C	5.66	127.24	113.10
1	A	547	LEU	CA-CB-CG	-5.26	103.20	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4323	0	4267	305	0
2	B	3312	0	3308	175	0
All	All	7635	0	7575	468	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

The worst 5 of 468 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:465:THR:HG22	1:A:467:LEU:H	1.15	1.11
1:A:274:ARG:HH22	1:A:320:THR:HG21	1.16	1.10
2:B:157:GLN:NE2	2:B:197:ASN:HD22	1.50	1.10
1:A:47:ARG:HB2	1:A:47:ARG:HH11	1.16	1.07
2:B:212:VAL:HB	2:B:232:MET:HE3	1.37	1.06

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	552/655 (84%)	433 (78%)	88 (16%)	31 (6%)	2 7
2	B	425/555 (77%)	374 (88%)	39 (9%)	12 (3%)	6 24
All	All	977/1210 (81%)	807 (83%)	127 (13%)	43 (4%)	3 12

5 of 43 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	145	ALA
1	A	164	SER
1	A	283	GLN
1	A	409	ALA
1	A	412	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/496 (86%)	390 (91%)	37 (9%)	13	36
2	B	331/418 (79%)	309 (93%)	22 (7%)	21	51
All	All	758/914 (83%)	699 (92%)	59 (8%)	16	41

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	446	GLU
1	A	559	ASP
2	B	455	ARG
1	A	452	LEU
1	A	525	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	540	ASN
2	B	64	HIS
2	B	403	ASN
1	A	617	GLN
1	A	272	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	558/655 (85%)	0.13	28 (5%) 32 26	48, 87, 147, 168	0
2	B	437/555 (78%)	-0.03	19 (4%) 39 32	40, 63, 123, 172	0
All	All	995/1210 (82%)	0.06	47 (4%) 35 29	40, 77, 142, 172	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	241	LEU	7.0
2	B	176	ALA	6.3
2	B	240	PHE	6.3
1	A	245	LEU	5.5
1	A	207	ALA	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.