



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:34 PM GMT

PDB ID : 1UA7  
Title : Crystal Structure Analysis of Alpha-Amylase from Bacillus Subtilis complexed with Acarbose  
Authors : Kagawa, M.; Fujimoto, Z.; Momma, M.; Takase, K.; Mizuno, H.  
Deposited on : 2003-03-03  
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4405 (2.24-2.20)
Clashscore	102246	5146 (2.24-2.20)
Ramachandran outliers	100387	5065 (2.24-2.20)
Sidechain outliers	100360	5066 (2.24-2.20)
RSRZ outliers	91569	4414 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div></div> <div>78%</div> <div>20%</div> <div>.</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3808 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-amylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	422	Total	C	N	O	S	0	0	0
			3303	2054	581	660	8			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	105	PHE	SER	SEE REMARK 999	UNP P00691
A	163	GLU	ASP	SEE REMARK 999	UNP P00691
A	356	GLN	ASN	ENGINEERED	UNP P00691

- Molecule 2 is a polymer of unknown type called SUGAR (6-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	6	Total	C	N	O	0	0
			65	38	2	25		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Ca	0	0
			3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	437	Total	O	0	0
			437	437		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Alpha-amylase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.32Å 74.20Å 115.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.97 – 2.21 45.63 – 2.21	Depositor EDS
% Data completeness (in resolution range)	95.5 (19.97-2.21) 95.6 (45.63-2.21)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.38 (at 2.20Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.208 , 0.265 0.212 , 0.269	Depositor DCC
$R_{free}$ test set	2959 reflections (9.98%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.0	Xtriage
Anisotropy	0.964	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 63.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	1 of 29787 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3808	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, ACI, GLC, GLD, G6D, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/3383	0.59	0/4596

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3303	0	3087	63	0
2	A	65	0	60	2	0
3	A	3	0	0	0	0
4	A	437	0	0	8	0
All	All	3808	0	3147	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:LYS:HE2	4:A:611:HOH:O	1.87	0.74
1:A:391:ASP:HA	1:A:405:VAL:HG23	1.76	0.68
1:A:127:ILE:HG13	1:A:140:SER:O	1.98	0.64
1:A:151:ASN:O	1:A:154:VAL:HG22	2.00	0.62
1:A:78:GLN:O	1:A:82:GLU:HG3	2.00	0.61
1:A:197:ILE:HD12	1:A:197:ILE:C	2.23	0.59
1:A:373:LEU:HB2	1:A:420:ALA:HB3	1.85	0.59
1:A:383:ILE:HG23	1:A:412:GLY:O	2.03	0.58
1:A:250:SER:O	1:A:296:ARG:NH2	2.36	0.57
1:A:392:GLY:N	1:A:405:VAL:HG23	2.20	0.57
1:A:221:ALA:HA	1:A:224:MET:O	2.05	0.56
1:A:14:ALA:CB	1:A:22:LEU:HD21	2.37	0.55
1:A:228:ALA:HB2	1:A:261:LEU:HG	1.90	0.54
1:A:362:PHE:HB3	1:A:374:ALA:HB3	1.90	0.53
2:A:506:BGC:H6C1	4:A:960:HOH:O	2.08	0.53
1:A:13:HIS:HB2	1:A:302:LEU:HD22	1.89	0.53
1:A:162:LEU:O	1:A:166:LEU:HD22	2.09	0.53
1:A:352:LEU:N	1:A:352:LEU:HD23	2.24	0.53
1:A:7:LYS:HE2	1:A:259:ASP:O	2.10	0.52
1:A:185:ASP:CG	1:A:216:ARG:HH22	2.14	0.52
1:A:247:SER:HB2	4:A:705:HOH:O	2.10	0.51
1:A:390:PRO:O	1:A:405:VAL:HG21	2.11	0.50
1:A:332:ASP:OD2	1:A:334:ALA:HB3	2.11	0.50
1:A:414:ILE:HG23	1:A:414:ILE:O	2.12	0.50
1:A:394:TYR:HE2	1:A:405:VAL:HG21	1.76	0.50
1:A:259:ASP:OD1	1:A:260:LYS:HG3	2.13	0.49
1:A:251:HIS:CD2	1:A:253:ALA:H	2.31	0.49
1:A:340:ARG:O	1:A:344:VAL:HG22	2.13	0.48
1:A:348:GLN:HG3	1:A:367:GLY:HA3	1.96	0.48
1:A:395:ASP:HB2	4:A:880:HOH:O	2.13	0.48
1:A:195:PRO:HG3	1:A:223:TYR:CE1	2.50	0.47
1:A:391:ASP:CA	1:A:405:VAL:HG23	2.44	0.47
1:A:151:ASN:HB3	1:A:154:VAL:HG13	1.97	0.47
1:A:102:HIS:CD2	1:A:144:LEU:HD13	2.51	0.46
1:A:270:THR:HB	1:A:277:GLU:HG3	1.98	0.45
1:A:348:GLN:HG3	1:A:349:PRO:HD2	1.98	0.45
1:A:25:ASN:O	1:A:29:ILE:HG13	2.17	0.44
1:A:37:ILE:HD12	1:A:37:ILE:C	2.36	0.44
1:A:345:MET:O	1:A:348:GLN:HB2	2.16	0.44
1:A:348:GLN:HE22	1:A:369:HIS:CE1	2.36	0.44
1:A:384:ASN:C	1:A:384:ASN:HD22	2.20	0.44
1:A:209:ILE:HD12	1:A:226:VAL:HG12	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:122:HIS:HB2	1:A:146:ASP:HB3	2.00	0.44
1:A:336:THR:O	1:A:340:ARG:HG3	2.18	0.43
1:A:405:VAL:HG23	1:A:405:VAL:O	2.18	0.43
1:A:158:LEU:O	1:A:161:PHE:HB3	2.18	0.43
1:A:383:ILE:O	1:A:411:THR:HA	2.19	0.43
1:A:62:TYR:HB3	2:A:504:ACI:O6	2.19	0.43
1:A:396:ASN:OD1	1:A:420:ALA:HA	2.19	0.43
1:A:81:LYS:HE2	1:A:168:ASP:OD1	2.19	0.43
1:A:122:HIS:NE2	1:A:148:ASN:HA	2.34	0.42
1:A:383:ILE:HD11	1:A:410:LEU:HD23	2.01	0.42
1:A:7:LYS:HD2	1:A:300:THR:OG1	2.18	0.42
1:A:394:TYR:HE2	1:A:405:VAL:CG2	2.32	0.42
1:A:76:THR:OG1	1:A:79:GLU:HG3	2.20	0.42
1:A:248:ASN:HB2	4:A:1010:HOH:O	2.19	0.42
1:A:6:ILE:HG13	4:A:1004:HOH:O	2.19	0.42
1:A:194:TRP:CZ3	1:A:197:ILE:HD11	2.54	0.42
1:A:349:PRO:O	1:A:366:ARG:HA	2.20	0.41
1:A:377:GLY:HA3	4:A:700:HOH:O	2.20	0.41
1:A:296:ARG:HD2	1:A:350:GLU:OE1	2.21	0.41
1:A:35:THR:HB	4:A:889:HOH:O	2.19	0.41
1:A:421:VAL:C	1:A:422:LEU:HD13	2.41	0.40
1:A:345:MET:O	1:A:346:ALA:C	2.60	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	420/422 (100%)	405 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.



### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	351/351 (100%)	338 (96%)	13 (4%)	41	50

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	SER
1	A	166	LEU
1	A	225	ASP
1	A	242	ARG
1	A	261	LEU
1	A	264	TRP
1	A	302	LEU
1	A	348	GLN
1	A	352	LEU
1	A	356	GLN
1	A	383	ILE
1	A	384	ASN
1	A	422	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	112	ASN
1	A	126	GLN
1	A	153	GLN
1	A	342	HIS
1	A	348	GLN
1	A	356	GLN
1	A	384	ASN
1	A	406	ASN
1	A	415	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

6 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	ACI	A	501	2	11,12,12	5.11	5 (45%)	9,17,17	0.97	0
2	GLD	A	502	2	10,10,10	2.46	5 (50%)	13,14,14	0.99	0
2	GLC	A	503	2	11,11,12	1.93	5 (45%)	15,15,17	1.03	1 (6%)
2	ACI	A	504	2	10,11,12	2.91	5 (50%)	8,15,17	1.67	1 (12%)
2	G6D	A	505	2	10,10,11	2.10	5 (50%)	13,14,16	1.06	0
2	BGC	A	506	2	11,11,12	1.90	4 (36%)	15,15,17	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACI	A	501	2	-	0/2/22/22	0/1/1/1
2	GLD	A	502	2	-	0/0/16/16	0/1/1/1
2	GLC	A	503	2	-	0/2/18/22	0/1/1/1
2	ACI	A	504	2	-	0/1/18/22	0/1/1/1
2	G6D	A	505	2	-	0/0/16/20	0/1/1/1
2	BGC	A	506	2	-	0/2/18/22	0/1/1/1

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	506	BGC	C3-C2	2.02	1.55	1.52
2	A	503	GLC	O5-C1	2.03	1.46	1.43
2	A	506	BGC	O5-C5	2.10	1.48	1.44
2	A	501	ACI	C4-C5	2.15	1.57	1.50
2	A	501	ACI	C3-C2	2.35	1.58	1.52
2	A	505	G6D	C4A-C5A	2.37	1.56	1.51
2	A	503	GLC	C4-C5	2.40	1.58	1.52
2	A	505	G6D	C3A-C2	2.46	1.55	1.52
2	A	504	ACI	C4-C3	2.46	1.58	1.52
2	A	505	G6D	O5-C5A	2.50	1.49	1.44
2	A	503	GLC	O5-C5	2.63	1.49	1.44
2	A	502	GLD	O5-C5	2.72	1.49	1.44
2	A	503	GLC	C3-C2	2.84	1.56	1.52
2	A	506	BGC	C4-C3	2.87	1.57	1.52
2	A	504	ACI	C4-C5	2.92	1.56	1.50
2	A	505	G6D	C4A-C3A	3.06	1.57	1.52
2	A	502	GLD	C3-C2	3.07	1.56	1.52
2	A	502	GLD	O5-C1	3.11	1.48	1.43
2	A	505	G6D	O5-C1	3.21	1.49	1.43
2	A	503	GLC	C4-C3	3.32	1.58	1.52
2	A	501	ACI	C1-C7	3.64	1.58	1.49
2	A	506	BGC	O5-C1	3.69	1.50	1.43
2	A	502	GLD	C4-C3	3.71	1.58	1.52
2	A	502	GLD	C4-C5	3.85	1.59	1.51
2	A	504	ACI	C1-C7	3.97	1.59	1.49
2	A	501	ACI	C3-C4	4.27	1.58	1.53
2	A	504	ACI	C3-C2	4.54	1.58	1.52
2	A	504	ACI	C7-C5	5.37	1.53	1.33
2	A	501	ACI	C7-C5	15.52	1.56	1.32

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	504	ACI	O6-C6-C5	-3.79	103.67	111.75
2	A	503	GLC	C3-C4-C5	-2.32	106.70	111.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	504	ACI	1	0
2	A	506	BGC	1	0

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	422/422 (100%)	-0.34	6 (1%) 78 77	9, 19, 34, 56	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	405	VAL	3.7
1	A	406	ASN	2.8
1	A	390	PRO	2.8
1	A	407	ASP	2.6
1	A	393	ARG	2.2
1	A	369	HIS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	BGC	A	506	11/12	0.91	0.14	1.36	21,29,41,50	0
2	ACI	A	504	11/12	0.95	0.14	0.78	12,15,18,19	0
2	GLC	A	503	11/12	0.97	0.10	-0.66	10,17,21,21	0
2	G6D	A	505	10/11	0.97	0.10	-0.81	16,21,24,26	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GLD	A	502	10/10	0.95	0.09	-1.00	19,22,24,26	0
2	ACI	A	501	12/12	0.95	0.10	-	18,24,29,34	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	A	602	1/1	0.99	0.05	-1.75	19,19,19,19	0
3	CA	A	601	1/1	0.99	0.03	-3.80	13,13,13,13	0
3	CA	A	603	1/1	0.99	0.03	-	11,11,11,11	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.