



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 07:32 PM GMT

PDB ID : 4UFC
Title : Crystal structure of the GH95 enzyme BACOVA_03438
Authors : Rogowski, A.; Briggs, J.A.; Mortimer, J.C.; Tryfona, T.; Terrapon, N.; Lowe, E.C.; Basle, A.; Morland, C.; Day, A.M.; Zheng, H.; Rogers, T.E.; Thompson, P.; Hawkins, A.R.; Yadav, M.P.; Henrissat, B.; Martens, E.C.; Dupree, P.; Gilbert, H.J.; Bolam, D.N.
Deposited on : 2015-03-16
Resolution : 2.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2676 (2.84-2.80)
Clashscore	102246	3124 (2.84-2.80)
Ramachandran outliers	100387	3072 (2.84-2.80)
Sidechain outliers	100360	3074 (2.84-2.80)
RSRZ outliers	91569	2690 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	811	 90% 7% •
1	B	811	 90% 7% •

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 12568 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GH95.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	787	Total	C	N	O	S	0	0	0
			6261	3980	1087	1169	25			
1	B	788	Total	C	N	O	S	0	0	0
			6270	3985	1089	1171	25			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Ca	0	0
			2	2		
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is SUGAR (BETA-L-GALACTOPYRANOSE) (three-letter code: GIV) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			12	6	6		
3	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is CACODYLATE ION (three-letter code: CAC) (formula: $\text{C}_2\text{H}_6\text{AsO}_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	As	C	O	0	0
			5	1	2	2		
4	B	1	Total	As	C	O	0	0
			5	1	2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GH95



- Molecule 1: GH95



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.69Å 179.12Å 205.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	134.93 – 2.81 49.31 – 2.81	Depositor EDS
% Data completeness (in resolution range)	97.4 (134.93-2.81) 97.4 (49.31-2.81)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.186 , 0.228 0.190 , 0.228	Depositor DCC
R_{free} test set	2441 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	44.1	Xtriage
Anisotropy	0.811	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 22.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 48216 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12568	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, GIV, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/6425	0.69	1/8740 (0.0%)
1	B	0.52	0/6434	0.69	3/8752 (0.0%)
All	All	0.53	0/12859	0.69	4/17492 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	385	THR	N-CA-C	-6.51	93.42	111.00
1	B	659	ASN	CB-CA-C	-6.34	97.72	110.40
1	B	421	HIS	CB-CA-C	-6.07	98.25	110.40
1	A	659	ASN	CB-CA-C	-5.95	98.50	110.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	688	ASP	Peptide
1	B	688	ASP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6261	0	6102	32	0
1	B	6270	0	6110	34	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
3	A	12	0	12	0	0
3	B	12	0	12	0	0
4	A	5	0	0	1	0
4	B	5	0	0	1	0
All	All	12568	0	12236	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:20:GLN:NE2	1:B:139:GLU:OE2	1.94	1.00
1:B:337:ARG:HH11	1:B:385:THR:HG23	1.38	0.86
1:B:36:ALA:O	1:B:659:ASN:ND2	2.20	0.74
1:A:24:LEU:HA	1:A:313:THR:HG21	1.71	0.73
1:A:574:HIS:O	1:A:606:ARG:NH1	2.22	0.72
1:B:24:LEU:HA	1:B:313:THR:HG21	1.71	0.71
1:B:574:HIS:O	1:B:606:ARG:NH1	2.24	0.70
1:A:41:ASN:O	1:A:42:SER:HB3	1.90	0.70
1:B:337:ARG:HD2	1:B:385:THR:HG23	1.79	0.63
1:B:112:SER:HA	1:B:366:ASP:HB3	1.83	0.61
1:B:104:PHE:CD2	1:B:440:MET:HE1	2.38	0.58
1:A:104:PHE:CD2	1:A:440:MET:HE1	2.38	0.58
1:A:112:SER:HA	1:A:366:ASP:HB3	1.85	0.58
1:B:381:PRO:O	1:B:385:THR:HB	2.07	0.55
1:B:486:HIS:CE1	1:B:488:VAL:HG22	2.42	0.54
1:B:41:ASN:O	1:B:42:SER:HB3	2.08	0.53
1:B:664:HIS:N	1:B:665:PRO:HA	2.24	0.52
1:B:104:PHE:CD2	1:B:440:MET:CE	2.92	0.52
1:B:41:ASN:O	1:B:42:SER:CB	2.57	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:752:LEU:HD12	1:A:794:ILE:HG12	1.92	0.52
1:A:104:PHE:CD2	1:A:440:MET:CE	2.92	0.51
1:B:752:LEU:HD12	1:B:794:ILE:HG12	1.92	0.50
1:A:664:HIS:N	1:A:665:PRO:HA	2.27	0.50
1:A:68:SER:OG	1:A:213:LYS:O	2.30	0.49
1:B:366:ASP:OD2	4:B:1811:CAC:AS	2.90	0.49
1:A:51:GLY:O	1:A:156:ARG:NH2	2.46	0.48
1:B:609:LYS:O	1:B:610:ALA:HB3	2.14	0.48
1:A:609:LYS:O	1:A:610:ALA:HB3	2.13	0.48
1:B:68:SER:OG	1:B:213:LYS:O	2.31	0.47
1:B:500:PRO:HA	1:B:501:GLU:HA	1.78	0.47
1:B:337:ARG:HG2	1:B:385:THR:HG21	1.97	0.47
1:B:383:GLU:O	1:B:385:THR:O	2.32	0.47
1:A:368:LYS:HG2	1:A:424:THR:O	2.16	0.46
1:B:368:LYS:HG2	1:B:424:THR:O	2.16	0.46
1:A:36:ALA:O	1:A:659:ASN:OD1	2.34	0.46
1:A:187:CYS:SG	1:A:188:PRO:HD2	2.56	0.45
1:A:500:PRO:HA	1:A:501:GLU:HA	1.77	0.45
1:B:664:HIS:HB3	1:B:665:PRO:C	2.37	0.45
1:A:132:PHE:CE2	1:A:134:ARG:HG3	2.52	0.44
1:B:511:MET:SD	1:B:576:SER:HB3	2.57	0.44
1:B:732:ARG:NH1	1:B:734:ASN:OD1	2.50	0.44
1:B:110:GLY:HA3	1:B:368:LYS:HD2	1.99	0.44
1:A:659:ASN:HB3	1:A:661:LEU:HG	2.00	0.44
1:A:41:ASN:O	1:A:42:SER:CB	2.64	0.43
1:A:366:ASP:OD2	4:A:1810:CAC:AS	2.96	0.43
1:A:689:GLY:HA2	1:A:765:PHE:O	2.18	0.43
1:A:752:LEU:CD1	1:A:794:ILE:HD11	2.48	0.43
1:A:143:THR:CG2	1:A:158:THR:HB	2.48	0.43
1:B:187:CYS:SG	1:B:188:PRO:HD2	2.59	0.43
1:A:511:MET:SD	1:A:576:SER:HB3	2.58	0.42
1:A:25:TRP:H	1:A:313:THR:HG23	1.84	0.42
1:A:104:PHE:HD2	1:A:440:MET:HE1	1.84	0.42
1:B:370:THR:O	1:B:374:ASN:HB2	2.19	0.42
1:B:752:LEU:CD1	1:B:794:ILE:HD11	2.49	0.42
1:A:370:THR:O	1:A:374:ASN:HB2	2.20	0.42
1:B:193:VAL:CG1	1:B:200:LEU:HD11	2.50	0.42
1:A:313:THR:N	1:A:314:PRO:CD	2.83	0.41
1:A:110:GLY:HA3	1:A:368:LYS:HD2	2.01	0.41
1:B:65:TRP:CD1	1:B:426:LEU:HD22	2.55	0.41
1:A:193:VAL:CG1	1:A:200:LEU:HD11	2.49	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:353:LEU:HB2	1:B:367:SER:HA	2.02	0.41
1:A:353:LEU:HB2	1:A:367:SER:HA	2.02	0.41
1:A:582:TYR:CG	1:A:583:PRO:HA	2.56	0.41
1:B:380:TRP:N	1:B:381:PRO:CD	2.84	0.41
1:B:689:GLY:HA2	1:B:765:PHE:O	2.20	0.40
1:A:664:HIS:HB3	1:A:665:PRO:C	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	785/811 (97%)	750 (96%)	33 (4%)	2 (0%)	46	78
1	B	786/811 (97%)	746 (95%)	38 (5%)	2 (0%)	46	78
All	All	1571/1622 (97%)	1496 (95%)	71 (4%)	4 (0%)	46	78

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	373	ILE
1	B	373	ILE
1	A	42	SER
1	B	42	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	669/691 (97%)	658 (98%)	11 (2%)	70	92
1	B	670/691 (97%)	661 (99%)	9 (1%)	76	94
All	All	1339/1382 (97%)	1319 (98%)	20 (2%)	72	93

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	42	SER
1	A	182	THR
1	A	196	GLN
1	A	213	LYS
1	A	311	LEU
1	A	502	HIS
1	A	556	ASN
1	A	702	GLU
1	A	759	GLU
1	A	783	SER
1	B	20	GLN
1	B	58	GLN
1	B	231	ARG
1	B	385	THR
1	B	422	HIS
1	B	502	HIS
1	B	556	ASN
1	B	777	LYS
1	B	807	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	372	ASN
1	A	423	ASN
1	B	423	ASN
1	B	556	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GIV	A	1809	-	12,12,12	1.15	1 (8%)	17,17,17	1.64	4 (23%)
4	CAC	A	1810	-	0,4,4	0.00	-	0,6,6	0.00	-
3	GIV	B	1810	-	12,12,12	0.94	0	17,17,17	1.42	3 (17%)
4	CAC	B	1811	-	0,4,4	0.00	-	0,6,6	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GIV	A	1809	-	-	0/2/22/22	0/1/1/1
4	CAC	A	1810	-	-	0/0/0/0	0/0/0/0
3	GIV	B	1810	-	-	0/2/22/22	0/1/1/1
4	CAC	B	1811	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1809	GIV	O1-C1	2.12	1.47	1.39

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1809	GIV	O2-C2-C3	-2.79	104.07	110.36
3	B	1810	GIV	C3-C4-C5	-2.53	105.72	110.23
3	A	1809	GIV	O1-C1-O5	-2.43	103.56	110.33
3	B	1810	GIV	O2-C2-C3	-2.15	105.51	110.36
3	A	1809	GIV	O4-C4-C5	2.20	115.02	109.23
3	B	1810	GIV	C1-C2-C3	3.25	116.01	110.68
3	A	1809	GIV	C1-C2-C3	3.74	116.81	110.68

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1810	CAC	1	0
4	B	1811	CAC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	787/811 (97%)	-0.61	0 100 100	28, 38, 56, 93	0
1	B	788/811 (97%)	-0.46	4 (0%) 91 88	30, 44, 69, 100	0
All	All	1575/1622 (97%)	-0.53	4 (0%) 94 92	28, 41, 65, 100	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	756	ASN	5.4
1	B	127	GLN	2.8
1	B	757	GLY	2.7
1	B	197	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	CAC	B	1811	5/5	0.97	0.13	0.02	67,70,77,82	0
3	GIV	B	1810	12/12	0.96	0.12	-0.74	26,30,32,33	0
4	CAC	A	1810	5/5	0.95	0.11	-1.19	58,64,65,78	0
3	GIV	A	1809	12/12	0.97	0.11	-1.66	16,17,18,18	0
2	CA	A	1808	1/1	0.94	0.08	-2.83	55,55,55,55	0
2	CA	B	1809	1/1	0.90	0.09	-3.40	78,78,78,78	0
2	CA	B	1808	1/1	0.87	0.07	-	48,48,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.