



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:39 PM GMT

PDB ID : 3UK1
Title : Crystal structure of a transketolase from Burkholderia thailandensis with an oxidized cysteinesulfonic acid in the active site
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2011-11-08
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	711	
1	B	711	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	B	692	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10538 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transketolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	651	Total	C	N	O	S	0	1	0
			4894	3096	864	916	18			
1	B	669	Total	C	N	O	S	0	0	0
			4991	3150	887	936	18			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	EXPRESSION TAG	UNP Q2SZA7
A	-19	ALA	-	EXPRESSION TAG	UNP Q2SZA7
A	-18	HIS	-	EXPRESSION TAG	UNP Q2SZA7
A	-17	HIS	-	EXPRESSION TAG	UNP Q2SZA7
A	-16	HIS	-	EXPRESSION TAG	UNP Q2SZA7
A	-15	HIS	-	EXPRESSION TAG	UNP Q2SZA7
A	-14	HIS	-	EXPRESSION TAG	UNP Q2SZA7
A	-13	HIS	-	EXPRESSION TAG	UNP Q2SZA7
A	-12	MET	-	EXPRESSION TAG	UNP Q2SZA7
A	-11	GLY	-	EXPRESSION TAG	UNP Q2SZA7
A	-10	THR	-	EXPRESSION TAG	UNP Q2SZA7
A	-9	LEU	-	EXPRESSION TAG	UNP Q2SZA7
A	-8	GLU	-	EXPRESSION TAG	UNP Q2SZA7
A	-7	ALA	-	EXPRESSION TAG	UNP Q2SZA7
A	-6	GLN	-	EXPRESSION TAG	UNP Q2SZA7
A	-5	THR	-	EXPRESSION TAG	UNP Q2SZA7
A	-4	GLN	-	EXPRESSION TAG	UNP Q2SZA7
A	-3	GLY	-	EXPRESSION TAG	UNP Q2SZA7
A	-2	PRO	-	EXPRESSION TAG	UNP Q2SZA7
A	-1	GLY	-	EXPRESSION TAG	UNP Q2SZA7
A	0	SER	-	EXPRESSION TAG	UNP Q2SZA7
B	-20	MET	-	EXPRESSION TAG	UNP Q2SZA7
B	-19	ALA	-	EXPRESSION TAG	UNP Q2SZA7
B	-18	HIS	-	EXPRESSION TAG	UNP Q2SZA7
B	-17	HIS	-	EXPRESSION TAG	UNP Q2SZA7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	EXPRESSION TAG	UNP Q2SZA7
B	-15	HIS	-	EXPRESSION TAG	UNP Q2SZA7
B	-14	HIS	-	EXPRESSION TAG	UNP Q2SZA7
B	-13	HIS	-	EXPRESSION TAG	UNP Q2SZA7
B	-12	MET	-	EXPRESSION TAG	UNP Q2SZA7
B	-11	GLY	-	EXPRESSION TAG	UNP Q2SZA7
B	-10	THR	-	EXPRESSION TAG	UNP Q2SZA7
B	-9	LEU	-	EXPRESSION TAG	UNP Q2SZA7
B	-8	GLU	-	EXPRESSION TAG	UNP Q2SZA7
B	-7	ALA	-	EXPRESSION TAG	UNP Q2SZA7
B	-6	GLN	-	EXPRESSION TAG	UNP Q2SZA7
B	-5	THR	-	EXPRESSION TAG	UNP Q2SZA7
B	-4	GLN	-	EXPRESSION TAG	UNP Q2SZA7
B	-3	GLY	-	EXPRESSION TAG	UNP Q2SZA7
B	-2	PRO	-	EXPRESSION TAG	UNP Q2SZA7
B	-1	GLY	-	EXPRESSION TAG	UNP Q2SZA7
B	0	SER	-	EXPRESSION TAG	UNP Q2SZA7

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Cl 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	410	Total 410	O 410	0	0
4	B	232	Total 232	O 232	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Transketolase



• Molecule 1: Transketolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	145.35Å 145.35Å 142.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.15 47.95 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-2.15) 99.2 (47.95-2.15)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.48 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.185 , 0.225 0.184 , 0.224	Depositor DCC
R_{free} test set	4121 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.2	EDS
Estimated twinning fraction	0.025 for -h,-l,-k 0.024 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 82279 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10538	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OCS, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	6/5004 (0.1%)	0.76	0/6818
1	B	0.69	8/5103 (0.2%)	0.71	1/6958 (0.0%)
All	All	0.73	14/10107 (0.1%)	0.74	1/13776 (0.0%)

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	516	TRP	CD2-CE2	5.82	1.48	1.41
1	B	645	TRP	CD2-CE2	5.60	1.48	1.41
1	B	230	TRP	CD2-CE2	5.57	1.48	1.41
1	B	313	TRP	CD2-CE2	5.45	1.47	1.41
1	B	426	TRP	CD2-CE2	5.43	1.47	1.41
1	A	216	TRP	CD2-CE2	5.39	1.47	1.41
1	A	409	TRP	CD2-CE2	5.38	1.47	1.41
1	A	298	TRP	CD2-CE2	5.33	1.47	1.41
1	A	230	TRP	CD2-CE2	5.32	1.47	1.41
1	B	409	TRP	CD2-CE2	5.16	1.47	1.41
1	A	326	TRP	CD2-CE2	5.09	1.47	1.41
1	B	570	TRP	CD2-CE2	5.08	1.47	1.41
1	B	326	TRP	CD2-CE2	5.06	1.47	1.41
1	B	298	TRP	CD2-CE2	5.03	1.47	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	219	ASP	CB-CG-OD2	-5.16	113.65	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4894	0	4713	29	0
1	B	4991	0	4769	34	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	B	1	0	0	0	0
4	A	410	0	0	3	0
4	B	232	0	0	0	0
All	All	10538	0	9482	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:373:THR:HG22	1:B:549:PHE:H	1.23	1.03
1:A:252:LYS:HE2	4:A:921:HOH:O	1.83	0.77
1:A:610:MET:HE1	1:A:645:TRP:HZ3	1.51	0.74
1:A:223:LYS:HE2	1:B:226:GLU:HB2	1.71	0.71
1:A:610:MET:HE1	1:A:645:TRP:CZ3	2.29	0.67
1:A:223:LYS:HE3	1:B:223:LYS:HG3	1.78	0.66
1:B:373:THR:HG22	1:B:549:PHE:N	2.06	0.62
1:B:222:PRO:O	1:B:226:GLU:HG3	2.02	0.59
1:A:150:GLU:HG3	1:A:167:HIS:CD2	2.39	0.57
1:B:376:THR:HG23	1:B:547:LEU:HD12	1.87	0.57
1:A:472:VAL:HG11	4:A:903:HOH:O	2.06	0.56
1:A:376:THR:HG23	1:A:547:LEU:HD12	1.87	0.56
1:A:652:GLU:OE1	1:A:652:GLU:N	2.25	0.55
1:A:210:ASP:HB2	1:B:400:ASP:OD1	2.09	0.53
1:A:215:ASN:HB2	1:A:216:TRP:CE3	2.43	0.53
1:A:610:MET:CE	1:A:645:TRP:CZ3	2.91	0.53
1:B:306:PRO:HG2	1:B:309:VAL:HG23	1.91	0.53
1:A:223:LYS:HG3	1:B:223:LYS:HE3	1.90	0.53
1:A:49:MET:HE3	1:A:82:LEU:HD11	1.93	0.50
1:A:469:ALA:HA	1:A:472:VAL:HG12	1.91	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:489:ILE:HA	1:A:499:GLN:HG2	1.92	0.50
1:B:49:MET:C	1:B:49:MET:SD	2.91	0.49
1:B:489:ILE:HA	1:B:499:GLN:HG2	1.93	0.49
1:A:305:ILE:HB	1:A:310:TYR:CE2	2.48	0.48
1:B:166:ASP:OD1	1:B:197:LYS:NZ	2.44	0.48
1:B:381:GLN:HA	1:B:384:ILE:HD12	1.96	0.48
1:B:570:TRP:CD2	1:B:578:LYS:HE3	2.49	0.47
1:A:534:HIS:HB2	4:A:789:HOH:O	2.15	0.47
1:A:476:MET:O	1:A:477:LYS:HB2	2.16	0.45
1:A:49:MET:C	1:A:49:MET:SD	2.95	0.45
1:A:88:SER:HB3	1:A:121:PRO:HD3	1.98	0.45
1:B:625:GLU:OE2	1:B:629:PRO:HA	2.17	0.44
1:A:184:HIS:CE1	1:B:224:ARG:CZ	3.01	0.44
1:A:512:ASN:OD1	1:A:536:HIS:ND1	2.51	0.44
1:A:461:THR:HG23	1:A:503:HIS:CE1	2.52	0.44
1:A:224:ARG:CZ	1:B:184:HIS:CE1	3.01	0.44
1:B:468:ASN:O	1:B:472:VAL:HG23	2.18	0.43
1:B:562:LYS:HA	1:B:619:GLN:NE2	2.33	0.43
1:B:461:THR:HG23	1:B:503:HIS:CE1	2.54	0.43
1:B:106:GLU:HG3	1:B:116:LYS:HD2	2.00	0.43
1:B:474:ALA:HA	1:B:537:PRO:HG3	2.01	0.42
1:B:633:ARG:C	1:B:634:ARG:HG3	2.39	0.42
1:B:391:LEU:HD11	1:B:532:VAL:HG11	2.00	0.42
1:A:49:MET:HB3	1:A:50:PRO:HD3	2.01	0.42
1:B:457:GLY:HA2	1:B:483:VAL:O	2.20	0.42
1:A:49:MET:HE1	1:A:53:MET:SD	2.60	0.42
1:B:522:VAL:O	1:B:526:VAL:HG23	2.20	0.41
1:B:646:ARG:HA	1:B:646:ARG:HD3	1.92	0.41
1:B:110:PHE:CE2	1:B:111:ARG:HD2	2.55	0.41
1:B:191:GLY:HA3	1:B:228:TYR:O	2.19	0.41
1:A:457:GLY:HA2	1:A:483:VAL:O	2.21	0.41
1:B:86:HIS:HA	1:B:120:HIS:CD2	2.55	0.41
1:B:131:THR:HG23	1:B:133:THR:HG23	2.04	0.40
1:B:499:GLN:OE1	1:B:662:GLY:HA3	2.21	0.40
1:B:460:LEU:HA	1:B:484:PHE:HB3	2.04	0.40
1:A:383:THR:HG21	1:A:528:TRP:HB2	2.03	0.40
1:B:383:THR:HG21	1:B:528:TRP:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	647/711 (91%)	631 (98%)	16 (2%)	0	100	100
1	B	666/711 (94%)	647 (97%)	19 (3%)	0	100	100
All	All	1313/1422 (92%)	1278 (97%)	35 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	482/547 (88%)	477 (99%)	5 (1%)	82	87
1	B	486/547 (89%)	480 (99%)	6 (1%)	78	83
All	All	968/1094 (88%)	957 (99%)	11 (1%)	80	85

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	MET
1	A	84	ASN
1	A	420	ASP
1	A	470	LEU
1	A	500	SER
1	B	49	MET
1	B	67	HIS
1	B	322	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	345	GLU
1	B	390	VAL
1	B	592	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	231	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	OCS	A	177	1	7,8,9	3.13	2 (28%)	7,11,13	1.20	1 (14%)
1	OCS	B	177	1	7,8,9	3.74	2 (28%)	7,11,13	2.88	3 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	A	177	1	-	0/4/7/9	0/0/0/0
1	OCS	B	177	1	-	0/4/7/9	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	177	OCS	CB-SG	-7.77	1.66	1.77
1	A	177	OCS	CB-SG	-4.78	1.70	1.77
1	B	177	OCS	OD2-SG	5.85	1.61	1.46
1	A	177	OCS	OD2-SG	6.51	1.63	1.46

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	177	OCS	CB-CA-C	-2.65	104.20	111.46
1	A	177	OCS	O-C-CA	-2.12	119.97	125.49
1	B	177	OCS	OD2-SG-OD3	-2.03	106.90	111.61
1	B	177	OCS	OD1-SG-CB	6.43	112.36	106.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	691	-	4,4,4	0.13	0	6,6,6	0.89	0
2	SO4	B	691	-	4,4,4	0.31	0	6,6,6	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	691	-	-	0/0/0/0	0/0/0/0
2	SO4	B	691	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	650/711 (91%)	0.33	65 (10%)	9	15	20, 29, 49, 70	0
1	B	668/711 (93%)	0.39	62 (9%)	11	17	25, 39, 60, 97	0
All	All	1318/1422 (92%)	0.36	127 (9%)	10	16	20, 34, 57, 97	0

All (127) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	162	ALA	6.2
1	B	575	VAL	5.1
1	A	437	PHE	4.8
1	A	182	ILE	4.8
1	A	470	LEU	4.7
1	B	182	ILE	4.6
1	B	437	PHE	4.5
1	B	470	LEU	4.3
1	A	482	PHE	4.0
1	A	290	ALA	4.0
1	A	465	TYR	4.0
1	B	574	ILE	4.0
1	A	466	SER	3.9
1	A	286	ALA	3.9
1	A	462	PHE	3.9
1	A	443	ILE	3.8
1	A	211	GLY	3.8
1	B	419	ALA	3.7
1	A	575	VAL	3.7
1	A	507	LEU	3.5
1	B	465	TYR	3.5
1	B	631	GLY	3.5
1	A	690	ALA	3.5
1	A	295	ALA	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	469	ALA	3.5
1	A	510	ILE	3.4
1	A	421	GLY	3.4
1	B	573	GLU	3.4
1	A	574	ILE	3.4
1	B	576	ALA	3.4
1	B	154	ALA	3.3
1	A	441	ALA	3.3
1	A	294	GLU	3.3
1	A	292	THR	3.3
1	A	434	VAL	3.2
1	A	463	SER	3.2
1	A	458	THR	3.2
1	B	420	ASP	3.1
1	B	342	ALA	3.1
1	A	420	ASP	3.1
1	A	439	MET	3.1
1	B	159	ARG	3.1
1	B	460	LEU	3.0
1	B	462	PHE	3.0
1	B	181	GLY	3.0
1	B	505	ALA	3.0
1	B	164	ILE	3.0
1	B	468	ASN	3.0
1	B	469	ALA	3.0
1	A	136	LEU	2.9
1	A	484	PHE	2.9
1	B	440	SER	2.9
1	B	153	LEU	2.9
1	A	181	GLY	2.9
1	A	467	ARG	2.9
1	B	467	ARG	2.8
1	A	301	ALA	2.8
1	B	333	TYR	2.8
1	A	399	ALA	2.8
1	A	440	SER	2.8
1	B	389	ALA	2.8
1	B	22	ALA	2.8
1	B	507	LEU	2.8
1	A	212	ASP	2.8
1	B	337	TYR	2.8
1	A	483	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	466	SER	2.7
1	A	442	ALA	2.7
1	B	461	THR	2.7
1	B	690	ALA	2.7
1	A	135	PRO	2.7
1	B	161	ASP	2.7
1	B	678	VAL	2.7
1	A	296	LEU	2.6
1	B	136	LEU	2.6
1	A	513	LEU	2.6
1	A	539	CYS	2.6
1	B	141	ALA	2.6
1	B	328	ALA	2.6
1	B	335	ALA	2.6
1	A	505	ALA	2.6
1	A	289	ILE	2.5
1	A	468	ASN	2.5
1	A	438	GLY	2.5
1	A	541	ILE	2.5
1	B	441	ALA	2.5
1	A	459	PHE	2.5
1	B	539	CYS	2.5
1	A	474	ALA	2.4
1	A	538	SER	2.4
1	B	158	ASN	2.4
1	A	464	ASP	2.4
1	B	463	SER	2.4
1	A	473	ALA	2.4
1	A	540	LEU	2.4
1	B	439	MET	2.4
1	A	141	ALA	2.3
1	B	464	ASP	2.3
1	B	482	PHE	2.3
1	A	40	GLN	2.3
1	A	287	ASP	2.3
1	A	456	GLY	2.3
1	A	461	THR	2.3
1	B	443	ILE	2.2
1	B	157	PHE	2.2
1	B	510	ILE	2.2
1	A	436	GLU	2.2
1	B	599	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	300	TRP	2.2
1	B	438	GLY	2.2
1	B	458	THR	2.2
1	A	602	GLY	2.2
1	B	341	ALA	2.2
1	A	137	GLY	2.1
1	B	336	LYS	2.1
1	B	506	SER	2.1
1	B	138	GLN	2.1
1	B	331	ALA	2.1
1	A	481	ILE	2.1
1	A	291	LYS	2.1
1	B	388	ALA	2.1
1	B	165	VAL	2.1
1	B	503	HIS	2.0
1	A	433	GLY	2.0
1	A	480	SER	2.0
1	B	344	PHE	2.0
1	A	457	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	OCS	A	177	9/10	0.98	0.13	-	25,27,30,32	0
1	OCS	B	177	9/10	0.96	0.12	-	26,29,38,42	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	CL	B	692	1/1	0.97	0.25	6.24	52,52,52,52	0
2	SO4	B	691	5/5	0.97	0.09	-1.42	51,53,57,65	0
2	SO4	A	691	5/5	0.98	0.11	-	39,40,42,47	0

6.5 Other polymers

There are no such residues in this entry.