



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:41 PM GMT

PDB ID : 3UNP
Title : Structure of human SUN2 SUN domain
Authors : Zhou, Z.C.; Greene, M.I.
Deposited on : 2011-11-16
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	203	<div> <div>8%</div> <div>46%</div> <div>32%</div> <div>13%</div> <div>• 5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACE	A	1	-	-	-	X

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1541 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called SUN domain-containing protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	193	1479	941	259	275	4	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	717	HIS	-	EXPRESSION TAG	UNP Q9UH99
A	718	HIS	-	EXPRESSION TAG	UNP Q9UH99
A	719	HIS	-	EXPRESSION TAG	UNP Q9UH99
A	720	HIS	-	EXPRESSION TAG	UNP Q9UH99
A	721	HIS	-	EXPRESSION TAG	UNP Q9UH99

- Molecule 2 is ACETYL GROUP (three-letter code: ACE) (formula: C₂H₄O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	3	2	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	59	Total	O	0	0
			59	59		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SUN domain-containing protein 2



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	78.92Å 78.92Å 197.92Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.47 – 2.39 34.25 – 2.39	Depositor EDS
% Data completeness (in resolution range)	96.9 (39.47-2.39) 96.9 (34.25-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.94 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.218 , 0.266 0.211 , 0.259	Depositor DCC
R_{free} test set	967 reflections (11.48%)	DCC
Wilson B-factor (Å ²)	44.0	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 69.0	EDS
Estimated twinning fraction	0.016 for $-1/3^*h+1/3^*k+1/3^*l, -k, 8/3^*h+4/3^*k+1/3^*l$ 0.021 for $-2/3^*h-1/3^*k-1/3^*l, -1/3^*h-2/3^*k+1/3^*l, -4/3^*h+4/3^*k+1/3^*l$ 0.023 for $-h, 1/3^*h-1/3^*k-1/3^*l, -4/3^*h-8/3^*k+1/3^*l$	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 9393 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1541	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.35	9/1521 (0.6%)	1.28	16/2077 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	700	PRO	N-CD	9.01	1.60	1.47
1	A	700	PRO	CA-C	-8.33	1.36	1.52
1	A	571	ALA	C-O	6.58	1.35	1.23
1	A	554	GLY	N-CA	6.10	1.55	1.46
1	A	700	PRO	CG-CD	5.83	1.69	1.50
1	A	553	GLY	N-CA	5.39	1.54	1.46
1	A	704	CYS	CB-SG	-5.31	1.73	1.81
1	A	555	ALA	CA-C	5.18	1.66	1.52
1	A	700	PRO	N-CA	-5.14	1.38	1.47

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	700	PRO	CA-CB-CG	-11.15	82.81	104.00
1	A	700	PRO	CA-N-CD	-9.77	97.82	111.50
1	A	555	ALA	N-CA-C	8.53	134.02	111.00
1	A	597	PRO	C-N-CA	-7.00	107.60	122.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	555	ALA	N-CA-CB	-6.32	101.26	110.10
1	A	559	SER	N-CA-C	-6.00	94.80	111.00
1	A	557	VAL	N-CA-C	-5.76	95.44	111.00
1	A	594	ASP	CB-CG-OD1	5.70	123.43	118.30
1	A	583	HIS	N-CA-C	5.54	125.94	111.00
1	A	613	ARG	NE-CZ-NH2	5.51	123.05	120.30
1	A	716	HIS	N-CA-C	5.38	125.53	111.00
1	A	669	ASP	C-N-CA	-5.34	111.08	122.30
1	A	700	PRO	O-C-N	5.28	131.15	122.70
1	A	651	ASP	CB-CG-OD1	-5.18	113.64	118.30
1	A	553	GLY	CA-C-N	5.05	126.29	116.20
1	A	692	ARG	NE-CZ-NH1	5.00	122.80	120.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	554	GLY	Peptide
1	A	567	GLU	Peptide
1	A	568	THR	Peptide
1	A	652	GLU	Peptide
1	A	715	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1479	0	1384	106	2
2	A	3	0	3	0	0
3	A	59	0	0	9	0
All	All	1541	0	1387	106	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 37.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:583:HIS:O	1:A:584:SER:HB2	1.34	1.12
1:A:542:ARG:CG	1:A:542:ARG:HH11	1.63	1.10
1:A:692:ARG:HG3	1:A:692:ARG:HH11	0.94	1.08
1:A:557:VAL:HG12	1:A:558:ILE:H	1.11	1.08
1:A:542:ARG:NH1	1:A:542:ARG:HG2	1.39	1.06
1:A:570:THR:O	1:A:577:GLY:CA	2.09	1.00
1:A:567:GLU:HG2	1:A:567:GLU:O	1.69	0.92
1:A:652:GLU:HA	1:A:652:GLU:OE1	1.67	0.91
1:A:692:ARG:HG3	1:A:692:ARG:NH1	1.69	0.89
1:A:563:SER:H	1:A:604:GLN:NE2	1.71	0.88
1:A:570:THR:O	1:A:577:GLY:HA2	1.72	0.88
1:A:583:HIS:O	1:A:584:SER:CB	2.19	0.82
1:A:643:LYS:HG2	1:A:697:TRP:CE2	2.14	0.82
1:A:562:CYS:O	1:A:563:SER:CB	2.27	0.81
1:A:557:VAL:CG1	1:A:558:ILE:H	1.93	0.78
1:A:585:GLN:HE21	1:A:585:GLN:HA	1.49	0.78
1:A:557:VAL:HG12	1:A:558:ILE:N	1.92	0.78
1:A:585:GLN:NE2	1:A:585:GLN:HA	1.99	0.77
1:A:563:SER:H	1:A:604:GLN:HE22	1.33	0.76
1:A:603:PHE:HZ	1:A:693:ILE:HG12	1.51	0.76
1:A:562:CYS:O	1:A:563:SER:HB3	1.85	0.75
1:A:523:THR:HG22	1:A:526:GLN:CG	2.17	0.74
1:A:690:GLU:OE2	1:A:692:ARG:NH1	2.21	0.73
1:A:558:ILE:O	1:A:558:ILE:HG13	1.87	0.73
1:A:523:THR:HG22	1:A:526:GLN:HG3	1.72	0.72
1:A:585:GLN:HE21	1:A:585:GLN:CA	2.04	0.71
1:A:714:PRO:C	1:A:716:HIS:H	1.95	0.69
1:A:570:THR:O	1:A:577:GLY:HA3	1.93	0.69
1:A:558:ILE:HD12	1:A:560:THR:HB	1.78	0.66
1:A:604:GLN:H	1:A:604:GLN:CD	1.99	0.66
1:A:550:LEU:HD13	1:A:552:SER:HB2	1.77	0.66
1:A:692:ARG:CG	1:A:692:ARG:HH11	1.89	0.66
1:A:549:ALA:O	1:A:591:LEU:HD22	1.98	0.64
1:A:714:PRO:C	1:A:716:HIS:N	2.52	0.63
1:A:536:GLN:O	1:A:540:GLU:HG3	1.99	0.62
1:A:656:GLN:O	1:A:657:GLU:O	2.18	0.61
1:A:619:ARG:NH1	3:A:34:HOH:O	2.17	0.61
1:A:542:ARG:NH1	1:A:542:ARG:CG	2.31	0.60
1:A:558:ILE:CD1	1:A:560:THR:HB	2.32	0.59
1:A:652:GLU:C	1:A:654:LEU:H	2.06	0.59
1:A:668:GLN:HG2	1:A:697:TRP:CD2	2.37	0.58
1:A:714:PRO:O	1:A:716:HIS:N	2.36	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:585:GLN:NE2	1:A:585:GLN:CA	2.63	0.57
1:A:652:GLU:HB2	1:A:656:GLN:HG2	1.87	0.55
1:A:569:LYS:O	1:A:576:PHE:HA	2.07	0.55
1:A:652:GLU:C	1:A:654:LEU:N	2.62	0.53
1:A:570:THR:O	1:A:577:GLY:N	2.42	0.52
1:A:568:THR:HA	1:A:570:THR:H	1.75	0.52
1:A:603:PHE:CZ	1:A:693:ILE:HG12	2.39	0.52
1:A:663:LYS:HD2	3:A:12:HOH:O	2.10	0.52
1:A:595:VAL:HG22	1:A:706:TYR:HB3	1.91	0.51
1:A:663:LYS:CD	3:A:12:HOH:O	2.58	0.51
1:A:652:GLU:HB2	1:A:656:GLN:CG	2.41	0.51
1:A:542:ARG:HD3	3:A:57:HOH:O	2.10	0.50
1:A:716:HIS:O	3:A:46:HOH:O	2.18	0.50
1:A:547:ASP:OD2	3:A:5:HOH:O	2.19	0.50
1:A:655:GLN:OE1	1:A:655:GLN:C	2.50	0.50
1:A:597:PRO:O	1:A:598:GLY:C	2.50	0.50
1:A:523:THR:CG2	1:A:526:GLN:H	2.25	0.49
1:A:557:VAL:CG1	1:A:558:ILE:N	2.62	0.49
1:A:656:GLN:O	1:A:657:GLU:C	2.51	0.49
1:A:585:GLN:HG3	1:A:600:CYS:O	2.13	0.49
1:A:582:TYR:C	1:A:582:TYR:CD2	2.86	0.49
1:A:570:THR:HA	1:A:577:GLY:HA2	1.94	0.48
1:A:643:LYS:HG2	1:A:697:TRP:CZ2	2.49	0.48
1:A:579:PRO:O	1:A:582:TYR:HB3	2.14	0.48
1:A:524:GLU:O	1:A:528:HIS:HD2	1.97	0.47
1:A:542:ARG:HG2	1:A:542:ARG:HH11	0.66	0.47
1:A:692:ARG:NH1	1:A:692:ARG:CG	2.57	0.47
1:A:643:LYS:NZ	3:A:44:HOH:O	2.47	0.47
1:A:643:LYS:HG2	1:A:697:TRP:NE1	2.30	0.47
1:A:585:GLN:HE21	1:A:585:GLN:N	2.13	0.46
1:A:643:LYS:HB3	1:A:695:THR:OG1	2.16	0.46
1:A:626:GLU:HA	1:A:672:PRO:O	2.15	0.46
1:A:595:VAL:O	1:A:595:VAL:CG1	2.64	0.45
1:A:595:VAL:O	1:A:595:VAL:HG13	2.16	0.45
1:A:603:PHE:CE2	1:A:703:THR:HG23	2.51	0.45
1:A:643:LYS:HB3	1:A:644:ASP:H	1.60	0.44
1:A:566:TYR:O	1:A:567:GLU:HB3	2.17	0.44
1:A:619:ARG:HG3	1:A:683:MET:HA	1.99	0.44
1:A:527:VAL:O	1:A:531:VAL:HG23	2.18	0.44
1:A:565:THR:OG1	1:A:567:GLU:OE1	2.28	0.43
1:A:523:THR:HG23	1:A:526:GLN:H	1.83	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:558:ILE:O	1:A:558:ILE:CG1	2.61	0.43
1:A:618:ILE:HD12	1:A:712:GLY:HA3	2.00	0.43
1:A:570:THR:C	1:A:577:GLY:HA2	2.36	0.43
1:A:643:LYS:HE3	1:A:697:TRP:CZ3	2.54	0.42
1:A:643:LYS:HE3	1:A:697:TRP:CH2	2.55	0.42
1:A:523:THR:CG2	1:A:526:GLN:HG3	2.44	0.42
1:A:578:ILE:HD12	1:A:578:ILE:H	1.84	0.42
1:A:559:SER:HB3	3:A:15:HOH:O	2.18	0.42
1:A:558:ILE:HD12	1:A:560:THR:H	1.85	0.41
1:A:648:PHE:O	1:A:689:VAL:HA	2.19	0.41
1:A:667:ASP:O	1:A:669:ASP:O	2.37	0.41
1:A:559:SER:CB	3:A:15:HOH:O	2.68	0.41
1:A:582:TYR:CG	1:A:583:HIS:N	2.87	0.41
1:A:523:THR:HG22	1:A:526:GLN:CD	2.40	0.41
1:A:696:ASN:OD1	1:A:696:ASN:C	2.59	0.41
1:A:702:TYR:HE1	1:A:704:CYS:HG	1.66	0.41
1:A:570:THR:CA	1:A:577:GLY:HA2	2.51	0.41
1:A:625:LEU:HA	1:A:625:LEU:HD23	1.99	0.40
1:A:635:ASN:O	1:A:636:SER:HB2	2.21	0.40
1:A:669:ASP:C	1:A:670:GLY:O	2.59	0.40
1:A:656:GLN:HG3	1:A:656:GLN:H	1.59	0.40
1:A:563:SER:N	1:A:604:GLN:NE2	2.52	0.40
1:A:646:ALA:O	1:A:647:ILE:HD13	2.22	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:583:HIS:NE2	1:A:583:HIS:NE2[6_766]	1.78	0.42
1:A:583:HIS:CD2	1:A:583:HIS:NE2[6_766]	2.04	0.16

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	189/203 (93%)	160 (85%)	18 (10%)	11 (6%)	2 1

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	558	ILE
1	A	563	SER
1	A	584	SER
1	A	653	ASP
1	A	657	GLU
1	A	700	PRO
1	A	552	SER
1	A	567	GLU
1	A	715	ALA
1	A	557	VAL
1	A	578	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	151/174 (87%)	115 (76%)	36 (24%)	1 1

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	525	GLU
1	A	533	GLN
1	A	537	ARG
1	A	542	ARG
1	A	550	LEU
1	A	556	SER
1	A	558	ILE
1	A	559	SER
1	A	562	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	564	GLU
1	A	567	GLU
1	A	570	THR
1	A	584	SER
1	A	585	GLN
1	A	594	ASP
1	A	595	VAL
1	A	604	GLN
1	A	606	PRO
1	A	628	VAL
1	A	630	LYS
1	A	632	LEU
1	A	643	LYS
1	A	652	GLU
1	A	655	GLN
1	A	656	GLN
1	A	661	LEU
1	A	663	LYS
1	A	664	PHE
1	A	668	GLN
1	A	671	GLU
1	A	675	THR
1	A	692	ARG
1	A	693	ILE
1	A	700	PRO
1	A	703	THR
1	A	716	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	585	GLN
1	A	604	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACE	A	1	-	2,2,2	1.34	0	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACE	A	1	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	193/203 (95%)	0.40	17 (8%) 12 12	21, 37, 68, 74	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	520	ILE	7.9
1	A	571	ALA	5.0
1	A	577	GLY	3.6
1	A	700	PRO	3.6
1	A	653	ASP	3.3
1	A	716	HIS	3.3
1	A	522	VAL	3.1
1	A	565	THR	3.0
1	A	652	GLU	3.0
1	A	583	HIS	2.8
1	A	715	ALA	2.6
1	A	562	CYS	2.5
1	A	651	ASP	2.5
1	A	639	SER	2.2
1	A	654	LEU	2.1
1	A	521	GLY	2.1
1	A	655	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ACE	A	1	3/3	0.79	0.31	15.00	56,56,57,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.