



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 05:06 PM BST

PDB ID : 1USS  
Title : YEAST HISTONE H1 GLOBULAR DOMAIN II, HHO1P GII, SOLUTION NMR STRUCTURES  
Authors : Ali, T.; Coles, P.; Stevens, T.J.; Stott, K.; Thomas, J.O.  
Deposited on : 2003-11-30

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 82%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	88	

## 2 Ensemble composition and analysis

This entry contains 10 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:178-A:251 (74)	0.77	3

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 5, 7, 8, 9, 10
2	2, 6

### 3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 1421 atoms, of which 739 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called HISTONE H1.


Mol	Chain	Residues	Atoms						Trace
1	A	88	Total	C	H	N	O	S	0
			1421	435	739	115	129	3	

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: HISTONE H1




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: HISTONE H1




#### 4.2.2 Score per residue for model 2

- Molecule 1: HISTONE H1




### 4.2.3 Score per residue for model 3 (medoid)

- Molecule 1: HISTONE H1




### 4.2.4 Score per residue for model 4

- Molecule 1: HISTONE H1




### 4.2.5 Score per residue for model 5

- Molecule 1: HISTONE H1




### 4.2.6 Score per residue for model 6

- Molecule 1: HISTONE H1




### 4.2.7 Score per residue for model 7

- Molecule 1: HISTONE H1




### 4.2.8 Score per residue for model 8

- Molecule 1: HISTONE H1




### 4.2.9 Score per residue for model 9

- Molecule 1: HISTONE H1



### 4.2.10 Score per residue for model 10

- Molecule 1: HISTONE H1



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *DISTANCE GEOMETRY*, *BOND ANGLE GEOMETRY*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *MINIMUM ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
ARIA	structure solution	
CNS	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 6162
Number of chemical shift lists	1
Total number of shifts	1008
Number of shifts mapped to atoms	1008
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	82%

No validations of the models with respect to experimental NMR restraints is performed at this time.



## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.37±0.06	0±1/590 (0.0±0.1%)	0.39±0.02	0±0/788 (0.0±0.0%)
All	All	0.38	2/5900 (0.0%)	0.39	0/7880 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.1±0.3
All	All	0	1

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	180	TYR	CE2-CZ	-6.73	1.29	1.38	6	1
1	A	180	TYR	CE1-CZ	6.09	1.46	1.38	6	1

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	222	TYR	Sidechain	1

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	581	621	621	12±4
All	All	5810	6210	6210	115

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:214:LEU:HD13	1:A:216:THR:H	0.89	1.27	9	1
1:A:238:VAL:HB	1:A:250:ASN:HB2	0.80	1.51	1	1
1:A:238:VAL:HB	1:A:248:LYS:HB2	0.79	1.55	6	1
1:A:231:CYS:HB3	1:A:237:LEU:HD23	0.73	1.61	8	3
1:A:213:LYS:HE2	1:A:213:LYS:HA	0.66	1.67	8	1
1:A:180:TYR:OH	1:A:216:THR:HA	0.66	1.90	6	1
1:A:188:MET:HB3	1:A:189:PRO:HD3	0.64	1.70	8	1
1:A:214:LEU:HD13	1:A:216:THR:N	0.62	2.04	9	2
1:A:220:PHE:O	1:A:224:PHE:HB2	0.58	1.97	4	4
1:A:188:MET:HB2	1:A:189:PRO:HD3	0.57	1.76	9	4
1:A:199:ARG:HD3	1:A:247:ILE:HD11	0.57	1.75	8	1
1:A:224:PHE:O	1:A:228:ILE:HG13	0.56	1.99	10	6
1:A:231:CYS:O	1:A:234:ASN:HB2	0.56	2.00	4	5
1:A:202:LEU:HD13	1:A:247:ILE:HD12	0.55	1.78	5	1
1:A:212:SER:O	1:A:213:LYS:HG2	0.54	2.03	5	1
1:A:214:LEU:HB3	1:A:216:THR:HG23	0.54	1.77	7	1
1:A:184:ILE:HG23	1:A:202:LEU:HD21	0.54	1.80	7	1
1:A:232:VAL:HA	1:A:237:LEU:O	0.54	2.04	8	3
1:A:206:VAL:HA	1:A:210:PHE:CD1	0.53	2.39	8	3
1:A:180:TYR:O	1:A:184:ILE:HG12	0.53	2.03	6	1
1:A:202:LEU:O	1:A:206:VAL:HG23	0.53	2.02	4	2
1:A:199:ARG:HG3	1:A:247:ILE:HD11	0.52	1.81	3	1
1:A:214:LEU:CD1	1:A:216:THR:HB	0.52	2.35	1	1
1:A:214:LEU:HD22	1:A:215:LYS:N	0.52	2.19	1	1
1:A:206:VAL:O	1:A:210:PHE:HB2	0.51	2.05	9	1
1:A:249:LEU:HD12	1:A:251:LYS:HD2	0.51	1.83	4	1
1:A:184:ILE:HA	1:A:202:LEU:HD11	0.50	1.83	4	1
1:A:231:CYS:HB2	1:A:237:LEU:HD23	0.50	1.83	5	1
1:A:184:ILE:O	1:A:188:MET:HG2	0.50	2.07	2	2
1:A:219:ASN:O	1:A:223:LEU:HG	0.49	2.07	5	1
1:A:180:TYR:O	1:A:184:ILE:HG13	0.49	2.08	8	1

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:197:SER:O	1:A:247:ILE:HB	0.49	2.08	7	5
1:A:192:ASN:ND2	1:A:194:GLY:H	0.48	2.07	10	1
1:A:187:SER:O	1:A:191:LEU:HB2	0.48	2.08	9	2
1:A:239:GLN:HB3	1:A:244:SER:HA	0.47	1.85	2	1
1:A:183:MET:HB3	1:A:206:VAL:CG2	0.47	2.39	2	2
1:A:232:VAL:HG13	1:A:238:VAL:HA	0.47	1.85	8	1
1:A:229:LYS:O	1:A:233:GLU:HG3	0.47	2.09	7	2
1:A:232:VAL:HG21	1:A:239:GLN:CG	0.47	2.40	5	1
1:A:197:SER:OG	1:A:201:VAL:HB	0.47	2.10	8	1
1:A:181:LYS:H	1:A:181:LYS:HD3	0.47	1.70	5	1
1:A:196:GLY:HA3	1:A:248:LYS:HA	0.46	1.87	4	1
1:A:186:LYS:O	1:A:190:GLN:HG2	0.46	2.10	3	1
1:A:219:ASN:HB3	1:A:223:LEU:HD12	0.46	1.88	7	1
1:A:187:SER:O	1:A:191:LEU:HG	0.45	2.10	1	1
1:A:188:MET:HB2	1:A:189:PRO:CD	0.45	2.42	2	1
1:A:237:LEU:HD11	1:A:247:ILE:HG23	0.45	1.88	2	1
1:A:229:LYS:O	1:A:233:GLU:HG2	0.45	2.11	9	1
1:A:188:MET:CB	1:A:189:PRO:HD3	0.45	2.41	8	1
1:A:228:ILE:O	1:A:232:VAL:HG23	0.45	2.11	8	2
1:A:238:VAL:HG12	1:A:248:LYS:O	0.44	2.12	10	2
1:A:210:PHE:O	1:A:214:LEU:HB3	0.44	2.12	9	1
1:A:231:CYS:HB3	1:A:237:LEU:CD2	0.44	2.43	9	1
1:A:186:LYS:O	1:A:190:GLN:HG3	0.43	2.13	6	1
1:A:232:VAL:HB	1:A:239:GLN:HE21	0.43	1.74	1	1
1:A:210:PHE:O	1:A:214:LEU:HB2	0.43	2.14	7	2
1:A:212:SER:O	1:A:215:LYS:HG3	0.43	2.14	4	1
1:A:231:CYS:HA	1:A:234:ASN:HD22	0.42	1.74	9	1
1:A:231:CYS:HB3	1:A:237:LEU:HD22	0.42	1.91	10	1
1:A:183:MET:HB3	1:A:206:VAL:HG21	0.42	1.90	3	1
1:A:216:THR:HG21	1:A:220:PHE:HB2	0.42	1.90	4	1
1:A:196:GLY:HA3	1:A:248:LYS:HG2	0.42	1.90	8	1
1:A:183:MET:SD	1:A:214:LEU:HG	0.42	2.55	7	1
1:A:202:LEU:O	1:A:206:VAL:HG22	0.42	2.14	8	1
1:A:181:LYS:HG2	1:A:231:CYS:SG	0.42	2.55	4	1
1:A:202:LEU:HB3	1:A:224:PHE:HZ	0.42	1.74	8	1
1:A:232:VAL:HG21	1:A:239:GLN:HG2	0.41	1.92	7	1
1:A:188:MET:HB3	1:A:189:PRO:CD	0.41	2.45	1	2
1:A:214:LEU:O	1:A:214:LEU:HD22	0.41	2.15	9	1
1:A:228:ILE:HD13	1:A:247:ILE:HD11	0.41	1.92	10	1
1:A:229:LYS:O	1:A:232:VAL:HG12	0.41	2.16	1	1
1:A:185:LEU:C	1:A:185:LEU:HD13	0.41	2.36	5	1

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:236:GLU:HB2	1:A:250:ASN:O	0.41	2.16	2	1
1:A:181:LYS:O	1:A:185:LEU:HB2	0.41	2.15	9	1
1:A:188:MET:SD	1:A:197:SER:HB3	0.40	2.56	8	1
1:A:219:ASN:O	1:A:223:LEU:HB2	0.40	2.16	1	1
1:A:180:TYR:OH	1:A:220:PHE:HA	0.40	2.15	2	1
1:A:178:LEU:HD21	1:A:213:LYS:HB3	0.40	1.94	10	1
1:A:199:ARG:H	1:A:199:ARG:HD3	0.40	1.77	3	1
1:A:236:GLU:HB3	1:A:237:LEU:HD22	0.40	1.91	8	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	74/88 (84%)	66±2 (89±3%)	8±2 (10±3%)	1±1 (1±1%)	21	68
All	All	740/880 (84%)	656 (89%)	75 (10%)	9 (1%)	21	68

All 7 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	192	ASN	2
1	A	216	THR	2
1	A	244	SER	1
1	A	194	GLY	1
1	A	193	ASP	1
1	A	217	SER	1
1	A	218	SER	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	68/81 (84%)	63±1 (93±2%)	5±1 (7±2%)	23 68
All	All	680/810 (84%)	632 (93%)	48 (7%)	23 68

All 27 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	249	LEU	6
1	A	251	LYS	4
1	A	195	LYS	3
1	A	246	ILE	3
1	A	214	LEU	3
1	A	188	MET	3
1	A	191	LEU	2
1	A	199	ARG	2
1	A	230	LYS	2
1	A	181	LYS	2
1	A	202	LEU	2
1	A	186	LYS	1
1	A	204	LYS	1
1	A	190	GLN	1
1	A	236	GLU	1
1	A	206	VAL	1
1	A	219	ASN	1
1	A	225	ASN	1
1	A	237	LEU	1
1	A	238	VAL	1
1	A	213	LYS	1
1	A	203	LYS	1
1	A	193	ASP	1
1	A	201	VAL	1
1	A	179	THR	1
1	A	216	THR	1
1	A	229	LYS	1

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 82% for the well-defined parts and 82% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 6162

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1008
Number of shifts mapped to atoms	1008
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	88	$-0.23 \pm 0.11$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	83	$0.23 \pm 0.09$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	0	—	—
$^{15}\text{N}$	82	$0.73 \pm 0.44$	None needed (imprecise)

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 82%, i.e. 767 atoms were assigned a chemical shift out of a possible 936. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	287/364 (79%)	143/145 (99%)	74/148 (50%)	70/71 (99%)
Sidechain	478/521 (92%)	299/309 (97%)	172/189 (91%)	7/23 (30%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	2/51 (4%)	2/27 (7%)	0/24 (0%)	0/0 (—%)
Overall	767/936 (82%)	444/481 (92%)	246/361 (68%)	77/94 (82%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 82%, i.e. 901 atoms were assigned a chemical shift out of a possible 1101. 1 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	338/432 (78%)	168/172 (98%)	88/176 (50%)	82/84 (98%)
Sidechain	561/618 (91%)	349/368 (95%)	205/223 (92%)	7/27 (26%)
Aromatic	2/51 (4%)	2/27 (7%)	0/24 (0%)	0/0 (—%)
Overall	901/1101 (82%)	519/567 (92%)	293/423 (69%)	89/111 (80%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

