



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 09:03 PM GMT

PDB ID : 4US8
Title : Aldehyde Oxidoreductase from Desulfovibrio gigas (MOP), soaked with benzaldehyde
Authors : Correia, H.D.; Romao, M.J.; Santos-Silva, T.
Deposited on : 2014-07-03
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
9	HBX	A	922[A]	-	-	-	X
9	HBX	A	923[B]	-	-	-	X
9	HBX	A	924[C]	-	-	X	X
9	HBX	A	925[D]	-	-	-	X

2 Entry composition [\(i\)](#)


There are 10 unique types of molecules in this entry. The entry contains 8459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALDEHYDE OXIDOREDUCTASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	907	Total	C 7080	N 4502	O 1195	S 1336	47	0	50	0

- Molecule 2 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	Fe 4	S 2	2	0	0
2	A	1	Total	Fe 4	S 2	2	0	0

- Molecule 3 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 7 4 3	0	0

- Molecule 4 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C_3H_8O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 3 1	0	0

- Molecule 5 is BICARBONATE ION (three-letter code: BCT) (formula: CHO_3).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 1 3	0	0


- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	4	Total Mg 4 4	0	0

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	2	Total Cl 2 2	0	0

- Molecule 8 is (MOLYBDOPTERIN-CYTOSINE DINUCLEOTIDE-S,S)-DIOXO-AQUA-MOLYBDENUM(V) (three-letter code: PCD) (formula: C₁₉H₂₆MoN₈O₁₆P₂S₂).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
			Total	C	Mo	N	O	P	S		
8	A	1	48	19	1	8	16	2	2	0	0

- Molecule 9 is BENZALDEHYDE (three-letter code: HBX) (formula: C₇H₆O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
9	A	1	8	7	1	0	1
9	A	1	8	7	1	0	1
9	A	1	8	7	1	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total C O 8 7 1	0	1


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	1270	Total O 1270 1270	0	0

3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ALDEHYDE OXIDOREDUCTASE



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	913	IPA	4	0
8	A	921	PCD	1	0
9	A	922[A]	HBX	1	0
9	A	924[C]	HBX	4	0
9	A	925[D]	HBX	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
9	HBX	A	925[D]	8/8	0.88	0.24	17.29	12,15,16,20	8
9	HBX	A	923[B]	8/8	0.92	0.20	10.07	18,20,22,22	8
9	HBX	A	922[A]	8/8	0.96	0.18	9.39	6,7,8,8	8
9	HBX	A	924[C]	8/8	0.97	0.16	8.70	13,14,14,18	8
3	PGE	A	912	7/10	0.91	0.09	1.79	31,33,44,44	0
4	IPA	A	913	4/4	0.89	0.08	0.65	24,25,27,27	0
5	BCT	A	914	4/4	0.98	0.07	-0.46	8,11,13,14	0
8	PCD	A	921	48/48	1.00	0.06	-1.00	5,7,8,13	2
2	FES	A	908	4/4	1.00	0.05	-1.18	6,7,7,7	0
2	FES	A	909	4/4	1.00	0.03	-1.78	10,10,10,10	0
6	MG	A	915	1/1	1.00	0.10	-	23,23,23,23	1
7	CL	A	920	1/1	1.00	0.03	-	15,15,15,15	0
6	MG	A	917	1/1	1.00	0.04	-	11,11,11,11	0
6	MG	A	916	1/1	0.99	0.21	-	11,11,11,11	1
7	CL	A	919	1/1	1.00	0.03	-	14,14,14,14	0
6	MG	A	918	1/1	1.00	0.07	-	19,19,19,19	1

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.