



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:04 PM GMT

PDB ID : 4USI
Title : Nitrogen regulatory protein PII from Chlamydomonas reinhardtii in complex with MgATP and 2-oxoglutarate
Authors : Chellamuthu, V.R.; Forchhammer, K.; Hartmann, M.D.
Deposited on : 2014-07-08
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1278 (1.48-1.44)
Clashscore	102246	1336 (1.48-1.44)
Ramachandran outliers	100387	1320 (1.48-1.44)
Sidechain outliers	100360	1320 (1.48-1.44)
RSRZ outliers	91569	1279 (1.48-1.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	154	<div> <div>9%</div> <div>61%</div> <div>13%</div> <div>25%</div> </div>
1	B	154	<div> <div>16%</div> <div>60%</div> <div>11%</div> <div>28%</div> </div>
1	C	154	<div> <div>11%</div> <div>62%</div> <div>8%</div> <div>29%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2984 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITROGEN REGULATORY PROTEIN PII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	115	Total	C	N	O	S	0	8	0
			911	595	155	160	1			
1	B	111	Total	C	N	O	S	0	3	0
			861	561	150	149	1			
1	C	109	Total	C	N	O	S	0	2	0
			838	544	145	148	1			

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP A8JI83
A	145	SER	-	EXPRESSION TAG	UNP A8JI83
A	146	ALA	-	EXPRESSION TAG	UNP A8JI83
A	147	TRP	-	EXPRESSION TAG	UNP A8JI83
A	148	SER	-	EXPRESSION TAG	UNP A8JI83
A	149	HIS	-	EXPRESSION TAG	UNP A8JI83
A	150	PRO	-	EXPRESSION TAG	UNP A8JI83
A	151	GLN	-	EXPRESSION TAG	UNP A8JI83
A	152	PHE	-	EXPRESSION TAG	UNP A8JI83
A	153	GLU	-	EXPRESSION TAG	UNP A8JI83
A	154	LYS	-	EXPRESSION TAG	UNP A8JI83
B	1	MET	-	EXPRESSION TAG	UNP A8JI83
B	145	SER	-	EXPRESSION TAG	UNP A8JI83
B	146	ALA	-	EXPRESSION TAG	UNP A8JI83
B	147	TRP	-	EXPRESSION TAG	UNP A8JI83
B	148	SER	-	EXPRESSION TAG	UNP A8JI83
B	149	HIS	-	EXPRESSION TAG	UNP A8JI83
B	150	PRO	-	EXPRESSION TAG	UNP A8JI83
B	151	GLN	-	EXPRESSION TAG	UNP A8JI83
B	152	PHE	-	EXPRESSION TAG	UNP A8JI83
B	153	GLU	-	EXPRESSION TAG	UNP A8JI83
B	154	LYS	-	EXPRESSION TAG	UNP A8JI83
C	1	MET	-	EXPRESSION TAG	UNP A8JI83

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	145	SER	-	EXPRESSION TAG	UNP A8JI83
C	146	ALA	-	EXPRESSION TAG	UNP A8JI83
C	147	TRP	-	EXPRESSION TAG	UNP A8JI83
C	148	SER	-	EXPRESSION TAG	UNP A8JI83
C	149	HIS	-	EXPRESSION TAG	UNP A8JI83
C	150	PRO	-	EXPRESSION TAG	UNP A8JI83
C	151	GLN	-	EXPRESSION TAG	UNP A8JI83
C	152	PHE	-	EXPRESSION TAG	UNP A8JI83
C	153	GLU	-	EXPRESSION TAG	UNP A8JI83
C	154	LYS	-	EXPRESSION TAG	UNP A8JI83

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O P 31 10 5 13 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 4 is 2-OXOGLUTARIC ACID (three-letter code: AKG) (formula: $C_5H_6O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	5	5		
4	B	1	Total	C	O	0	0
			10	5	5		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	120	Total	O	0	0
			120	120		
6	B	104	Total	O	0	0
			104	104		
6	C	61	Total	O	0	0
			61	61		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: NITROGEN REGULATORY PROTEIN PII



• Molecule 1: NITROGEN REGULATORY PROTEIN PII



• Molecule 1: NITROGEN REGULATORY PROTEIN PII



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.04Å 90.17Å 46.18Å 90.00° 96.47° 90.00°	Depositor
Resolution (Å)	37.90 – 1.45 32.78 – 1.45	Depositor EDS
% Data completeness (in resolution range)	97.8 (37.90-1.45) 97.8 (32.78-1.45)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.185 , 0.216 0.193 , 0.222	Depositor DCC
R_{free} test set	2971 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	13.3	Xtriage
Anisotropy	0.968	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 59337 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2984	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, AKG, SO4, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.18	2/949 (0.2%)	1.30	8/1284 (0.6%)
1	B	1.19	1/887 (0.1%)	1.24	6/1201 (0.5%)
1	C	1.19	1/860 (0.1%)	1.31	9/1166 (0.8%)
All	All	1.18	4/2696 (0.1%)	1.28	23/3651 (0.6%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	77	GLU	CD-OE1	-8.30	1.16	1.25
1	A	100	TYR	CE1-CZ	-5.44	1.31	1.38
1	B	20	ARG	CZ-NH1	-5.41	1.26	1.33
1	A	100	TYR	CG-CD2	-5.00	1.32	1.39

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	79	LEU	CB-CG-CD1	8.65	125.70	111.00
1	A	25	PHE	CB-CG-CD1	8.60	126.82	120.80
1	C	85	ARG	NE-CZ-NH2	-7.53	116.54	120.30
1	A	32[A]	PHE	CB-CG-CD2	-7.50	115.55	120.80
1	A	32[B]	PHE	CB-CG-CD2	-7.50	115.55	120.80
1	C	26	ARG	NE-CZ-NH1	7.33	123.97	120.30
1	A	25	PHE	CB-CG-CD2	-7.25	115.72	120.80
1	B	85	ARG	NE-CZ-NH2	-7.17	116.71	120.30
1	A	29	ARG	NE-CZ-NH1	6.96	123.78	120.30
1	C	29	ARG	NE-CZ-NH2	6.94	123.77	120.30
1	C	121	ARG	NE-CZ-NH2	-6.85	116.87	120.30
1	C	119	ARG	NE-CZ-NH2	-6.08	117.26	120.30
1	B	26	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	B	40[A]	TYR	CB-CA-C	-5.83	98.74	110.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	40[B]	TYR	CB-CA-C	-5.83	98.74	110.40
1	A	121	ARG	NE-CZ-NH2	-5.67	117.46	120.30
1	C	32	PHE	CB-CG-CD1	5.44	124.61	120.80
1	A	32[A]	PHE	CB-CG-CD1	5.42	124.59	120.80
1	A	32[B]	PHE	CB-CG-CD1	5.42	124.59	120.80
1	C	106	ASP	CB-CG-OD1	5.25	123.03	118.30
1	C	85	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	B	80	ASP	CB-CG-OD1	-5.07	113.74	118.30
1	C	26	ARG	NE-CZ-NH2	-5.01	117.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	911	0	954	10	0
1	B	861	0	897	15	0
1	C	838	0	863	4	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	31	0	12	1	0
3	B	31	0	12	2	0
4	A	10	0	4	0	0
4	B	10	0	4	0	0
5	C	5	0	0	1	0
6	A	120	0	0	0	1
6	B	104	0	0	6	0
6	C	61	0	0	3	1
All	All	2984	0	2746	27	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:69:GLY:H	1:B:72:ASN:HD22	1.30	0.77
1:A:69:GLY:H	1:A:72:ASN:HD22	1.35	0.73
1:A:78[A]:LYS:HD3	1:A:80[A]:ASP:OD1	1.92	0.69
1:B:76:LYS:CE	6:B:2076:HOH:O	2.42	0.66
1:B:76:LYS:HD2	6:B:2076:HOH:O	1.96	0.64
1:B:123:ALA:HA	6:B:2101:HOH:O	2.02	0.59
1:C:67:GLU:N	6:C:2039:HOH:O	2.35	0.58
1:B:76:LYS:CD	6:B:2076:HOH:O	2.51	0.57
1:B:107:GLY:HA2	3:B:1152:ATP:H5'1	1.85	0.57
1:B:17:LYS:NZ	6:B:2014:HOH:O	2.35	0.52
1:B:8:CYS:HB2	1:B:89:ASP:OD1	2.10	0.51
1:B:40[A]:TYR:CD2	1:B:94:LEU:HD11	2.47	0.50
1:B:121:ARG:NH2	5:C:1152:SO4:O1	2.40	0.50
1:A:51:LYS:HG3	6:C:2033:HOH:O	2.14	0.48
1:B:107:GLY:N	3:B:1152:ATP:O2A	2.42	0.47
1:B:118:VAL:HB	1:C:109:ILE:HB	1.96	0.46
1:A:107:GLY:HA2	3:A:1152:ATP:H5'1	1.97	0.46
1:A:38[A]:SER:HG	1:B:68:PHE:N	2.15	0.45
1:A:122:THR:O	1:A:123:ALA:HB3	2.18	0.44
1:C:73:LEU:N	1:C:73:LEU:HD22	2.33	0.43
1:B:76:LYS:NZ	6:B:2076:HOH:O	2.40	0.43
1:A:78[A]:LYS:HG2	1:A:79:LEU:N	2.33	0.43
1:A:30:LEU:HG	1:A:34:ILE:HD12	2.01	0.43
1:A:70:PRO:HD3	6:C:2023:HOH:O	2.20	0.41
1:A:118:VAL:HG22	1:A:125:THR:HG22	2.01	0.41
1:B:40[A]:TYR:CE2	1:B:94:LEU:HD11	2.55	0.41
1:C:73:LEU:N	1:C:73:LEU:CD2	2.84	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2012:HOH:O	6:C:2046:HOH:O[1_655]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	119/154 (77%)	118 (99%)	1 (1%)	0	100	100
1	B	110/154 (71%)	110 (100%)	0	0	100	100
1	C	107/154 (70%)	106 (99%)	1 (1%)	0	100	100
All	All	336/462 (73%)	334 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	99/127 (78%)	95 (96%)	4 (4%)	38	6
1	B	92/127 (72%)	92 (100%)	0	100	100
1	C	89/127 (70%)	87 (98%)	2 (2%)	60	22
All	All	280/381 (74%)	274 (98%)	6 (2%)	72	23

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6[A]	ILE
1	A	6[B]	ILE
1	A	32[A]	PHE
1	A	32[B]	PHE
1	C	71	SER
1	C	72	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	72	ASN
1	C	72	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ATP	A	1152	2	24,33,33	1.27	2 (8%)	31,52,52	2.64	5 (16%)
4	AKG	A	1153	2	3,9,9	1.75	1 (33%)	4,11,11	1.31	1 (25%)
3	ATP	B	1152	2	24,33,33	1.24	1 (4%)	31,52,52	1.97	8 (25%)
4	AKG	B	1153	2	3,9,9	0.96	0	4,11,11	1.86	3 (75%)
5	SO4	C	1152	-	4,4,4	0.89	0	6,6,6	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	1152	2	-	0/18/38/38	0/3/3/3
4	AKG	A	1153	2	-	0/3/9/9	0/0/0/0
3	ATP	B	1152	2	-	0/18/38/38	0/3/3/3
4	AKG	B	1153	2	-	0/3/9/9	0/0/0/0
5	SO4	C	1152	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1152	ATP	PG-O3G	-2.42	1.46	1.54
4	A	1153	AKG	O5-C2	2.50	1.26	1.22
3	A	1152	ATP	C5-C4	2.69	1.46	1.40
3	A	1152	ATP	C2-N3	4.02	1.39	1.32

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1152	ATP	N3-C2-N1	-11.98	119.72	128.89
3	B	1152	ATP	N3-C2-N1	-7.38	123.24	128.89
3	A	1152	ATP	C1'-N9-C4	-3.39	121.83	126.94
3	B	1152	ATP	PA-O3A-PB	-3.17	123.83	132.73
3	B	1152	ATP	O3A-PA-O5'	-3.15	94.57	102.94
3	B	1152	ATP	PB-O3B-PG	-2.40	124.61	132.67
3	A	1152	ATP	O3A-PA-O5'	-2.33	96.75	102.94
3	B	1152	ATP	C1'-N9-C4	-2.26	123.53	126.94
3	A	1152	ATP	PA-O3A-PB	-2.21	126.52	132.73
4	B	1153	AKG	C4-C3-C2	-2.11	107.64	112.98
4	B	1153	AKG	C3-C4-C5	-2.08	108.94	112.75
4	A	1153	AKG	O5-C2-C3	2.04	124.14	120.28
3	B	1152	ATP	O3G-PG-O3B	2.14	114.80	105.09
4	B	1153	AKG	O5-C2-C3	2.17	124.37	120.28
3	B	1152	ATP	O3G-PG-O1G	2.18	117.60	110.58
3	B	1152	ATP	O2A-PA-O1A	2.32	125.09	112.53
3	A	1152	ATP	C2-N1-C6	4.92	127.56	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1152	ATP	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1152	ATP	2	0
5	C	1152	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	115/154 (74%)	0.65	14 (12%)	5 5	10, 17, 45, 51	0
1	B	111/154 (72%)	0.88	25 (22%)	1 1	11, 16, 42, 70	0
1	C	109/154 (70%)	1.00	17 (15%)	3 3	11, 18, 45, 68	0
All	All	335/462 (72%)	0.84	56 (16%)	2 2	10, 17, 45, 70	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	70	PRO	8.7
1	C	127	LEU	8.2
1	C	104	ILE	7.1
1	B	6	ILE	6.7
1	A	6[A]	ILE	6.2
1	A	5	SER	6.2
1	A	59	SER	6.1
1	B	59	SER	5.7
1	C	69	GLY	5.6
1	A	43	ARG	4.9
1	B	7	GLN	4.4
1	C	102	GLY	4.4
1	C	68	PHE	4.0
1	A	125	THR	4.0
1	B	55	VAL	3.9
1	A	126	GLY	3.9
1	B	8	CYS	3.8
1	C	125	THR	3.8
1	C	100	TYR	3.7
1	C	126	GLY	3.6
1	A	7	GLN	3.6
1	B	123	ALA	3.5
1	C	8	CYS	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	7	GLN	3.3
1	B	122	THR	3.3
1	C	71	SER	3.2
1	B	58	GLY	3.2
1	B	57	GLY	3.2
1	B	124	GLU	3.2
1	B	54	GLY	3.1
1	B	73	LEU	3.1
1	B	9	ASP	3.1
1	C	103	GLU	3.0
1	C	9	ASP	2.8
1	A	40	TYR	2.8
1	A	8	CYS	2.7
1	B	71	SER	2.7
1	C	72	ASN	2.7
1	B	121	ARG	2.6
1	A	123	ALA	2.6
1	B	68	PHE	2.3
1	B	56	GLN	2.3
1	C	54	GLY	2.3
1	A	68	PHE	2.2
1	B	53	VAL	2.2
1	B	109	ILE	2.2
1	A	120	ILE	2.2
1	B	120	ILE	2.2
1	B	40[A]	TYR	2.2
1	A	21[A]	ILE	2.1
1	B	81	ILE	2.1
1	B	119	ARG	2.1
1	B	70	PRO	2.0
1	C	101	THR	2.0
1	A	71	SER	2.0
1	B	11	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	SO4	C	1152	5/5	0.95	0.26	0.70	38,41,62,65	0
4	AKG	A	1153	10/10	0.96	0.13	0.10	16,18,21,21	0
3	ATP	B	1152	31/31	0.94	0.13	-0.01	15,18,23,25	0
4	AKG	B	1153	10/10	0.86	0.15	-0.30	18,20,21,21	10
3	ATP	A	1152	31/31	0.98	0.06	-1.46	11,13,15,18	0
2	MG	A	1151	1/1	0.99	0.05	-	16,16,16,16	0
2	MG	B	1151	1/1	0.91	0.10	-	24,24,24,24	0

6.5 Other polymers

There are no such residues in this entry.