



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:45 PM GMT

PDB ID : 3UTE  
Title : Crystal structure of Aspergillus fumigatus UDP galactopyranose mutase sulfate complex  
Authors : Dhatwalia, R.; Singh, H.; Tanner, J.J.  
Deposited on : 2011-11-25  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*





The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	513	 89% 9% •
1	B	513	 89% 9% •
1	C	513	 92% 6% •
1	D	513	 88% 10% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	512	-	-	-	X
3	SO4	B	512	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 16420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-galactopyranose mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	505	Total	C	N	O	S	13	1	0
			3917	2489	667	739	22			
1	B	505	Total	C	N	O	S	6	1	0
			3916	2487	663	744	22			
1	C	504	Total	C	N	O	S	0	2	0
			3878	2466	656	734	22			
1	D	504	Total	C	N	O	S	0	1	0
			3886	2468	660	736	22			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	ALA	-	EXPRESSION TAG	UNP Q4W1X2
A	-1	ILE	-	EXPRESSION TAG	UNP Q4W1X2
A	0	ALA	-	EXPRESSION TAG	UNP Q4W1X2
A	344	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
A	345	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
A	429	THR	ALA	SEE REMARK 999	UNP Q4W1X2
B	-2	ALA	-	EXPRESSION TAG	UNP Q4W1X2
B	-1	ILE	-	EXPRESSION TAG	UNP Q4W1X2
B	0	ALA	-	EXPRESSION TAG	UNP Q4W1X2
B	344	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
B	345	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
B	429	THR	ALA	SEE REMARK 999	UNP Q4W1X2
C	-2	ALA	-	EXPRESSION TAG	UNP Q4W1X2
C	-1	ILE	-	EXPRESSION TAG	UNP Q4W1X2
C	0	ALA	-	EXPRESSION TAG	UNP Q4W1X2
C	344	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
C	345	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
C	429	THR	ALA	SEE REMARK 999	UNP Q4W1X2
D	-2	ALA	-	EXPRESSION TAG	UNP Q4W1X2
D	-1	ILE	-	EXPRESSION TAG	UNP Q4W1X2
D	0	ALA	-	EXPRESSION TAG	UNP Q4W1X2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	344	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
D	345	ALA	LYS	ENGINEERED MUTATION	UNP Q4W1X2
D	429	THR	ALA	SEE REMARK 999	UNP Q4W1X2

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	158	Total	O	0	0
			158	158		
6	B	146	Total	O	0	0
			146	146		
6	C	94	Total	O	0	0
			94	94		
6	D	118	Total	O	0	0
			118	118		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: UDP-galactopyranose mutase




#### • Molecule 1: UDP-galactopyranose mutase



#### • Molecule 1: UDP-galactopyranose mutase



#### • Molecule 1: UDP-galactopyranose mutase



L254	L260	L261	Y262	Y269	F273	R303	Y326	L346	L351	A352	D353	S378	Q384	P403	E406	H412	R413	R414	F415	R447	M463	D470	D485	R496	K506	SER	LVS	ALA	GIN
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	217.84Å 217.84Å 319.69Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.42 – 2.35 49.73 – 2.35	Depositor EDS
% Data completeness (in resolution range)	90.4 (49.42-2.35) 90.2 (49.73-2.35)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.95 (at 2.34Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.2_432)	Depositor
R, $R_{free}$	0.188 , 0.218 0.185 , 0.213	Depositor DCC
$R_{free}$ test set	8374 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.6	Xtriage
Anisotropy	0.469	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 167333 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	16420	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, FAD, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/4018	0.52	0/5472
1	B	0.40	0/4017	0.52	0/5473
1	C	0.36	0/3982	0.52	0/5433
1	D	0.37	0/3987	0.51	0/5438
All	All	0.38	0/16004	0.52	0/21816

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3917	0	3793	36	0
1	B	3916	0	3776	30	0
1	C	3878	0	3709	21	0
1	D	3886	0	3729	33	0
2	A	53	0	31	4	0
2	B	53	0	30	7	0
2	C	53	0	31	7	0
2	D	53	0	31	5	0
3	A	25	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	25	0	0	0	0
3	C	15	0	0	0	0
3	D	20	0	0	0	0
4	B	4	0	3	0	0
5	B	6	0	8	0	0
6	A	158	0	0	0	0
6	B	146	0	0	2	0
6	C	94	0	0	0	0
6	D	118	0	0	0	0
All	All	16420	0	15141	125	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 125 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:146:ILE:HG23	1:D:159:MET:HE3	1.45	0.98
1:B:46:LEU:HD23	2:B:2001:FAD:C8M	2.14	0.78
1:B:46:LEU:HD23	2:B:2001:FAD:HM82	1.66	0.77
1:A:146:ILE:HG23	1:A:159:MET:HE3	1.67	0.76
1:B:159:MET:HG2	6:B:583:HOH:O	1.94	0.68

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	504/513 (98%)	497 (99%)	7 (1%)	0	100	100
1	B	504/513 (98%)	493 (98%)	11 (2%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	504/513 (98%)	497 (99%)	7 (1%)	0	100	100
1	D	503/513 (98%)	496 (99%)	7 (1%)	0	100	100
All	All	2015/2052 (98%)	1983 (98%)	32 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	415/432 (96%)	411 (99%)	4 (1%)	82	91
1	B	415/432 (96%)	412 (99%)	3 (1%)	88	95
1	C	405/432 (94%)	399 (98%)	6 (2%)	72	85
1	D	409/432 (95%)	403 (98%)	6 (2%)	72	85
All	All	1644/1728 (95%)	1625 (99%)	19 (1%)	78	89

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	69	TYR
1	C	351	LEU
1	D	175	GLN
1	C	7	SER
1	D	351	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	289	GLN
1	D	289	GLN
1	D	384	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

23 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FAD	A	2001	-	48,58,58	1.87	17 (35%)	54,89,89	2.78	22 (40%)
3	SO4	A	511	-	4,4,4	0.14	0	6,6,6	0.14	0
3	SO4	A	512	-	4,4,4	0.11	0	6,6,6	0.18	0
3	SO4	A	513	-	4,4,4	0.13	0	6,6,6	0.20	0
3	SO4	A	514	-	4,4,4	0.18	0	6,6,6	0.20	0
3	SO4	A	515	-	4,4,4	0.14	0	6,6,6	0.18	0
2	FAD	B	2001	-	48,58,58	1.93	15 (31%)	54,89,89	2.78	19 (35%)
3	SO4	B	511	-	4,4,4	0.10	0	6,6,6	0.30	0
3	SO4	B	512	-	4,4,4	0.14	0	6,6,6	0.13	0
3	SO4	B	513	-	4,4,4	0.15	0	6,6,6	0.18	0
3	SO4	B	514	-	4,4,4	0.25	0	6,6,6	0.12	0
3	SO4	B	515	-	4,4,4	0.13	0	6,6,6	0.24	0
4	ACT	B	516	-	1,3,3	1.34	0	0,3,3	0.00	-
5	GOL	B	517	-	5,5,5	0.29	0	5,5,5	0.78	0
2	FAD	C	2001	-	48,58,58	1.89	16 (33%)	54,89,89	2.57	17 (31%)
3	SO4	C	511	-	4,4,4	0.13	0	6,6,6	0.13	0
3	SO4	C	512	-	4,4,4	0.16	0	6,6,6	0.26	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	C	513	-	4,4,4	0.19	0	6,6,6	0.15	0
2	FAD	D	2001	-	48,58,58	1.91	14 (29%)	54,89,89	2.47	16 (29%)
3	SO4	D	511	-	4,4,4	0.21	0	6,6,6	0.16	0
3	SO4	D	512	-	4,4,4	0.24	0	6,6,6	0.25	0
3	SO4	D	513	-	4,4,4	0.16	0	6,6,6	0.12	0
3	SO4	D	514	-	4,4,4	0.15	0	6,6,6	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	2001	-	-	0/30/50/50	0/6/6/6
3	SO4	A	511	-	-	0/0/0/0	0/0/0/0
3	SO4	A	512	-	-	0/0/0/0	0/0/0/0
3	SO4	A	513	-	-	0/0/0/0	0/0/0/0
3	SO4	A	514	-	-	0/0/0/0	0/0/0/0
3	SO4	A	515	-	-	0/0/0/0	0/0/0/0
2	FAD	B	2001	-	-	0/30/50/50	0/6/6/6
3	SO4	B	511	-	-	0/0/0/0	0/0/0/0
3	SO4	B	512	-	-	0/0/0/0	0/0/0/0
3	SO4	B	513	-	-	0/0/0/0	0/0/0/0
3	SO4	B	514	-	-	0/0/0/0	0/0/0/0
3	SO4	B	515	-	-	0/0/0/0	0/0/0/0
4	ACT	B	516	-	-	0/0/0/0	0/0/0/0
5	GOL	B	517	-	-	0/4/4/4	0/0/0/0
2	FAD	C	2001	-	-	0/30/50/50	0/6/6/6
3	SO4	C	511	-	-	0/0/0/0	0/0/0/0
3	SO4	C	512	-	-	0/0/0/0	0/0/0/0
3	SO4	C	513	-	-	0/0/0/0	0/0/0/0
2	FAD	D	2001	-	-	0/30/50/50	0/6/6/6
3	SO4	D	511	-	-	0/0/0/0	0/0/0/0
3	SO4	D	512	-	-	0/0/0/0	0/0/0/0
3	SO4	D	513	-	-	0/0/0/0	0/0/0/0
3	SO4	D	514	-	-	0/0/0/0	0/0/0/0

The worst 5 of 62 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2001	FAD	O4'-C4'	-3.31	1.35	1.43
2	B	2001	FAD	PA-O2A	-2.73	1.43	1.54

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	FAD	O4'-C4'	-2.70	1.37	1.43
2	C	2001	FAD	O2'-C2'	-2.58	1.37	1.43
2	D	2001	FAD	O2'-C2'	-2.37	1.38	1.43

The worst 5 of 74 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2001	FAD	N3A-C2A-N1A	-11.93	119.76	128.89
2	A	2001	FAD	N3A-C2A-N1A	-11.65	119.97	128.89
2	D	2001	FAD	N3A-C2A-N1A	-11.48	120.10	128.89
2	C	2001	FAD	N3A-C2A-N1A	-11.04	120.44	128.89
2	B	2001	FAD	C4-C4X-C10	-6.39	115.85	119.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2001	FAD	4	0
2	B	2001	FAD	7	0
2	C	2001	FAD	7	0
2	D	2001	FAD	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	505/513 (98%)	-0.33	2 (0%) 93 97	18, 31, 53, 97	3 (0%)
1	B	505/513 (98%)	-0.33	3 (0%) 90 95	16, 30, 51, 94	2 (0%)
1	C	504/513 (98%)	-0.32	3 (0%) 90 95	19, 36, 59, 82	0
1	D	504/513 (98%)	-0.32	1 (0%) 95 98	19, 35, 59, 77	0
All	All	2018/2052 (98%)	-0.33	9 (0%) 93 97	16, 32, 58, 97	5 (0%)

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	358	GLN	3.2
1	C	6	ILE	2.4
1	C	3	HIS	2.4
1	D	4	PRO	2.3
1	C	279	ASN	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	A	512	5/5	0.95	0.14	5.53	68,71,75,83	0
3	SO4	B	512	5/5	0.97	0.12	4.31	65,72,75,76	0
3	SO4	D	512	5/5	0.99	0.14	1.05	29,36,39,42	0
2	FAD	A	2001	53/53	0.97	0.13	0.98	14,28,41,42	0
2	FAD	C	2001	53/53	0.96	0.14	0.86	23,36,48,52	0
3	SO4	B	511	5/5	0.98	0.13	0.82	37,39,43,45	0
3	SO4	D	511	5/5	0.97	0.14	0.76	79,82,85,88	0
2	FAD	B	2001	53/53	0.97	0.12	0.74	19,29,43,44	0
2	FAD	D	2001	53/53	0.96	0.14	0.73	19,34,44,47	0
3	SO4	C	512	5/5	0.99	0.13	0.11	34,35,37,42	0
3	SO4	B	514	5/5	0.98	0.12	0.07	67,69,74,80	0
5	GOL	B	517	6/6	0.90	0.11	-0.11	34,48,52,53	0
3	SO4	A	511	5/5	0.99	0.11	-0.82	29,30,38,41	0
3	SO4	C	511	5/5	0.98	0.11	-0.95	54,68,72,77	0
3	SO4	D	513	5/5	0.96	0.16	-	68,70,75,84	0
3	SO4	A	515	5/5	0.93	0.14	-	87,90,94,97	0
3	SO4	A	513	5/5	0.98	0.13	-	61,64,71,72	0
3	SO4	D	514	5/5	0.98	0.20	-	57,63,70,72	0
3	SO4	A	514	5/5	0.95	0.29	-	75,90,95,97	0
3	SO4	B	513	5/5	0.93	0.14	-	88,88,92,94	0
4	ACT	B	516	4/4	0.91	0.26	-	50,51,53,55	4
3	SO4	C	513	5/5	0.98	0.18	-	69,76,78,80	0
3	SO4	B	515	5/5	0.98	0.13	-	51,57,63,64	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.