



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:05 PM GMT

PDB ID : 4UUX  
Title : Competence or damage-inducible protein CinA from *Thermus thermophilus*  
Authors : Derrick, J.; Karuppiah, V.  
Deposited on : 2014-07-31  
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	<div> <div>4%</div> <div>89%</div> <div>11%</div> </div>
1	B	394	<div> <div>6%</div> <div>90%</div> <div>8%</div> </div>

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CINA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	394	Total	C	N	O	S	0	0	0
			3023	1912	546	557	8			
1	B	394	Total	C	N	O	S	0	0	0
			3023	1912	546	557	8			

- Molecule 2 is ADENOSINE-5-DIPHOSPHORIBOSE (three-letter code: APR) (formula:  $C_{15}H_{23}N_5O_{14}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			36	15	5	14	2		
2	B	1	Total	C	N	O	P	0	0
			36	15	5	14	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	232	Total O 232 232	0	0
5	B	225	Total O 225 225	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: CINA



#### • Molecule 1: CINA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.39 Å 94.03 Å 132.38 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.93 – 1.99 28.93 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.0 (28.93-1.99) 99.0 (28.93-1.99)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.39 (at 2.00 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.176 , 0.213 0.174 , 0.211	Depositor DCC
$R_{free}$ test set	3064 reflections (5.12%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.5	Xtriage
Anisotropy	0.575	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 51.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 62877 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6587	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: APR, MG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.11	3/3082 (0.1%)	1.03	12/4182 (0.3%)
1	B	1.09	9/3082 (0.3%)	1.00	8/4182 (0.2%)
All	All	1.10	12/6164 (0.2%)	1.02	20/8364 (0.2%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	157	TRP	CD2-CE2	8.18	1.51	1.41
1	B	203	GLU	CD-OE2	7.37	1.33	1.25
1	B	203	GLU	CD-OE1	6.73	1.33	1.25
1	A	127	TRP	CD2-CE2	6.31	1.49	1.41
1	B	2	GLU	CD-OE1	6.15	1.32	1.25
1	B	138	TRP	CD2-CE2	6.05	1.48	1.41
1	B	56	GLU	CD-OE2	6.03	1.32	1.25
1	A	369	ARG	CZ-NH1	5.57	1.40	1.33
1	B	183	TRP	CD2-CE2	5.51	1.48	1.41
1	B	161	TRP	CD2-CE2	5.43	1.47	1.41
1	B	157	TRP	CD2-CE2	5.29	1.47	1.41
1	B	203	GLU	CG-CD	5.03	1.59	1.51

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	105	ARG	NE-CZ-NH2	-10.65	114.98	120.30
1	A	370	ARG	NE-CZ-NH1	-9.43	115.58	120.30
1	B	321	ARG	NE-CZ-NH1	9.11	124.86	120.30
1	A	369	ARG	NE-CZ-NH2	-8.19	116.20	120.30
1	A	270	LEU	CB-CG-CD1	-7.29	98.61	111.00
1	B	105	ARG	NE-CZ-NH1	7.17	123.89	120.30
1	A	200	ARG	NE-CZ-NH1	7.09	123.84	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	254	LYS	CD-CE-NZ	-6.55	96.63	111.70
1	A	52	ARG	NE-CZ-NH1	6.36	123.48	120.30
1	A	200	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	B	39	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	A	220	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	52	ARG	NE-CZ-NH2	-5.46	117.57	120.30
1	A	39	ARG	NE-CZ-NH1	5.45	123.03	120.30
1	B	52	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	173	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	B	256	ARG	NE-CZ-NH1	-5.27	117.67	120.30
1	B	200	ARG	NE-CZ-NH1	5.08	122.84	120.30
1	A	66	LEU	CB-CG-CD1	-5.05	102.41	111.00
1	B	303	ARG	NE-CZ-NH1	-5.05	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3023	0	3095	23	0
1	B	3023	0	3095	29	0
2	A	36	0	21	1	0
2	B	36	0	21	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	232	0	0	5	0
5	B	225	0	0	6	1
All	All	6587	0	6232	52	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:225:ARG:HH11	1:B:225:ARG:HG3	1.42	0.83
5:A:2126:HOH:O	2:B:400:APR:H5R1	1.89	0.73
1:B:301:LYS:NZ	1:B:314:THR:HG22	2.05	0.72
1:B:194:LEU:O	1:B:197:LEU:HD13	1.92	0.69
1:B:225:ARG:NH1	1:B:225:ARG:HG3	2.06	0.68
1:B:367:GLU:OE2	5:B:2206:HOH:O	2.11	0.67
1:A:82:VAL:HG21	1:A:149:LEU:HD11	1.74	0.67
1:B:110:ALA:HA	5:B:2069:HOH:O	2.02	0.59
1:A:225:ARG:NH2	5:A:2135:HOH:O	2.40	0.54
1:A:379:THR:O	1:A:383:ARG:HG2	2.09	0.53
1:B:197:LEU:HD12	1:B:197:LEU:N	2.24	0.52
1:B:101:GLU:OE2	1:B:105:ARG:NH2	2.42	0.52
1:B:203:GLU:OE1	1:B:225:ARG:NH1	2.29	0.52
1:B:197:LEU:HD12	1:B:197:LEU:H	1.75	0.52
1:B:82:VAL:HG21	1:B:149:LEU:HD11	1.91	0.51
1:B:197:LEU:HG	1:B:232:ARG:HD3	1.93	0.51
1:A:121:ARG:HA	1:A:138:TRP:CZ2	2.46	0.51
1:B:199:VAL:HG12	1:B:201:GLY:H	1.75	0.50
1:A:48:ALA:HB3	1:A:49:PRO:HD3	1.94	0.50
1:A:19:LEU:HD11	1:B:19:LEU:HD23	1.93	0.49
1:B:295:SER:HB2	1:B:301:LYS:HG3	1.94	0.49
1:B:109:ARG:O	5:B:2069:HOH:O	2.20	0.48
1:B:197:LEU:H	1:B:197:LEU:CD1	2.27	0.47
1:A:308:GLN:HB3	5:A:2172:HOH:O	2.14	0.47
1:A:298:VAL:HG23	5:A:2167:HOH:O	2.13	0.47
1:B:225:ARG:CG	1:B:225:ARG:HH11	2.20	0.47
1:B:287:ARG:NH2	5:B:2136:HOH:O	2.43	0.46
1:B:201:GLY:HA3	1:B:203:GLU:OE2	2.16	0.46
1:A:301:LYS:HE3	5:A:2157:HOH:O	2.16	0.45
1:B:138:TRP:CE3	1:B:149:LEU:HD22	2.51	0.45
1:A:222:ARG:NH1	1:A:224:ASP:OD2	2.50	0.45
1:A:224:ASP:OD1	1:A:225:ARG:N	2.49	0.45
1:B:47:VAL:HG22	1:B:77:VAL:HB	1.98	0.45
1:A:187:GLU:O	1:A:191:VAL:HG23	2.16	0.45
1:A:222:ARG:NH2	1:A:225:ARG:HG3	2.32	0.44
1:A:193:ARG:HD2	1:A:236:LYS:O	2.17	0.44
1:B:154:PRO:HA	1:B:157:TRP:CD1	2.52	0.43
1:A:267:MET:HG3	1:A:293:VAL:HG23	2.00	0.43
1:A:268:GLU:CD	1:A:275:LEU:HB3	2.39	0.43
1:A:206:VAL:HG22	1:A:219:VAL:HG22	2.00	0.43
1:A:257:MET:HG3	1:A:288:PHE:CD1	2.53	0.43
1:B:220:ARG:NH1	5:B:2110:HOH:O	2.27	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:267:MET:O	1:A:338:ALA:HA	2.19	0.42
1:A:118:GLN:NE2	2:A:400:APR:O1D	2.48	0.42
1:A:270:LEU:HD21	1:A:339:THR:HB	2.03	0.41
1:B:109:ARG:NH1	5:B:2070:HOH:O	2.50	0.41
1:B:358:VAL:HG23	1:B:391:LEU:HD11	2.02	0.41
1:B:121:ARG:HA	1:B:138:TRP:CZ2	2.56	0.41
1:B:289:TYR:CE2	1:B:291:GLY:HA2	2.56	0.41
1:B:154:PRO:HA	1:B:157:TRP:NE1	2.35	0.40
1:A:270:LEU:HD22	1:A:340:THR:C	2.42	0.40
1:A:173:ARG:O	1:A:173:ARG:HG3	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:2026:HOH:O	5:B:2162:HOH:O[4_445]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/394 (100%)	384 (98%)	8 (2%)	0	100	100
1	B	392/394 (100%)	380 (97%)	12 (3%)	0	100	100
All	All	784/788 (100%)	764 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/307 (100%)	305 (99%)	2 (1%)	88	91
1	B	307/307 (100%)	301 (98%)	6 (2%)	63	65
All	All	614/614 (100%)	606 (99%)	8 (1%)	76	79

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	GLU
1	A	188	SER
1	B	197	LEU
1	B	200	ARG
1	B	203	GLU
1	B	232	ARG
1	B	239	LYS
1	B	301	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	APR	A	400	3	32,39,39	1.15	3 (9%)	39,60,60	2.28	9 (23%)
4	SO4	A	402	-	4,4,4	1.07	0	6,6,6	0.40	0
2	APR	B	400	3	32,39,39	1.01	2 (6%)	39,60,60	2.26	10 (25%)
4	SO4	B	402	-	4,4,4	1.08	0	6,6,6	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	APR	A	400	3	-	0/18/54/54	0/4/4/4
4	SO4	A	402	-	-	0/0/0/0	0/0/0/0
2	APR	B	400	3	-	0/18/54/54	0/4/4/4
4	SO4	B	402	-	-	0/0/0/0	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	400	APR	C3D-C4D	-2.15	1.47	1.53
2	A	400	APR	C2-N1	2.28	1.38	1.33
2	A	400	APR	C5-C4	2.53	1.46	1.40
2	B	400	APR	C5-C4	2.72	1.46	1.40
2	A	400	APR	O4'-C1'	4.15	1.46	1.41

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	400	APR	N3-C2-N1	-9.48	121.64	128.89
2	A	400	APR	N3-C2-N1	-7.40	123.23	128.89
2	A	400	APR	C2'-C1'-N9	-6.75	103.98	114.29
2	B	400	APR	C1D-C2D-C3D	-4.57	96.40	102.45
2	B	400	APR	C2'-C1'-N9	-4.27	107.77	114.29
2	A	400	APR	C1D-C2D-C3D	-4.20	96.90	102.45

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	400	APR	C1'-N9-C4	-4.03	120.86	126.94
2	A	400	APR	PB-O3A-PA	-3.86	121.88	132.73
2	B	400	APR	C1'-N9-C4	-3.34	121.90	126.94
2	B	400	APR	O3D-C3D-C4D	-3.06	101.86	111.05
2	B	400	APR	O2D-C2D-C1D	-2.94	103.63	111.62
2	B	400	APR	PB-O3A-PA	-2.86	124.69	132.73
2	A	400	APR	C4-C5-N7	-2.59	107.10	109.48
2	B	400	APR	C5D-C4D-C3D	-2.23	106.37	115.21
2	B	400	APR	C4-C5-N7	-2.07	107.57	109.48
2	A	400	APR	C5D-C4D-C3D	-2.03	107.15	115.21
2	A	400	APR	O4'-C1'-N9	2.43	113.19	108.10
2	B	400	APR	O2B-PB-O3A	2.64	117.09	105.09
2	A	400	APR	O1D-C1D-O4D	3.87	116.30	111.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	400	APR	1	0
2	B	400	APR	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	394/394 (100%)	0.24	14 (3%)	46 48	16, 28, 50, 65	0
1	B	394/394 (100%)	0.27	24 (6%)	25 26	16, 27, 53, 82	0
All	All	788/788 (100%)	0.26	38 (4%)	34 36	16, 28, 52, 82	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	232	ARG	4.7
1	A	19	LEU	4.3
1	A	110	ALA	4.3
1	B	43	VAL	4.0
1	A	6	ILE	3.6
1	B	199	VAL	3.6
1	B	196	PRO	3.5
1	B	202	GLU	3.1
1	A	109	ARG	3.0
1	A	108	GLY	3.0
1	A	149	LEU	2.9
1	B	7	LEU	2.9
1	A	105	ARG	2.9
1	B	41	LEU	2.9
1	B	235	LYS	2.9
1	B	92	LEU	2.8
1	B	203	GLU	2.8
1	B	9	VAL	2.8
1	A	7	LEU	2.7
1	B	150	LEU	2.7
1	B	200	ARG	2.6
1	A	65	VAL	2.6
1	A	238	LEU	2.6
1	B	275	LEU	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	104	PHE	2.5
1	A	66	LEU	2.5
1	B	195	GLY	2.4
1	A	148	VAL	2.4
1	B	238	LEU	2.4
1	B	293	VAL	2.3
1	B	294	VAL	2.3
1	B	149	LEU	2.3
1	B	148	VAL	2.2
1	B	231	GLU	2.2
1	B	228	GLU	2.2
1	A	112	PRO	2.2
1	B	6	ILE	2.2
1	B	65	VAL	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	APR	B	400	36/36	0.97	0.11	-0.16	21,29,44,71	0
2	APR	A	400	36/36	0.97	0.10	-0.29	22,30,39,44	0
4	SO4	A	402	5/5	1.00	0.07	-0.48	21,22,24,24	0
4	SO4	B	402	5/5	1.00	0.07	-0.62	22,24,27,27	0
3	MG	B	401	1/1	0.98	0.08	-1.04	31,31,31,31	0
3	MG	A	401	1/1	0.96	0.10	-	37,37,37,37	0

## 6.5 Other polymers

There are no such residues in this entry.