



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:43 PM GMT

PDB ID : 1UWT  
Title : Structure of beta-glycosidase from *Sulfolobus solfataricus* in complex with D-galactohydroxymo-1,5-lactam  
Authors : Gloster, T.M.; Roberts, S.; Ducros, V.M.-A.; Perugino, G.; Rossi, M.; Hoos, R.; Moracci, M.; Vasella, A.; Davies, G.J.  
Deposited on : 2004-02-11  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	489	<div> <div>4%</div> <div>86%</div> <div>11%</div> <div>..</div> </div>
1	B	489	<div> <div>4%</div> <div>87%</div> <div>10%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	B	1490	-	-	-	X

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 8905 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-GALACTOSIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	485	Total	C	N	O	S	0	10	0
			3983	2561	673	737	12			
1	B	484	Total	C	N	O	S	0	11	0
			3977	2555	681	729	12			

- Molecule 2 is (2E,3R,4R,5R,6S)-3,4,5-TRIHYDROXY-6-(HYDROXYMETHYL)-2-PIPERIDINONE (three-letter code: GTL) (formula:  $C_6H_{12}N_2O_5$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			13	6	2	5		
2	B	1	Total	C	N	O	0	0
			13	6	2	5		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	432	Total	O	0	0
			432	432		
4	B	483	Total	O	0	0
			483	483		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: BETA-GALACTOSIDASE



#### • Molecule 1: BETA-GALACTOSIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	168.30 Å 168.30 Å 95.67 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	141.42 – 1.95 38.51 – 1.95	Depositor EDS
% Data completeness (in resolution range)	96.9 (141.42-1.95) 97.0 (38.51-1.95)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.12 (at 1.95 Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.201 , 0.239 0.202 , 0.239	Depositor DCC
$R_{free}$ test set	5517 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.5	Xtriage
Anisotropy	0.199	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 55.4	EDS
Estimated twinning fraction	0.013 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	1 of 110240 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8905	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GTL, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.74	0/4152	0.86	18/5644 (0.3%)
1	B	0.74	0/4150	0.83	12/5641 (0.2%)
All	All	0.74	0/8302	0.85	30/11285 (0.3%)

There are no bond length outliers.

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	245	ARG	NE-CZ-NH2	-8.84	115.88	120.30
1	A	79	ARG	NE-CZ-NH2	-8.82	115.89	120.30
1	B	195	ASP	CB-CG-OD2	6.85	124.47	118.30
1	B	411	ARG	NE-CZ-NH2	-6.83	116.89	120.30
1	B	245	ARG	NE-CZ-NH1	6.30	123.45	120.30
1	A	137[A]	LEU	CA-CB-CG	6.27	129.72	115.30
1	A	137[B]	LEU	CA-CB-CG	6.27	129.72	115.30
1	A	86	ARG	NE-CZ-NH2	-6.27	117.17	120.30
1	A	39	ASP	CB-CG-OD2	5.98	123.68	118.30
1	A	370	ASP	CB-CG-OD2	5.95	123.65	118.30
1	A	308	ASP	CB-CG-OD2	5.89	123.60	118.30
1	A	119	ASP	CB-CG-OD2	5.84	123.55	118.30
1	A	286	ARG	NE-CZ-NH2	-5.71	117.45	120.30
1	A	196	ASP	CB-CG-OD2	5.65	123.39	118.30
1	B	199	ASP	CB-CG-OD2	5.60	123.34	118.30
1	A	79	ARG	NE-CZ-NH1	5.53	123.06	120.30
1	A	393	ASP	CB-CG-OD1	5.47	123.22	118.30
1	B	39	ASP	CB-CG-OD2	5.42	123.18	118.30
1	B	473	ASP	CB-CG-OD2	5.37	123.13	118.30
1	A	199	ASP	CB-CG-OD2	5.35	123.12	118.30
1	A	185	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	B	159	ASP	CB-CG-OD2	5.32	123.09	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	358	ASP	CB-CG-OD2	5.18	122.96	118.30
1	B	104	ASP	CB-CG-OD2	5.17	122.95	118.30
1	A	383	MET	CG-SD-CE	-5.09	92.05	100.20
1	B	196	ASP	CB-CG-OD2	5.08	122.87	118.30
1	B	274	ASP	CB-CG-OD2	5.08	122.88	118.30
1	A	286	ARG	NE-CZ-NH1	5.05	122.83	120.30
1	B	315	ASP	CB-CG-OD2	5.05	122.84	118.30
1	A	195	ASP	CB-CG-OD2	5.04	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3983	0	3704	33	0
1	B	3977	0	3722	46	0
2	A	13	0	12	0	0
2	B	13	0	12	0	0
3	B	4	0	3	1	0
4	A	432	0	0	3	0
4	B	483	0	0	6	0
All	All	8905	0	7453	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:489[B]:HIS:ND1	1:B:489[B]:HIS:CD2	2.07	1.23
1:B:234[A]:ARG:HB2	1:B:234[A]:ARG:HH11	1.23	1.01
1:A:489[B]:HIS:CE1	1:B:489[B]:HIS:CD2	2.50	0.99
1:B:234[A]:ARG:NH2	4:B:2290:HOH:O	2.07	0.88
1:A:13:SER:OG	1:A:79:ARG:HD3	1.77	0.82

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:TYR:CE1	1:A:185:ARG:HD2	2.16	0.81
1:A:124:LYS:HG3	4:A:2176:HOH:O	1.81	0.80
1:B:234[A]:ARG:CZ	4:B:2290:HOH:O	2.29	0.79
1:A:270:LEU:HD22	1:A:337:LEU:HD11	1.67	0.76
1:A:489[B]:HIS:CE1	1:B:489[B]:HIS:HD2	2.03	0.73
1:B:68[B]:ASP:OD1	1:B:140[B]:ARG:NH1	2.25	0.69
1:A:192:TRP:HA	1:A:253:VAL:HG11	1.75	0.69
1:B:445[A]:LYS:NZ	4:B:2438:HOH:O	2.26	0.69
1:B:245:ARG:NH2	1:B:309:ASP:OD2	2.28	0.66
1:B:234[A]:ARG:NH1	4:B:2290:HOH:O	2.26	0.65
1:B:234[A]:ARG:HB2	1:B:234[A]:ARG:NH1	2.05	0.65
1:A:329:ARG:HD3	1:A:333:GLY:O	1.98	0.64
1:A:68[A]:ASP:OD1	1:A:140:ARG:NH1	2.30	0.64
1:A:286:ARG:HD3	4:A:2291:HOH:O	1.98	0.62
1:A:489[B]:HIS:ND1	1:B:489[B]:HIS:CG	2.69	0.58
1:B:104:ASP:HA	1:B:241:GLN:HE22	1.70	0.57
1:A:270:LEU:HD22	1:A:337:LEU:CD1	2.35	0.56
1:B:241:GLN:NE2	1:B:307:ARG:HH12	2.04	0.56
1:B:110:ILE:HA	1:B:114:GLU:OE1	2.06	0.55
1:B:14:GLN:HG2	1:B:80:LEU:HD23	1.89	0.55
1:A:92:LEU:HB3	1:A:93:PRO:HD2	1.89	0.55
1:A:253:VAL:O	1:A:253:VAL:HG12	2.07	0.54
1:B:148:MET:HG3	1:B:187[A]:SER:HB2	1.88	0.54
1:B:184:ALA:HB2	1:B:245:ARG:HB3	1.93	0.51
1:B:150:HIS:O	1:B:152:PRO:HD3	2.11	0.51
1:B:365:PRO:O	1:B:408:GLN:HG3	2.10	0.51
1:B:411:ARG:HG2	3:B:1490:ACT:H1	1.95	0.49
1:B:445[B]:LYS:HG3	1:B:456:ARG:NH2	2.28	0.49
1:B:17:PHE:HB2	1:B:152:PRO:HG2	1.94	0.49
1:A:148:MET:HB2	1:A:203:THR:O	2.12	0.49
1:B:105:VAL:H	1:B:241:GLN:NE2	2.11	0.49
1:B:484:VAL:O	1:B:489[A]:HIS:HE1	1.95	0.48
1:B:108:VAL:HG23	1:B:245:ARG:HG3	1.96	0.47
1:B:241:GLN:HE21	1:B:307:ARG:HH22	1.63	0.47
1:B:245:ARG:CD	4:B:2181:HOH:O	2.63	0.47
1:A:365:PRO:O	1:A:408:GLN:HG3	2.15	0.46
1:B:105:VAL:H	1:B:241:GLN:HE22	1.63	0.46
1:B:245:ARG:HD2	4:B:2181:HOH:O	2.15	0.46
1:B:241:GLN:HE21	1:B:307:ARG:HH12	1.61	0.46
1:A:151:TRP:HB2	1:A:152:PRO:HD3	1.96	0.46
1:B:138:LYS:HD2	1:B:138:LYS:HA	1.70	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:91:PRO:HA	1:B:156:TRP:CD2	2.51	0.45
1:A:12:TRP:CD1	1:A:427:LEU:HB2	2.53	0.44
1:A:191:ALA:O	1:A:195:ASP:HB2	2.17	0.44
1:B:148:MET:HB2	1:B:203:THR:O	2.18	0.43
1:A:13:SER:OG	1:A:79:ARG:CD	2.56	0.43
1:A:253:VAL:CG1	1:A:253:VAL:O	2.66	0.43
1:A:349:VAL:HG13	1:A:353:GLY:HA2	1.99	0.43
1:A:366:GLU:HB3	4:A:2346:HOH:O	2.18	0.43
1:B:245:ARG:HA	1:B:245:ARG:HD3	1.79	0.43
1:B:87:ILE:HD12	1:B:127:LEU:HD13	2.00	0.43
1:B:21:MET:HB3	1:B:54:GLU:HA	2.01	0.43
1:A:91:PRO:HA	1:A:156:TRP:CD2	2.54	0.43
1:A:425:TRP:HA	1:A:426:SER:HA	1.78	0.43
1:A:150:HIS:O	1:A:152:PRO:HD3	2.20	0.42
1:A:17:PHE:HB2	1:A:152:PRO:HG2	2.01	0.42
1:A:137[A]:LEU:HD22	1:A:142:LEU:HD13	2.01	0.42
1:A:79:ARG:HB3	1:A:423:LEU:HD23	2.02	0.41
1:B:98:PHE:HE2	1:B:100:GLU:HG2	1.84	0.41
1:B:136:ASP:O	1:B:140[A]:ARG:HG2	2.19	0.41
1:A:39:ASP:HA	1:A:40:PRO:HD3	1.94	0.41
1:A:31:THR:HG22	1:A:85:SER:HB3	2.02	0.41
1:B:80:LEU:HD12	1:B:144:PHE:CE1	2.56	0.41
1:B:67:HIS:HB3	1:B:140[A]:ARG:HG3	2.02	0.41
1:B:191:ALA:O	1:B:195:ASP:HB2	2.21	0.41
1:B:329:ARG:HD2	1:B:333:GLY:O	2.21	0.41
1:B:79:ARG:C	1:B:79:ARG:HD2	2.41	0.40
1:B:423:LEU:N	1:B:423:LEU:HD12	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	488/489 (100%)	476 (98%)	11 (2%)	1 (0%)	52	43
1	B	488/489 (100%)	476 (98%)	11 (2%)	1 (0%)	52	43
All	All	976/978 (100%)	952 (98%)	22 (2%)	2 (0%)	52	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	151	TRP
1	B	151	TRP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	415/421 (99%)	395 (95%)	20 (5%)	31	15
1	B	417/421 (99%)	404 (97%)	13 (3%)	47	34
All	All	832/842 (99%)	799 (96%)	33 (4%)	41	23

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	12	TRP
1	A	79	ARG
1	A	103	GLN
1	A	112[A]	GLU
1	A	112[B]	GLU
1	A	114[A]	GLU
1	A	114[B]	GLU
1	A	239	ILE
1	A	275	MET
1	A	276	GLU
1	A	303	GLU
1	A	329	ARG
1	A	359	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	374	LYS
1	A	383	MET
1	A	411	ARG
1	A	414	ASN
1	A	420	ARG
1	A	462	TYR
1	B	12	TRP
1	B	79	ARG
1	B	80	LEU
1	B	124	LYS
1	B	127	LEU
1	B	138	LYS
1	B	234[A]	ARG
1	B	234[B]	ARG
1	B	241	GLN
1	B	245	ARG
1	B	305	ILE
1	B	359	PHE
1	B	386	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	264	ASN
1	A	388	ASN
1	B	113	ASN
1	B	241	GLN
1	B	264	ASN
1	B	342	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GTL	A	1490	-	12,13,13	0.99	0	9,18,18	1.95	3 (33%)
3	ACT	B	1490	-	1,3,3	1.29	0	0,3,3	0.00	-
2	GTL	B	1491	-	12,13,13	0.80	0	9,18,18	1.77	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTL	A	1490	-	-	0/2/24/24	0/1/1/1
3	ACT	B	1490	-	-	0/0/0/0	0/0/0/0
2	GTL	B	1491	-	-	0/2/24/24	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1490	GTL	C3-C4-C5	-3.08	106.44	111.23
2	B	1491	GTL	C3-C4-C5	-2.88	106.75	111.23
2	A	1490	GTL	C4-C3-C2	2.83	114.18	110.73
2	A	1490	GTL	C2-C1-N5	3.46	125.66	118.06
2	B	1491	GTL	C2-C1-N5	3.56	125.89	118.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1490	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	485/489 (99%)	0.03	22 (4%)	37 48	23, 34, 52, 67	0
1	B	484/489 (98%)	-0.03	21 (4%)	39 50	22, 33, 52, 68	0
All	All	969/978 (99%)	0.00	43 (4%)	38 49	22, 34, 52, 68	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	330	THR	4.7
1	A	332	LYS	4.0
1	B	261	ILE	3.9
1	B	98	PHE	3.8
1	A	272	ASP	3.7
1	A	333	GLY	3.6
1	B	275	MET	3.5
1	A	271	THR	3.3
1	B	95	PRO	3.2
1	A	276	GLU	3.2
1	B	93	PRO	3.2
1	A	275	MET	3.2
1	A	334	TYR	3.1
1	B	272	ASP	2.9
1	A	329	ARG	2.9
1	B	331	GLU	2.9
1	A	261	ILE	2.9
1	A	6	ASN	2.9
1	B	329	ARG	2.8
1	B	276	GLU	2.7
1	B	333	GLY	2.6
1	B	260	ILE	2.6
1	B	80	LEU	2.5
1	B	113	ASN	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	149	TYR	2.5
1	A	149	TYR	2.5
1	B	145	ILE	2.4
1	A	270	LEU	2.4
1	A	331	GLU	2.3
1	A	416	GLY	2.3
1	B	423	LEU	2.2
1	B	300	ARG	2.2
1	A	80	LEU	2.2
1	A	274	ASP	2.2
1	A	2	TYR	2.2
1	A	277	ALA	2.1
1	B	100	GLU	2.1
1	B	489[A]	HIS	2.1
1	B	305	ILE	2.1
1	A	151	TRP	2.0
1	A	376	TRP	2.0
1	A	186	PHE	2.0
1	B	299	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ACT	B	1490	4/4	0.80	0.23	4.30	60,60,61,61	0
2	GTL	A	1490	13/13	0.96	0.14	0.16	22,29,36,43	0
2	GTL	B	1491	13/13	0.96	0.13	-0.17	21,24,31,41	0

## 6.5 Other polymers

There are no such residues in this entry.