



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:50 AM GMT

PDB ID : 2UZ3  
Title : CRYSTAL STRUCTURE OF THYMIDINE KINASE WITH DTTP FROM  
U. UREALYTICUM  
Authors : Welin, M.; Kosinska, U.; Mikkelsen, N.E.; Carnrot, C.; Zhu, C.; Wang, L.;  
Eriksson, S.; Munch-Petersen, B.; Eklund, H.  
Deposited on : 2007-04-24  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	
1	B	243	
1	C	243	
1	D	243	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6651 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THYMIDINE KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	190	Total	C	N	O	S	0	0	0
			1527	984	255	281	7			
1	B	207	Total	C	N	O	S	0	0	0
			1659	1061	283	308	7			
1	C	192	Total	C	N	O	S	0	0	0
			1540	992	257	284	7			
1	D	207	Total	C	N	O	S	0	0	0
			1659	1061	283	308	7			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	16	PHE	LEU	ENGINEERED MUTATION	UNP Q9PPP5
B	16	PHE	LEU	ENGINEERED MUTATION	UNP Q9PPP5
C	16	PHE	LEU	ENGINEERED MUTATION	UNP Q9PPP5
D	16	PHE	LEU	ENGINEERED MUTATION	UNP Q9PPP5

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		
2	D	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is THYMIDINE-5'-TRIPHOSPHATE (three-letter code: TTP) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>2</sub>O<sub>14</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			29	10	2	14	3		
3	B	1	Total	C	N	O	P	0	0
			29	10	2	14	3		
3	C	1	Total	C	N	O	P	0	0
			29	10	2	14	3		
3	D	1	Total	C	N	O	P	0	0
			29	10	2	14	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	18	Total	O	0	0
			18	18		
5	B	52	Total	O	0	0
			52	52		
5	C	31	Total	O	0	0
			31	31		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	43	Total	O	0	0
			43	43		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: THYMIDINE KINASE




#### • Molecule 1: THYMIDINE KINASE



#### • Molecule 1: THYMIDINE KINASE



#### • Molecule 1: THYMIDINE KINASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.79 Å 110.75 Å 73.03 Å 90.00° 109.86° 90.00°	Depositor
Resolution (Å)	69.01 – 2.50 55.38 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (69.01-2.50) 99.7 (55.38-2.50)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.05 (at 2.51 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.220 , 0.275 0.227 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	41.4	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 39.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 33022 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6651	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, TTP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.56	0/1561	0.61	0/2104
1	B	0.55	0/1695	0.70	1/2287 (0.0%)
1	C	0.57	0/1575	0.64	1/2124 (0.0%)
1	D	0.49	0/1695	0.68	1/2287 (0.0%)
All	All	0.54	0/6526	0.66	3/8802 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	203	LEU	CA-CB-CG	5.99	129.08	115.30
1	D	192	ARG	NE-CZ-NH2	-5.45	117.58	120.30
1	B	192	ARG	NE-CZ-NH2	-5.20	117.70	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1527	0	1509	23	0
1	B	1659	0	1646	21	0
1	C	1540	0	1521	21	0
1	D	1659	0	1645	23	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	29	0	13	0	0
3	B	29	0	13	0	0
3	C	29	0	13	0	0
3	D	29	0	13	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	A	18	0	0	3	0
5	B	52	0	0	1	0
5	C	31	0	0	1	0
5	D	43	0	0	3	0
All	All	6651	0	6373	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:ILE:HG23	1:A:12:GLY:H	1.12	1.05
1:C:104:ARG:HH11	1:C:104:ARG:HG3	1.24	0.99
1:A:11:ILE:HG23	1:A:12:GLY:N	1.86	0.90
1:B:50:ILE:HD11	1:B:100:PHE:HB3	1.58	0.84
1:C:152:ILE:HD11	1:D:35:ARG:HH21	1.43	0.83
1:A:11:ILE:CG2	1:A:12:GLY:H	1.94	0.80
1:D:31:ARG:NH2	5:D:2007:HOH:O	2.20	0.72
1:B:64:THR:HG22	1:B:65:SER:H	1.55	0.71
1:A:11:ILE:CG2	1:A:12:GLY:N	2.54	0.70
1:B:50:ILE:CD1	1:B:100:PHE:HB3	2.23	0.69
1:A:129:LYS:NZ	1:A:174:TYR:O	2.23	0.69
1:B:110:ASN:OD1	1:D:192:ARG:NH2	2.27	0.67
1:B:58:ILE:HD11	1:B:96:ASP:HB3	1.81	0.63
1:C:104:ARG:NH1	1:C:104:ARG:HG3	2.01	0.62
1:A:104:ARG:HD2	1:C:204:ASN:OD1	2.00	0.61
1:A:149:LEU:HD22	1:B:31:ARG:NH2	2.16	0.61
1:C:152:ILE:CD1	1:D:35:ARG:HH21	2.16	0.57
1:C:11:ILE:HD12	1:C:11:ILE:N	2.20	0.57
1:D:102:ASP:OD1	1:D:104:ARG:HG3	2.04	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:71:VAL:HG13	1:D:76:GLU:HB2	1.88	0.56
1:A:106:CYS:SG	1:A:139:LEU:HD13	2.47	0.55
1:B:192:ARG:NH2	1:D:110:ASN:OD1	2.40	0.54
5:B:2001:HOH:O	1:D:192:ARG:HD2	2.07	0.54
1:B:138:LYS:HG3	1:D:132:PRO:HG2	1.90	0.54
1:D:17:ILE:HD13	1:D:28:GLU:HG3	1.91	0.52
1:B:50:ILE:HD13	1:B:100:PHE:CG	2.44	0.52
1:C:102:ASP:OD1	1:C:104:ARG:HG3	2.09	0.52
1:A:38:TYR:OH	1:B:61:ARG:HD2	2.09	0.52
1:C:154:ASN:HD22	1:D:39:ALA:HA	1.75	0.51
1:B:60:SER:HB2	1:B:62:THR:HG22	1.92	0.51
1:C:176:ASP:O	5:C:2024:HOH:O	2.19	0.51
1:B:192:ARG:HD2	5:D:2034:HOH:O	2.10	0.50
1:C:149:LEU:HD22	1:D:31:ARG:NH2	2.27	0.50
1:B:50:ILE:HD11	1:B:100:PHE:CB	2.35	0.49
1:A:104:ARG:HD3	5:A:2009:HOH:O	2.12	0.49
1:B:58:ILE:HD11	1:B:96:ASP:CB	2.42	0.48
1:B:14:ILE:HG12	1:B:119:VAL:HG13	1.96	0.47
1:C:183:CYS:HB3	1:C:184:GLN:H	1.46	0.47
1:D:183:CYS:SG	1:D:184:GLN:HG2	2.55	0.46
1:B:128:PHE:HE2	1:B:178:ILE:HA	1.81	0.46
1:A:166:LYS:HA	1:A:170:LYS:O	2.15	0.46
1:A:110:ASN:OD1	1:C:192:ARG:NH2	2.49	0.46
1:C:185:GLU:H	1:C:185:GLU:HG3	1.38	0.46
1:D:183:CYS:SG	1:D:184:GLN:N	2.89	0.46
1:B:50:ILE:CD1	1:B:100:PHE:CB	2.92	0.45
1:B:140:PHE:HB3	1:D:140:PHE:CG	2.52	0.45
1:C:88:ASP:N	1:C:88:ASP:OD2	2.47	0.45
1:A:96:ASP:C	1:A:97:GLU:HG2	2.36	0.45
1:A:104:ARG:HG3	1:A:104:ARG:HH11	1.81	0.44
1:B:140:PHE:CG	1:D:140:PHE:HB3	2.53	0.44
1:C:152:ILE:HG13	1:D:36:LEU:HD23	1.99	0.44
1:B:98:VAL:HG11	1:B:121:ILE:CG2	2.47	0.44
1:A:35:ARG:NH1	5:A:2005:HOH:O	2.51	0.44
1:C:32:ARG:HA	1:C:35:ARG:HD3	1.99	0.43
1:A:141:THR:O	1:C:126:LYS:NZ	2.44	0.43
1:D:43:TYR:OH	1:D:66:LEU:HD22	2.18	0.43
1:A:185:GLU:HG3	1:A:185:GLU:H	1.39	0.43
1:A:153:CYS:SG	1:A:155:GLU:HB2	2.58	0.43
1:D:136:ILE:HD12	1:D:137:ALA:N	2.33	0.43
1:A:177:ASP:N	1:A:177:ASP:OD2	2.52	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:61:ARG:N	5:D:2018:HOH:O	2.51	0.42
1:A:78:LEU:HG	1:C:210:PHE:CD1	2.54	0.42
1:C:106:CYS:SG	1:C:139:LEU:HD13	2.59	0.42
1:D:14:ILE:HB	1:D:143:ALA:HA	2.01	0.42
1:A:49:LYS:HG2	1:A:70:GLU:OE1	2.19	0.42
1:D:184:GLN:H	1:D:184:GLN:HG2	1.60	0.42
1:B:148:LYS:NZ	1:D:140:PHE:O	2.53	0.42
5:A:2013:HOH:O	1:C:192:ARG:HD2	2.20	0.41
1:A:75:PRO:HB3	1:C:206:ASN:OD1	2.21	0.41
1:A:72:GLU:HB3	1:A:76:GLU:OE1	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/243 (76%)	183 (98%)	3 (2%)	0	100	100
1	B	205/243 (84%)	197 (96%)	5 (2%)	3 (2%)	13	22
1	C	188/243 (77%)	182 (97%)	6 (3%)	0	100	100
1	D	205/243 (84%)	196 (96%)	9 (4%)	0	100	100
All	All	784/972 (81%)	758 (97%)	23 (3%)	3 (0%)	39	61

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	63	GLY
1	B	62	THR
1	B	56	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/213 (78%)	152 (92%)	14 (8%)	14	25
1	B	182/213 (85%)	163 (90%)	19 (10%)	9	16
1	C	168/213 (79%)	148 (88%)	20 (12%)	6	12
1	D	182/213 (85%)	168 (92%)	14 (8%)	16	30
All	All	698/852 (82%)	631 (90%)	67 (10%)	10	19

All (67) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ILE
1	A	42	LYS
1	A	71	VAL
1	A	78	LEU
1	A	84	ASN
1	A	98	VAL
1	A	102	ASP
1	A	119	VAL
1	A	133	PHE
1	A	138	LYS
1	A	139	LEU
1	A	177	ASP
1	A	185	GLU
1	A	197	VAL
1	B	42	LYS
1	B	50	ILE
1	B	56	ARG
1	B	57	ASN
1	B	58	ILE
1	B	64	THR
1	B	84	ASN
1	B	97	GLU
1	B	98	VAL
1	B	119	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	133	PHE
1	B	138	LYS
1	B	154	ASN
1	B	155	GLU
1	B	180	LYS
1	B	185	GLU
1	B	196	LYS
1	B	214	PHE
1	B	217	LYS
1	C	11	ILE
1	C	31	ARG
1	C	66	LEU
1	C	70	GLU
1	C	82	MET
1	C	88	ASP
1	C	98	VAL
1	C	104	ARG
1	C	119	VAL
1	C	133	PHE
1	C	139	LEU
1	C	155	GLU
1	C	168	ASP
1	C	170	LYS
1	C	183	CYS
1	C	185	GLU
1	C	192	ARG
1	C	212	LYS
1	C	215	LYS
1	C	217	LYS
1	D	11	ILE
1	D	31	ARG
1	D	42	LYS
1	D	50	ILE
1	D	52	THR
1	D	61	ARG
1	D	98	VAL
1	D	133	PHE
1	D	154	ASN
1	D	184	GLN
1	D	185	GLU
1	D	214	PHE
1	D	215	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	217	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	B	59	GLN
1	B	79	ASN
1	B	84	ASN
1	B	154	ASN
1	B	171	HIS
1	C	79	ASN
1	C	84	ASN
1	C	154	ASN
1	D	79	ASN
1	D	154	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	TTP	A	1220	-	21,30,30	0.52	0	31,47,47	2.18	5 (16%)
3	TTP	B	2220	4	21,30,30	0.64	0	31,47,47	2.07	3 (9%)
3	TTP	C	3220	-	21,30,30	0.63	0	31,47,47	2.17	6 (19%)
3	TTP	D	4220	4	21,30,30	0.59	0	31,47,47	2.05	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TTP	A	1220	-	-	0/18/34/34	0/2/2/2
3	TTP	B	2220	4	-	0/18/34/34	0/2/2/2
3	TTP	C	3220	-	-	0/18/34/34	0/2/2/2
3	TTP	D	4220	4	-	0/18/34/34	0/2/2/2

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1220	TTP	PB-O3A-PA	-6.43	114.66	132.73
3	C	3220	TTP	C5-C4-N3	-5.93	118.53	125.14
3	B	2220	TTP	PB-O3A-PA	-5.82	116.38	132.73
3	D	4220	TTP	C5-C4-N3	-5.54	118.97	125.14
3	A	1220	TTP	C5-C4-N3	-5.47	119.05	125.14
3	B	2220	TTP	C5-C4-N3	-5.12	119.44	125.14
3	D	4220	TTP	PB-O3A-PA	-4.85	119.12	132.73
3	C	3220	TTP	PB-O3A-PA	-3.87	121.86	132.73
3	A	1220	TTP	PB-O3B-PG	-3.26	121.75	132.67
3	C	3220	TTP	PB-O3B-PG	-2.28	125.04	132.67
3	D	4220	TTP	PB-O3B-PG	-2.19	125.32	132.67
3	D	4220	TTP	C2'-C3'-C4'	2.04	107.01	102.77
3	C	3220	TTP	C2'-C3'-C4'	2.36	107.66	102.77
3	C	3220	TTP	O4'-C1'-N1	2.79	112.55	107.72
3	A	1220	TTP	O4'-C1'-N1	3.77	114.25	107.72
3	A	1220	TTP	C4-N3-C2	5.94	120.38	115.25
3	B	2220	TTP	C4-N3-C2	6.94	121.24	115.25
3	D	4220	TTP	C4-N3-C2	7.21	121.48	115.25
3	C	3220	TTP	C4-N3-C2	7.34	121.59	115.25

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	190/243 (78%)	0.74	27 (14%) <b>4</b> <b>3</b>	36, 49, 69, 85	0
1	B	207/243 (85%)	0.52	8 (3%) 43 48	36, 49, 66, 85	0
1	C	192/243 (79%)	0.44	13 (6%) <b>20</b> <b>23</b>	36, 49, 66, 85	0
1	D	207/243 (85%)	0.53	8 (3%) 43 48	35, 49, 66, 85	0
All	All	796/972 (81%)	0.56	56 (7%) <b>19</b> <b>22</b>	35, 49, 66, 85	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	214	PHE	7.0
1	C	210	PHE	5.3
1	A	217	LYS	4.8
1	A	50	ILE	4.6
1	D	217	LYS	4.3
1	A	183	CYS	3.9
1	B	63	GLY	3.9
1	A	168	ASP	3.7
1	B	217	LYS	3.6
1	D	203	LEU	3.5
1	D	216	ASN	3.4
1	C	11	ILE	3.4
1	B	215	LYS	3.3
1	C	214	PHE	3.3
1	A	89	GLU	3.2
1	A	172	ALA	3.2
1	A	169	GLY	3.1
1	A	215	LYS	3.1
1	D	205	SER	3.1
1	A	199	ASN	3.0
1	B	216	ASN	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	84	ASN	2.9
1	B	203	LEU	2.9
1	A	197	VAL	2.9
1	C	65	SER	2.9
1	A	212	LYS	2.8
1	A	91	LYS	2.7
1	D	11	ILE	2.7
1	B	168	ASP	2.6
1	C	217	LYS	2.5
1	A	167	ILE	2.5
1	A	88	ASP	2.5
1	A	84	ASN	2.4
1	A	202	TYR	2.4
1	A	198	PRO	2.4
1	C	193	HIS	2.4
1	C	77	ILE	2.3
1	A	193	HIS	2.3
1	C	215	LYS	2.3
1	A	85	SER	2.2
1	D	168	ASP	2.2
1	D	212	LYS	2.2
1	A	86	PHE	2.2
1	C	72	GLU	2.2
1	C	216	ASN	2.2
1	A	170	LYS	2.2
1	D	193	HIS	2.2
1	A	186	PHE	2.2
1	A	210	PHE	2.1
1	C	84	ASN	2.1
1	C	202	TYR	2.1
1	A	203	LEU	2.1
1	A	216	ASN	2.0
1	C	79	ASN	2.0
1	B	170	LYS	2.0
1	A	87	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	TTP	A	1220	29/29	0.92	0.15	-0.67	60,62,71,72	0
3	TTP	C	3220	29/29	0.96	0.13	-0.81	44,46,55,56	0
3	TTP	D	4220	29/29	0.96	0.15	-1.01	36,42,46,48	0
3	TTP	B	2220	29/29	0.98	0.15	-1.26	36,38,42,43	0
2	ZN	A	1218	1/1	1.00	0.07	-1.72	55,55,55,55	0
2	ZN	C	3218	1/1	0.99	0.07	-1.80	47,47,47,47	0
2	ZN	D	4218	1/1	1.00	0.09	-1.83	50,50,50,50	0
2	ZN	B	2218	1/1	0.99	0.07	-2.16	47,47,47,47	0
4	MG	B	2219	1/1	0.60	0.30	-	66,66,66,66	0
4	MG	D	4219	1/1	0.58	0.31	-	59,59,59,59	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.