



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:11 PM GMT

PDB ID : 4V3K  
Title : RNF38-UbcH5B-UB complex  
Authors : Buetow, L.; Gabrielsen, M.; Anthony, N.G.; Dou, H.; Patel, A.; Aitkenhead, H.; Sibbet, G.J.; Smith, B.O.; Huang, D.T.  
Deposited on : 2014-10-20  
Resolution : 2.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1192 (2.04-2.04)
Clashscore	102246	1269 (2.04-2.04)
Ramachandran outliers	100387	1258 (2.04-2.04)
Sidechain outliers	100360	1258 (2.04-2.04)
RSRZ outliers	91569	1194 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	146	<div> <div>90%</div> <div>10% •</div> </div>
1	D	146	<div> <div>89%</div> <div>10% •</div> </div>
2	B	81	<div> <div>5%</div> <div>89%</div> <div>5% • 5%</div> </div>
2	E	81	<div> <div>86%</div> <div>9% 5%</div> </div>
3	C	79	<div> <div>5%</div> <div>84%</div> <div>11% 5%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	79	 A horizontal bar chart showing the quality of chain F. The bar is divided into four segments: a small red segment at the beginning labeled '5%', a large green segment labeled '72%', a yellow segment labeled '16%', and a small grey segment at the end labeled '11%'.

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5105 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UBIQUITIN-CONJUGATING ENZYME E2 D2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	146	Total	C	N	O	S	0	2	0
			1185	762	207	210	6			
1	D	146	Total	C	N	O	S	0	4	0
			1189	765	208	210	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	ARG	SER	ENGINEERED MUTATION	UNP P62837
A	85	LYS	CYS	ENGINEERED MUTATION	UNP P62837
D	22	ARG	SER	ENGINEERED MUTATION	UNP P62837
D	85	LYS	CYS	ENGINEERED MUTATION	UNP P62837

- Molecule 2 is a protein called POLYUBIQUITIN-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	77	Total	C	N	O	S	0	1	0
			598	377	104	116	1			
2	E	77	Total	C	N	O	S	0	0	0
			603	379	106	117	1			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	EXPRESSION TAG	UNP P0CG48
B	-3	SER	-	EXPRESSION TAG	UNP P0CG48
B	-2	GLY	-	EXPRESSION TAG	UNP P0CG48
B	-1	GLY	-	EXPRESSION TAG	UNP P0CG48
B	0	SER	-	EXPRESSION TAG	UNP P0CG48
E	-4	GLY	-	EXPRESSION TAG	UNP P0CG48
E	-3	SER	-	EXPRESSION TAG	UNP P0CG48
E	-2	GLY	-	EXPRESSION TAG	UNP P0CG48

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	-1	GLY	-	EXPRESSION TAG	UNP P0CG48
E	0	SER	-	EXPRESSION TAG	UNP P0CG48

- Molecule 3 is a protein called E3 UBIQUITIN-PROTEIN LIGASE RNF38.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	75	Total	C	N	O	S	0	2	0
			591	365	108	110	8			
3	F	70	Total	C	N	O	S	0	2	0
			541	338	94	100	9			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	387	GLY	-	EXPRESSION TAG	UNP Q9H0F5
C	388	SER	-	EXPRESSION TAG	UNP Q9H0F5
F	387	GLY	-	EXPRESSION TAG	UNP Q9H0F5
F	388	SER	-	EXPRESSION TAG	UNP Q9H0F5

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Cl	0	0
			3	3		
4	D	3	Total	Cl	0	0
			3	3		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0
5	E	1	Total C O 4 2 2	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	2	Total Zn 2 2	0	0
6	F	2	Total Zn 2 2	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	111	Total O 111 111	0	0
7	B	44	Total O 44 44	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	46	Total 46	O 46	0	0
7	D	92	Total 92	O 92	0	0
7	E	47	Total 47	O 47	0	0
7	F	28	Total 28	O 28	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UBIQUITIN-CONJUGATING ENZYME E2 D2

Chain A: 




- Molecule 1: UBIQUITIN-CONJUGATING ENZYME E2 D2

Chain D: 



- Molecule 2: POLYUBIQUITIN-C

Chain B: 




- Molecule 2: POLYUBIQUITIN-C

Chain E: 



- Molecule 3: E3 UBIQUITIN-PROTEIN LIGASE RNF38


Chain C: 



- Molecule 3: E3 UBIQUITIN-PROTEIN LIGASE RNF38

Chain F: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	139.62Å 139.62Å 70.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.91 – 2.04 37.45 – 2.04	Depositor EDS
% Data completeness (in resolution range)	99.9 (34.91-2.04) 99.9 (37.45-2.04)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.53 (at 2.05Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.181 , 0.223 0.180 , 0.222	Depositor DCC
$R_{free}$ test set	2269 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.7	Xtriage
Anisotropy	0.476	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 48.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 44939 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5105	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/1228	0.56	0/1672
1	D	0.42	0/1239	0.60	0/1689
2	B	0.38	0/607	0.57	0/818
2	E	0.38	0/609	0.57	0/819
3	C	0.39	0/610	0.57	0/830
3	F	0.36	0/559	0.54	0/761
All	All	0.40	0/4852	0.57	0/6589

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1185	0	1177	9	0
1	D	1189	0	1173	11	0
2	B	598	0	621	2	0
2	E	603	0	630	5	0
3	C	591	0	533	5	0
3	F	541	0	488	8	0
4	A	3	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	3	0	0	0	0
5	A	4	0	6	0	0
5	B	8	0	12	0	0
5	D	4	0	6	0	0
5	E	4	0	6	0	0
6	C	2	0	0	0	0
6	F	2	0	0	0	0
7	A	111	0	0	0	0
7	B	44	0	0	1	0
7	C	46	0	0	0	0
7	D	92	0	0	0	0
7	E	47	0	0	1	0
7	F	28	0	0	0	0
All	All	5105	0	4652	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:29:ASP:OD2	1:D:32[B]:HIS:HD2	1.70	0.74
1:A:18:PRO:HG2	1:A:21:CYS:HB2	1.74	0.67
1:D:115:PRO:HB2	1:D:128:LYS:HE3	1.86	0.58
3:C:396:LEU:HB2	3:C:427:ARG:HD2	1.89	0.54
1:D:43:SER:HB2	1:D:44:PRO:HD2	1.93	0.49
3:F:431:CYS:O	3:F:432:ASN:HB2	2.14	0.48
2:E:73:LEU:HD12	2:E:73:LEU:C	2.34	0.48
1:A:118:PRO:HB3	1:A:124:ALA:HB2	1.96	0.48
3:F:453:CYS:O	3:F:454:ARG:HB2	2.14	0.47
1:D:118:PRO:HB3	1:D:124:ALA:HB2	1.97	0.45
1:A:11:ASN:O	1:A:15:ARG:HB2	2.16	0.45
3:C:459[B]:GLU:OE2	1:D:90:ARG:HB3	2.17	0.45
2:B:2:GLN:HA	2:B:15:LEU:O	2.17	0.45
7:B:2018:HOH:O	3:C:462[B]:ARG:NH2	2.13	0.43
2:E:11:LYS:NZ	7:E:2007:HOH:O	2.47	0.43
1:D:117:ASP:O	2:E:76:GLY:HA2	2.19	0.43
2:E:9:THR:HG22	3:F:452:ILE:HD13	2.00	0.42
1:D:60:TYR:CD1	1:D:61:PRO:HA	2.55	0.42
1:A:107:CYS:O	1:A:110:LEU:HB2	2.20	0.42
1:A:90:ARG:H	1:A:90:ARG:HG2	1.68	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:ALA:O	1:A:20:GLN:HB2	2.20	0.42
3:F:444:LEU:HA	3:F:444:LEU:HD23	1.89	0.42
3:F:428:VAL:HG22	3:F:434[A]:GLU:HG2	2.02	0.41
1:A:90:ARG:HG3	4:A:1149:CL:CL	2.57	0.41
1:D:62:PHE:CZ	3:F:442:LYS:HD2	2.55	0.41
1:A:38:MET:SD	1:A:49:VAL:HG22	2.61	0.41
2:B:48:LYS:HD3	2:B:59:TYR:HE1	1.85	0.41
3:C:400:ARG:HD3	3:C:423:ARG:O	2.21	0.41
1:A:62:PHE:CZ	3:C:442:LYS:HD3	2.56	0.41
3:F:429:LEU:HB3	3:F:430:PRO:CD	2.51	0.40
1:D:62:PHE:CE2	3:F:442:LYS:HD2	2.57	0.40
2:E:45:PHE:HB3	2:E:50:LEU:HD21	2.04	0.40
1:D:29:ASP:OD2	1:D:32[B]:HIS:CD2	2.61	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	146/146 (100%)	143 (98%)	3 (2%)	0	100	100
1	D	148/146 (101%)	146 (99%)	2 (1%)	0	100	100
2	B	76/81 (94%)	76 (100%)	0	0	100	100
2	E	75/81 (93%)	74 (99%)	1 (1%)	0	100	100
3	C	75/79 (95%)	73 (97%)	2 (3%)	0	100	100
3	F	70/79 (89%)	67 (96%)	3 (4%)	0	100	100
All	All	590/612 (96%)	579 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/130 (101%)	129 (98%)	2 (2%)	72	70
1	D	131/130 (101%)	127 (97%)	4 (3%)	47	39
2	B	67/70 (96%)	65 (97%)	2 (3%)	48	41
2	E	68/70 (97%)	67 (98%)	1 (2%)	72	70
3	C	64/73 (88%)	62 (97%)	2 (3%)	47	39
3	F	58/73 (80%)	56 (97%)	2 (3%)	44	36
All	All	519/546 (95%)	506 (98%)	13 (2%)	57	48

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LYS
1	A	90	ARG
2	B	48	LYS
2	B	64	GLU
3	C	394	GLU
3	C	408	SER
1	D	20	GLN
1	D	90	ARG
1	D	100[A]	SER
1	D	100[B]	SER
2	E	54	ARG
3	F	412	LEU
3	F	438	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	432	ASN
3	F	447	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 10 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	EDO	A	1151	-	3,3,3	0.44	0	2,2,2	0.56	0
5	EDO	B	1077	-	3,3,3	0.37	0	2,2,2	1.01	0
5	EDO	B	1078	-	3,3,3	0.43	0	2,2,2	0.74	0
5	EDO	D	1151	-	3,3,3	0.44	0	2,2,2	0.61	0
5	EDO	E	1077	-	3,3,3	0.43	0	2,2,2	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	1151	-	-	0/1/1/1	0/0/0/0
5	EDO	B	1077	-	-	0/1/1/1	0/0/0/0
5	EDO	B	1078	-	-	0/1/1/1	0/0/0/0
5	EDO	D	1151	-	-	0/1/1/1	0/0/0/0
5	EDO	E	1077	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	146/146 (100%)	-0.34	0 <span>100</span> <span>100</span>	17, 23, 38, 46	0
1	D	146/146 (100%)	-0.31	0 <span>100</span> <span>100</span>	14, 21, 34, 48	0
2	B	77/81 (95%)	-0.45	1 (1%) <span>79</span> <span>83</span>	17, 26, 41, 53	0
2	E	77/81 (95%)	-0.41	0 <span>100</span> <span>100</span>	15, 23, 38, 54	0
3	C	75/79 (94%)	-0.14	4 (5%) <span>30</span> <span>34</span>	16, 25, 50, 61	0
3	F	70/79 (88%)	0.00	4 (5%) <span>27</span> <span>31</span>	18, 29, 58, 64	0
All	All	591/612 (96%)	-0.29	9 (1%) <span>76</span> <span>81</span>	14, 24, 43, 64	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	391	ALA	6.5
3	F	391	ALA	5.2
3	C	394	GLU	4.0
3	C	395	GLN	3.6
3	C	392	ASP	3.2
3	F	394	GLU	3.0
3	F	392	ASP	2.3
3	F	406	HIS	2.1
2	B	0	SER	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	ZN	C	1466	1/1	1.00	0.08	0.23	18,18,18,18	0
6	ZN	F	1461	1/1	0.99	0.08	0.10	21,21,21,21	0
4	CL	A	1150	1/1	0.96	0.10	-	49,49,49,49	0
4	CL	D	1150	1/1	0.94	0.23	-	42,42,42,42	0
4	CL	D	1148	1/1	0.88	0.19	-	55,55,55,55	0
4	CL	A	1148	1/1	0.92	0.07	-	50,50,50,50	0
6	ZN	F	1460	1/1	0.99	0.08	-	25,25,25,25	0
5	EDO	E	1077	4/4	0.85	0.17	-	41,43,46,50	0
4	CL	D	1149	1/1	0.98	0.10	-	47,47,47,47	0
5	EDO	A	1151	4/4	0.90	0.13	-	28,38,39,46	0
4	CL	A	1149	1/1	0.96	0.07	-	47,47,47,47	0
6	ZN	C	1465	1/1	1.00	0.07	-	23,23,23,23	0
5	EDO	D	1151	4/4	0.96	0.09	-	40,41,45,49	0
5	EDO	B	1078	4/4	0.90	0.19	-	23,29,40,46	0
5	EDO	B	1077	4/4	0.87	0.17	-	31,35,37,45	0

## 6.5 Other polymers

There are no such residues in this entry.