



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:53 AM GMT

PDB ID : 2V6Q  
Title : CRYSTAL STRUCTURE OF A BHRF-1 : BIM BH3 COMPLEX  
Authors : Kvensakul, M.; Huang, D.C.S.; Colman, P.M.  
Deposited on : 2007-07-20  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	173	
2	B	26	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1473 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BHRF1 PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	156	Total	C	N	O	S	0	0	0
			1245	788	217	231	9			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	EXPRESSION TAG	UNP P03182
A	-11	GLY	-	EXPRESSION TAG	UNP P03182
A	-10	SER	-	EXPRESSION TAG	UNP P03182
A	-9	HIS	-	EXPRESSION TAG	UNP P03182
A	-8	HIS	-	EXPRESSION TAG	UNP P03182
A	-7	HIS	-	EXPRESSION TAG	UNP P03182
A	-6	HIS	-	EXPRESSION TAG	UNP P03182
A	-5	HIS	-	EXPRESSION TAG	UNP P03182
A	-4	HIS	-	EXPRESSION TAG	UNP P03182
A	-3	SER	-	EXPRESSION TAG	UNP P03182
A	-2	GLN	-	EXPRESSION TAG	UNP P03182
A	-1	ASP	-	EXPRESSION TAG	UNP P03182
A	0	PRO	-	EXPRESSION TAG	UNP P03182

- Molecule 2 is a protein called BCL-2-LIKE PROTEIN 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	23	Total	C	N	O	S	0	0	0
			196	124	35	36	1			

- Molecule 3 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	5	Total	Br	0	0
			5	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	26	Total 26	O 26	0	0
4	B	1	Total 1	O 1	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: BHRF1 PROTEIN



#### • Molecule 2: BCL-2-LIKE PROTEIN 11



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.74Å 62.74Å 92.52Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.22 – 2.70 35.22 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.1 (35.22-2.70) 98.4 (35.22-2.70)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.70 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.206 , 0.249 0.202 , 0.242	Depositor DCC
$R_{free}$ test set	295 reflections (5.13%)	DCC
Wilson B-factor (Å <sup>2</sup> )	37.4	Xtriage
Anisotropy	0.469	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 54.2	EDS
Estimated twinning fraction	0.102 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 6044 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	1473	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.73	2/1276 (0.2%)	0.80	0/1739
2	B	0.80	0/200	0.77	0/269
All	All	0.74	2/1476 (0.1%)	0.80	0/2008

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	43	GLU	CB-CG	6.33	1.64	1.52
1	A	43	GLU	CG-CD	5.52	1.60	1.51

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	51	ASP	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1245	0	1196	41	0
2	B	196	0	181	8	0
3	A	5	0	0	3	0
4	A	26	0	0	2	0
4	B	1	0	0	0	0
All	All	1473	0	1377	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:ARG:NH1	3:A:1160:BR:BR	2.34	1.15
1:A:15:ARG:HH11	1:A:48:LEU:CD2	1.86	0.89
1:A:26:HIS:CD2	1:A:28:VAL:HG13	2.08	0.88
1:A:26:HIS:HD2	1:A:28:VAL:H	1.28	0.79
1:A:1:MET:HG3	1:A:151:SER:OG	1.82	0.79
1:A:15:ARG:HH11	1:A:48:LEU:HD21	1.48	0.75
1:A:12:LEU:HD13	1:A:54:LEU:HD11	1.77	0.67
1:A:26:HIS:HD2	1:A:28:VAL:HG13	1.63	0.64
1:A:15:ARG:NH1	1:A:48:LEU:HD21	2.11	0.64
1:A:1:MET:CG	1:A:151:SER:OG	2.48	0.61
1:A:10:LEU:HD23	1:A:38:LEU:HG	1.80	0.61
1:A:12:LEU:HD13	1:A:54:LEU:CD1	2.32	0.59
1:A:36:THR:O	1:A:38:LEU:N	2.36	0.59
1:A:101:ALA:O	1:A:105:MET:HG2	2.02	0.59
1:A:1:MET:HG3	1:A:151:SER:CB	2.33	0.58
1:A:68:THR:HB	2:B:58:ILE:HG12	1.85	0.57
1:A:155:GLU:HB2	4:A:2007:HOH:O	2.04	0.57
1:A:15:ARG:NH1	1:A:48:LEU:CD2	2.66	0.55
1:A:79:VAL:HG23	3:A:1157:BR:BR	2.63	0.54
1:A:31:LEU:HD12	1:A:35:GLU:HG2	1.91	0.53
1:A:144:ILE:HD13	1:A:150:TRP:CD1	2.44	0.52
1:A:62:SER:HB2	3:A:1158:BR:BR	2.66	0.50
2:B:53:ARG:N	2:B:54:PRO:CD	2.76	0.49
2:B:71:ALA:O	2:B:72:TYR:HB2	2.13	0.49
2:B:57:TRP:O	2:B:60:GLN:HB2	2.13	0.48
1:A:20:HIS:CD2	1:A:113:ARG:HD2	2.49	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:72:TYR:O	2:B:72:TYR:HD1	1.97	0.47
2:B:72:TYR:O	2:B:73:TYR:CB	2.63	0.47
1:A:76:THR:HG21	1:A:79:VAL:HG22	1.96	0.46
1:A:110:HIS:HD2	4:A:2020:HOH:O	1.98	0.46
1:A:122:PRO:HG2	1:A:125:VAL:HG23	1.98	0.46
1:A:26:HIS:CD2	1:A:28:VAL:H	2.20	0.46
1:A:155:GLU:OE1	1:A:156:ASP:HB2	2.15	0.46
1:A:6:ARG:NH1	1:A:36:THR:OG1	2.48	0.45
1:A:143:TRP:O	1:A:147:GLN:HG2	2.17	0.45
1:A:54:LEU:O	1:A:58:ILE:HG13	2.17	0.45
1:A:1:MET:CE	1:A:149:GLY:HA3	2.48	0.44
1:A:48:LEU:O	1:A:52:VAL:HG23	2.18	0.43
1:A:18:ARG:HD3	1:A:23:GLY:O	2.18	0.42
2:B:51:ASP:O	2:B:52:MET:HB2	2.19	0.42
1:A:76:THR:CG2	1:A:79:VAL:HG22	2.50	0.42
1:A:19:VAL:HG12	1:A:20:HIS:ND1	2.35	0.42
2:B:67:ASP:HA	2:B:70:ASN:HD22	1.85	0.42
1:A:33:ALA:O	1:A:36:THR:O	2.38	0.41
1:A:44:ASP:OD1	1:A:45:THR:N	2.53	0.41
1:A:1:MET:HG3	1:A:151:SER:HB2	2.01	0.40
1:A:86:VAL:O	1:A:90:ILE:HG13	2.21	0.40
1:A:19:VAL:CG1	1:A:19:VAL:O	2.69	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	154/173 (89%)	141 (92%)	12 (8%)	1 (1%)	30	59
2	B	21/26 (81%)	17 (81%)	2 (10%)	2 (10%)	1	1
All	All	175/199 (88%)	158 (90%)	14 (8%)	3 (2%)	11	29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	72	TYR
2	B	52	MET
1	A	21	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/153 (88%)	126 (93%)	9 (7%)	20	44
2	B	19/22 (86%)	15 (79%)	4 (21%)	1	3
All	All	154/175 (88%)	141 (92%)	13 (8%)	14	30

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LEU
1	A	25	LEU
1	A	38	LEU
1	A	45	THR
1	A	78	HIS
1	A	80	ASP
1	A	86	VAL
1	A	118	ASN
1	A	155	GLU
2	B	55	GLU
2	B	56	ILE
2	B	68	GLU
2	B	72	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	61	ASN
1	A	75	HIS
1	A	110	HIS
1	A	118	ASN
1	A	147	GLN
2	B	70	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	156/173 (90%)	-0.10	11 (7%) 19 17	12, 32, 59, 70	0
2	B	23/26 (88%)	0.27	2 (8%) 13 10	33, 49, 72, 75	0
All	All	179/199 (89%)	-0.05	13 (7%) 18 16	12, 34, 63, 75	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	4.1
1	A	118	ASN	3.5
1	A	2	ALA	3.5
2	B	51	ASP	3.2
1	A	120	SER	3.2
1	A	21	GLY	3.0
1	A	155	GLU	2.8
2	B	52	MET	2.8
1	A	22	ASN	2.6
1	A	154	ILE	2.5
1	A	156	ASP	2.5
1	A	151	SER	2.4
1	A	3	TYR	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	BR	A	1161	1/1	0.96	0.12	-1.16	89,89,89,89	0
3	BR	A	1157	1/1	0.99	0.06	-3.31	58,58,58,58	0
3	BR	A	1160	1/1	0.98	0.06	-4.23	83,83,83,83	0
3	BR	A	1158	1/1	0.99	0.07	-	46,46,46,46	0
3	BR	A	1159	1/1	0.80	0.10	-	90,90,90,90	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.