wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 05:42 PM BST

PDB ID	:	1V95
Title	:	Solution Structure of Anticodon Binding Domain from Nuclear Receptor Coactivator 5 (Human KIAA1637 Protein)
Authors	:	Zhao, C.; Kigawa, T.; Tochio, N.; Koshiba, S.; Inoue, M.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on	:	2004-01-20

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 90%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	130	27% 38% 8% 27%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:9-A:43, A:48-A:107 (95)	0.20	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 6, 9, 10, 13, 14, 15, 16, 17, 18, 19, 20
2	7, 8
Single-model clusters	5; 11; 12

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1982 atoms, of which 984 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Nuclear receptor coactivator 5.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	130	1982	608	984	185	198	7	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	CLONING ARTIFACT	UNP Q9HCD5
A	2	SER	-	CLONING ARTIFACT	UNP Q9HCD5
A	3	SER	-	CLONING ARTIFACT	UNP Q9HCD5
A	4	GLY	-	CLONING ARTIFACT	UNP Q9HCD5
A	5	SER	-	CLONING ARTIFACT	UNP Q9HCD5
A	6	SER	-	CLONING ARTIFACT	UNP Q9HCD5
A	7	GLY	-	CLONING ARTIFACT	UNP Q9HCD5
A	125	SER	-	CLONING ARTIFACT	UNP Q9HCD5
A	126	GLY	-	CLONING ARTIFACT	UNP Q9HCD5
A	127	PRO	-	CLONING ARTIFACT	UNP Q9HCD5
A	128	SER	-	CLONING ARTIFACT	UNP Q9HCD5
A	129	SER	-	CLONING ARTIFACT	UNP Q9HCD5
A	130	GLY	-	CLONING ARTIFACT	UNP Q9HCD5

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Nuclear receptor coactivator 5

Chain A: 27% 38% 8% 27%

G1 S2 S3 G4 S5 S6 G7 P8 V9 D10 C11 S12 V13 I14 V15 V16 N17 K18 Q19 T20 K21 E25 S26 V27 G28 R29 K30 V31 L34 G35 M36 V37 V38 D39 L40 L43 N44 T45 E46 V47 S48 L49 S50 Q51 A52 D55 V56 S57 R58 G59 G60 S61 P62 F63 A64 I65

V66 I67 T68 Q69 Q70 H71 Q72 I73 H74 R75 S76 I81 M82 F83 G84 T85 P86 H89 R90 N91 Q94 A95 D96 A97 M98 V99 L100 V101 A102 R103 N104 Y105 E106 R107 Y108 K109 N110 E111 C112 R113 E114 K115 E116 R117 E118 E119 I120 A121 R122 Q123 A124 S125 G126 P127 S128 S129 G130

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: Nuclear receptor coactivator 5

Chain A: 28% 34% 11% 27%

G1 S2 S3 G4 S5 S6 G7 P8 C11 S12 V13 I14 V15 V16 N17 K18 Q19 Y23 A24 E25 S26 V27 G28 R29 K30 V31 L34 G35 M36 V37 V38 D39 L40 L43 N44 T45 E46 V47 S48 L49 S50 Q51 A52 L53 V56 S57 R58 S61 P62 F63 A64 I65 V66 I67

T68 Q69 Q70 H71 Q72 I73 H74 R75 S76 I81 M82 F83 G84 T85 P86 H89 R90 N91 Q94 A95 D96 A97 M98 V99 L100 V101 A102 R103 N104 Y105 E106 R107 Y108 K109 N110 E111 C112 R113 E114 K115 E116 R117 E118 E119 I120 A121 R122 Q123 A124 S125 G126 P127 S128 S129 G130

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations, target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	1.0.7
CYANA	refinement	1.0.7

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 11255
Number of chemical shift lists	1
Total number of shifts	1428
Number of shifts mapped to atoms	1428
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	90%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	742	748	748	45±5
All	All	14840	14960	14960	892

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

5 of 188 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:16:VAL:HG12	1:A:43:LEU:HD12	0.97	1.32	11	17
1:A:52:ALA:O	1:A:56:VAL:HG23	0.90	1.66	15	17
1:A:31:VAL:HB	1:A:38:VAL:HG21	0.81	1.51	9	20
1:A:25:GLU:CG	1:A:40:LEU:HD22	0.80	2.06	6	13
1:A:79:VAL:HG11	1:A:100:LEU:HD21	0.79	1.55	8	8

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	95/130 (73%)	78±3 (82±3%)	16±3 (17±3%)	2±1 (2±1%)	17 61
All	All	1900/2600 (73%)	1550 (82%)	320 (17%)	30 (2%)	17 61

5 of 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	91	ASN	15
1	A	85	THR	3
1	A	17	ASN	3
1	A	84	GLY	3
1	A	9	VAL	2

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	84/112 (75%)	62±2 (74±2%)	22±2 (26±2%)	3 24
All	All	1680/2240 (75%)	1241 (74%)	439 (26%)	3 24

5 of 47 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	101	VAL	20
1	A	100	LEU	20
1	A	49	LEU	20
1	A	94	GLN	20
1	A	27	VAL	20

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 90% for the well-defined parts and 81% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 11255

Chemical shift list name: *assigned_ chem_ shift_ list_ 1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1428
Number of shifts mapped to atoms	1428
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	112	-0.45 ± 0.12	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	109	0.03 ± 0.12	None needed (< 0.5 ppm)
$^{13}C'$	111	-0.28 ± 0.08	None needed (< 0.5 ppm)
^{15}N	105	-0.39 ± 0.41	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 90%, i.e. 1056 atoms were assigned a chemical shift out of a possible 1172. 0 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	461/469 (98%)	185/187 (99%)	187/190 (98%)	89/92 (97%)
Sidechain	540/639 (85%)	330/372 (89%)	200/232 (86%)	10/35 (29%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	55/64 (86%)	29/35 (83%)	26/26 (100%)	0/3 (0%)
Overall	1056/1172 (90%)	544/594 (92%)	413/448 (92%)	99/130 (76%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: