



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 10:49 PM GMT

PDB ID : 1VCH
Title : Crystal Structure of a Phosphoribosyltransferase-related protein from *Thermus thermophilus*
Authors : Rehse, P.H.; Tahirov, T.H.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2004-03-08
Resolution : 1.94 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2910 (1.96-1.92)
Clashscore	102246	3095 (1.96-1.92)
Ramachandran outliers	100387	3062 (1.96-1.92)
Sidechain outliers	100360	3062 (1.96-1.92)
RSRZ outliers	91569	2915 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	175	<div> <div>5%</div> <div>71%</div> <div>24%</div> <div>• •</div> </div>
1	B	175	<div> <div>%</div> <div>80%</div> <div>17%</div> <div>•</div> </div>
1	C	175	<div> <div>3%</div> <div>78%</div> <div>18%</div> <div>• •</div> </div>
1	D	175	<div> <div>3%</div> <div>80%</div> <div>18%</div> <div>• •</div> </div>
1	E	175	<div> <div>45%</div> <div>36%</div> <div>46%</div> <div>5%</div> <div>13%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CL	B	1002	-	-	-	X
2	CL	C	1003	-	-	-	X
2	CL	D	1004	-	-	-	X
2	CL	E	1005	-	-	-	X
4	ACY	A	2002	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6899 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoribosyltransferase-related protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	169	Total	C	N	O	S	0	0	1
			1317	844	234	235	4			
1	B	170	Total	C	N	O	S	0	0	0
			1327	850	236	236	5			
1	C	170	Total	C	N	O	S	0	0	1
			1322	847	235	235	5			
1	D	173	Total	C	N	O	S	0	0	0
			1349	865	239	240	5			
1	E	152	Total	C	N	O	S	0	0	0
			1189	762	216	207	4			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Cl	0	0
			2	2		
2	D	1	Total	Cl	0	0
			1	1		
2	C	1	Total	Cl	0	0
			1	1		
2	E	1	Total	Cl	0	0
			1	1		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Ca	0	0
			1	1		

- Molecule 4 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	69	Total	O	0	0
			69	69		
5	B	95	Total	O	0	0
			95	95		
5	C	81	Total	O	0	0
			81	81		
5	D	97	Total	O	0	0
			97	97		
5	E	31	Total	O	0	0
			31	31		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phosphoribosyltransferase-related protein




- Molecule 1: Phosphoribosyltransferase-related protein



- Molecule 1: Phosphoribosyltransferase-related protein



- Molecule 1: Phosphoribosyltransferase-related protein



Protein	Residue	Score	Category
MET	L70	7.72	High
	A71	7.72	High
	E72	7.72	High
	T135	7.72	High
	M136	7.72	High
	R137	7.72	High
	A138	7.72	High
	M139	7.72	High
	L140	7.72	High
	M142	7.72	High
	L144	7.72	High
	L145	7.72	High
	H149	7.72	High
	V150	7.72	High
	L153	7.72	High
	L154	7.72	High
MET	L155	7.72	High
	V156	7.72	High
	F157	7.72	High
	R158	7.72	High
	P159	7.72	High
	G160	7.72	High
	T161	7.72	High
	P162	7.72	High
	G163	7.72	High
	L164	7.72	High
	A165	7.72	High
	D167	7.72	High
	T168	7.72	High
	V169	7.72	High
	E171	7.72	High
	MET	L172	7.72
L175		7.72	High
L176		7.72	High
L177		7.72	High
L178		7.72	High
L179		7.72	High
L180		7.72	High
L181		7.72	High
L182		7.72	High
L183		7.72	High
L184		7.72	High
L185		7.72	High
L186		7.72	High
L187		7.72	High
L188		7.72	High
L189		7.72	High
MET	L190	7.72	High
	L191	7.72	High
	L192	7.72	High
	L193	7.72	High
	L194	7.72	High
	L195	7.72	High
	L196	7.72	High
	L197	7.72	High
	L198	7.72	High
	L199	7.72	High
	L200	7.72	High
	L201	7.72	High
	L202	7.72	High
	L203	7.72	High
	L204	7.72	High
	L205	7.72	High
MET	L206	7.72	High
	L207	7.72	High
	L208	7.72	High
	L209	7.72	High
	L210	7.72	High
	L211	7.72	High
	L212	7.72	High
	L213	7.72	High
	L214	7.72	High
	L215	7.72	High
	L216	7.72	High
	L217	7.72	High
	L218	7.72	High
	L219	7.72	High
	L220	7.72	High
	MET	L221	7.72
L222		7.72	High
L223		7.72	High
L224		7.72	High
L225		7.72	High
L226		7.72	High
L227		7.72	High
L228		7.72	High
L229		7.72	High
L230		7.72	High
L231		7.72	High
L232		7.72	High
L233		7.72	High
L234		7.72	High
L235		7.72	High
MET		L236	7.72
	L237	7.72	High
	L238	7.72	High
	L239	7.72	High
	L240	7.72	High
	L241	7.72	High
	L242	7.72	High
	L243	7.72	High
	L244	7.72	High
	L245	7.72	High
	L246	7.72	High
	L247	7.72	High
	L248	7.72	High
	L249	7.72	High
	L250	7.72	High
	MET	L251	7.72
L252		7.72	High
L253		7.72	High
L254		7.72	High
L255		7.72	High
L256		7.72	High
L257		7.72	High
L258		7.72	High
L259		7.72	High
L260		7.72	High
L261		7.72	High
L262		7.72	High
L263		7.72	High
L264		7.72	High
L265		7.72	High
MET		L266	7.72
	L267	7.72	High
	L268	7.72	High
	L269	7.72	High
	L270	7.72	High
	L271	7.72	High
	L272	7.72	High
	L273	7.72	High
	L274	7.72	High
	L275	7.72	High
	L276	7.72	High
	L277	7.72	High
	L278	7.72	High
	L279	7.72	High
	L280	7.72	High
	MET	L281	7.72
L282		7.72	High
L283		7.72	High

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	167.42Å 61.41Å 102.39Å 90.00° 93.97° 90.00°	Depositor
Resolution (Å)	44.98 – 1.94 44.98 – 1.90	Depositor EDS
% Data completeness (in resolution range)	95.8 (44.98-1.94) 94.2 (44.98-1.90)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 1.89Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.232 , 0.259 0.230 , 0.258	Depositor DCC
R_{free} test set	3854 reflections (5.50%)	DCC
Wilson B-factor (Å ²)	30.8	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 80673 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6899	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ACY, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/1343	0.62	0/1829
1	B	0.34	0/1353	0.63	0/1842
1	C	0.32	0/1348	0.62	0/1834
1	D	0.30	0/1375	0.63	0/1873
1	E	0.38	0/1208	0.78	2/1634 (0.1%)
All	All	0.33	0/6627	0.66	2/9012 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	52	GLU	N-CA-C	-7.53	90.68	111.00
1	E	84	ARG	NE-CZ-NH2	6.68	123.64	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1317	0	1372	43	0
1	B	1327	0	1386	20	0
1	C	1322	0	1378	31	0
1	D	1349	0	1413	26	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1189	0	1224	112	0
2	B	2	0	0	1	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
3	C	1	0	0	0	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0
4	C	4	0	3	0	0
4	D	4	0	3	0	0
5	A	69	0	0	2	0
5	B	95	0	0	2	0
5	C	81	0	0	1	0
5	D	97	0	0	4	0
5	E	31	0	0	2	0
All	All	6899	0	6785	223	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 223 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:57:PHE:HE2	1:E:139:MET:HB3	1.14	1.09
1:E:95:GLN:HB3	1:E:108:LEU:HD11	1.06	1.06
1:E:108:LEU:HD13	1:E:142:MET:SD	1.95	1.06
1:E:158:ARG:HB3	1:E:168:THR:HG21	1.46	0.97
1:C:51:LYS:H	1:C:51:LYS:HE3	1.31	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	165/175 (94%)	162 (98%)	3 (2%)	0	100	100
1	B	166/175 (95%)	161 (97%)	4 (2%)	1 (1%)	30	16
1	C	166/175 (95%)	164 (99%)	1 (1%)	1 (1%)	30	16
1	D	169/175 (97%)	165 (98%)	3 (2%)	1 (1%)	30	16
1	E	136/175 (78%)	120 (88%)	12 (9%)	4 (3%)	6	0
All	All	802/875 (92%)	772 (96%)	23 (3%)	7 (1%)	21	9

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	51	LYS
1	E	47	PRO
1	C	91	ASP
1	B	91	ASP
1	E	91	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/147 (97%)	139 (98%)	3 (2%)	61	52
1	B	143/147 (97%)	141 (99%)	2 (1%)	74	68
1	C	142/147 (97%)	141 (99%)	1 (1%)	88	87
1	D	146/147 (99%)	145 (99%)	1 (1%)	88	87
1	E	126/147 (86%)	121 (96%)	5 (4%)	38	22
All	All	699/735 (95%)	687 (98%)	12 (2%)	68	61

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	51	LYS
1	D	57	PHE
1	E	158	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	122	ARG
1	E	128	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	159	GLN
1	D	98	GLN
1	D	159	GLN
1	C	120	ASN
1	D	120	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACY	A	2002	-	1,3,3	2.23	1 (100%)	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ACY	B	2001	-	1,3,3	2.25	1 (100%)	0,3,3	0.00	-
4	ACY	C	2003	-	1,3,3	2.03	1 (100%)	0,3,3	0.00	-
4	ACY	D	2004	-	1,3,3	2.20	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ACY	A	2002	-	-	0/0/0/0	0/0/0/0
4	ACY	B	2001	-	-	0/0/0/0	0/0/0/0
4	ACY	C	2003	-	-	0/0/0/0	0/0/0/0
4	ACY	D	2004	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	2003	ACY	CH3-C	2.03	1.51	1.48
4	D	2004	ACY	CH3-C	2.20	1.51	1.48
4	A	2002	ACY	CH3-C	2.23	1.51	1.48
4	B	2001	ACY	CH3-C	2.25	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	169/175 (96%)	0.36	8 (4%) 35 44	18, 39, 73, 85	0
1	B	170/175 (97%)	0.12	2 (1%) 81 86	20, 37, 57, 77	0
1	C	170/175 (97%)	0.21	6 (3%) 48 57	21, 40, 70, 96	0
1	D	173/175 (98%)	0.37	6 (3%) 48 57	22, 36, 63, 95	0
1	E	152/175 (86%)	2.39	79 (51%) 0 0	44, 74, 91, 99	0
All	All	834/875 (95%)	0.65	101 (12%) 6 9	18, 41, 83, 99	0

The worst 5 of 101 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	123	VAL	10.2
1	E	124	VAL	9.3
1	E	108	LEU	8.8
1	E	66	LEU	7.0
1	A	99	THR	6.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CL	D	1004	1/1	0.98	0.26	6.55	50,50,50,50	0
2	CL	C	1003	1/1	0.97	0.33	5.61	59,59,59,59	0
2	CL	B	1002	1/1	0.93	0.17	2.86	56,56,56,56	0
2	CL	E	1005	1/1	0.78	0.26	2.43	84,84,84,84	0
4	ACY	A	2002	4/4	0.89	0.18	2.08	59,60,60,66	0
4	ACY	C	2003	4/4	0.96	0.14	0.44	64,65,69,70	0
4	ACY	D	2004	4/4	0.94	0.09	-0.13	35,40,41,44	0
4	ACY	B	2001	4/4	0.95	0.10	-0.34	37,40,40,45	0
3	CA	C	1006	1/1	0.98	0.06	-	36,36,36,36	0
2	CL	B	1001	1/1	0.91	0.21	-	68,68,68,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.