



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:59 AM GMT

PDB ID : 2VJO
Title : FORMYL-COA TRANSFERASE MUTANT VARIANT Q17A WITH
ASPARTYL-COA THIOESTER INTERMEDIATES AND OXALATE
Authors : Berthold, C.L.; Toyota, C.G.; Richards, N.G.J.; Lindqvist, Y.
Deposited on : 2007-12-11
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	428	
1	B	428	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	OXL	A	1430	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7210 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FORMYL-COENZYME A TRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	427	Total	C	N	O	S	0	6	0
			3338	2113	572	630	23			
1	B	427	Total	C	N	O	S	0	1	0
			3313	2097	568	625	23			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	17	ALA	GLN	ENGINEERED MUTATION	UNP O06644
B	17	ALA	GLN	ENGINEERED MUTATION	UNP O06644
A	186	ILE	MET	CONFLICT SEE REMARK 9	UNP O06644
B	186	ILE	MET	CONFLICT SEE REMARK 9	UNP O06644

- Molecule 2 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	
			48	21	7	16	3	1	
2	B	1	Total	C	N	O	P	S	
			48	21	7	16	3	1	

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is OXALATE ION (three-letter code: OXL) (formula: C₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O		
			6	2	4	0	0

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	249	Total	O	0	0
			249	249		
6	B	191	Total	O	0	0
			191	191		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FORMYL-COENZYME A TRANSFERASE



• Molecule 1: FORMYL-COENZYME A TRANSFERASE



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	153.59Å 153.59Å 98.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 39.07 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (30.00-2.20) 99.4 (39.07-2.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.206 , 0.248 0.202 , 0.242	Depositor DCC
R_{free} test set	2894 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	30.5	Xtriage
Anisotropy	0.070	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.2	EDS
Estimated twinning fraction	0.037 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 57456 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7210	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: COA, OXL, EPE, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/3431	0.56	0/4640
1	B	0.43	0/3390	0.55	0/4584
All	All	0.43	0/6821	0.56	0/9224

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3338	0	3283	19	0
1	B	3313	0	3255	29	0
2	A	48	0	31	0	0
2	B	48	0	31	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	6	0	0	0	0
5	A	15	0	17	0	0
6	A	249	0	0	1	0
6	B	191	0	0	3	0
All	All	7210	0	6617	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:418:LYS:H	1:A:418:LYS:HD2	1.55	0.72
1:A:186:ILE:HG23	1:B:4:PRO:HA	1.77	0.66
1:B:11:LEU:HD11	1:B:36:ILE:HD11	1.79	0.65
1:A:167:LEU:HD21	1:B:167:LEU:HD21	1.78	0.64
1:A:9[B]:ASN:HD22	1:A:90:ALA:HA	1.63	0.64
1:B:11:LEU:CD1	1:B:36:ILE:HD11	2.27	0.64
1:A:135[B]:HIS:CD2	1:A:135[B]:HIS:H	2.15	0.64
1:A:364:LYS:HE3	1:B:130:GLY:O	2.02	0.59
1:B:291:GLN:HB3	1:B:343:TYR:HB3	1.88	0.55
1:A:124:VAL:HG23	1:A:176:LEU:HD22	1.90	0.54
1:A:24:GLN:HE21	1:A:65:CYS:H	1.55	0.52
1:A:11:LEU:HD11	1:A:36:ILE:HD11	1.92	0.52
1:B:228:PRO:HA	1:B:231:GLN:HG2	1.91	0.51
1:A:331:LYS:HB3	1:A:335:GLU:HG3	1.93	0.50
1:B:228:PRO:HB3	1:B:234:PHE:HE2	1.77	0.50
1:A:29:LEU:HD22	1:A:391:PHE:HZ	1.76	0.49
1:B:323:PHE:O	1:B:326:THR:OG1	2.28	0.49
1:A:237:ASP:OD2	1:A:241:ASN:HB2	2.13	0.49
1:B:78:GLU:HB2	1:B:427:VAL:HG21	1.93	0.48
1:B:96:ASN:ND2	6:B:2049:HOH:O	2.46	0.47
1:B:292:ILE:O	1:B:296:ILE:HG12	2.15	0.47
1:B:95:GLU:HG2	1:B:102:LEU:HD22	1.98	0.46
1:A:409:LEU:HD11	1:A:428:VAL:HG21	1.99	0.45
1:B:234:PHE:CG	1:B:318:MET:HG3	2.52	0.45
1:B:121:LEU:HD23	1:B:195:LYS:HG3	1.98	0.45
1:B:324:ILE:O	1:B:324:ILE:HG13	2.17	0.44
1:B:75:LYS:HG2	2:B:1169:COA:H2A	1.99	0.44
1:B:343:TYR:HB2	1:B:345:ILE:HG13	2.00	0.43
1:A:384:ALA:HA	1:A:385:PRO:HD3	1.83	0.43
1:B:195:LYS:HE2	6:B:2058:HOH:O	2.18	0.43
1:B:122:ALA:HA	1:B:196:VAL:O	2.19	0.43
1:B:4:PRO:HD3	1:B:391:PHE:CE2	2.54	0.42
1:B:320:ILE:O	1:B:324:ILE:HG23	2.20	0.42
1:A:188:HIS:HE1	1:B:185:GLU:OE1	2.03	0.42
1:B:342:GLN:HG2	1:B:342:GLN:H	1.72	0.42
1:A:356:LEU:HD21	1:B:146:SER:O	2.19	0.42
1:B:234:PHE:CD2	1:B:318:MET:HG3	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:322:SER:HB2	6:B:2138:HOH:O	2.20	0.41
1:A:233[B]:ASN:ND2	6:A:2109:HOH:O	2.53	0.41
1:B:74:MET:HE2	2:B:1169:COA:C6A	2.51	0.41
1:A:16:VAL:O	1:A:20:PRO:HG2	2.21	0.40
1:B:135:HIS:H	1:B:135:HIS:CD2	2.38	0.40
1:A:231:GLN:OE1	1:A:317:LEU:HB2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	431/428 (101%)	420 (97%)	10 (2%)	1 (0%)	52	59
1	B	426/428 (100%)	412 (97%)	12 (3%)	2 (0%)	34	35
All	All	857/856 (100%)	832 (97%)	22 (3%)	3 (0%)	39	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	VAL
1	B	16	VAL
1	B	240	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	355/350 (101%)	338 (95%)	17 (5%)	31	37
1	B	350/350 (100%)	341 (97%)	9 (3%)	54	66
All	All	705/700 (101%)	679 (96%)	26 (4%)	41	50

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ARG
1	A	59	TYR
1	A	89	LYS
1	A	102	LEU
1	A	123	SER
1	A	186	ILE
1	A	208	VAL
1	A	247	ASN
1	A	314	VAL
1	A	317	LEU
1	A	386	PHE
1	A	391	PHE
1	A	392	GLN
1	A	410	LYS
1	A	418	LYS
1	A	419	ILE
1	A	428	VAL
1	B	15	HIS
1	B	59	TYR
1	B	81	GLU
1	B	138	VAL
1	B	169	ASP
1	B	309	THR
1	B	317	LEU
1	B	324	ILE
1	B	386	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	GLN
1	A	188	HIS
1	A	392	GLN
1	B	135	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	COA	A	1169	1	40,50,50	1.02	1 (2%)	50,75,75	1.78	5 (10%)
4	OXL	A	1430	-	0,5,5	0.00	-	0,6,6	0.00	-
5	EPE	A	1431	-	14,15,15	0.38	0	18,20,20	2.04	5 (27%)
2	COA	B	1169	1	40,50,50	1.08	2 (5%)	50,75,75	1.76	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	A	1169	1	-	0/44/64/64	0/3/3/3
4	OXL	A	1430	-	-	0/0/4/4	0/0/0/0
5	EPE	A	1431	-	-	0/9/19/19	0/1/1/1
2	COA	B	1169	1	-	0/44/64/64	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1169	COA	O4B-C1B	2.15	1.43	1.41
2	A	1169	COA	O9P-C9P	4.89	1.33	1.23
2	B	1169	COA	O9P-C9P	5.16	1.33	1.23

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1169	COA	N3A-C2A-N1A	-10.09	121.17	128.89
2	A	1169	COA	N3A-C2A-N1A	-10.04	121.20	128.89
2	B	1169	COA	P2A-O3A-P1A	-2.91	124.56	132.73
2	B	1169	COA	C4A-C5A-N7A	-2.52	107.16	109.48
2	A	1169	COA	C4A-C5A-N7A	-2.41	107.26	109.48
2	A	1169	COA	CDP-CBP-CCP	-2.26	105.57	108.50
2	A	1169	COA	P2A-O3A-P1A	-2.08	126.90	132.73
2	A	1169	COA	O4B-C1B-N9A	2.52	113.38	108.10
5	A	1431	EPE	O1S-S-C10	2.61	109.13	106.91
5	A	1431	EPE	C7-N4-C5	2.68	118.13	111.27
5	A	1431	EPE	C7-N4-C3	3.13	119.30	111.27
5	A	1431	EPE	O2S-S-C10	4.34	110.61	106.91
5	A	1431	EPE	C5-N4-C3	4.74	119.17	108.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1169	COA	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	427/428 (99%)	0.02	16 (3%) 45 44	15, 27, 51, 67	0
1	B	427/428 (99%)	0.55	60 (14%) 4 3	15, 29, 81, 94	0
All	All	854/856 (99%)	0.29	76 (8%) 12 11	15, 28, 65, 94	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	305	PRO	8.4
1	B	303	ASP	7.4
1	B	320	ILE	7.1
1	B	296	ILE	6.7
1	B	341	ALA	6.4
1	B	302	LYS	6.2
1	B	234	PHE	5.8
1	B	292	ILE	5.6
1	B	297	ASP	5.5
1	B	323	PHE	5.4
1	B	294	ASP	5.3
1	B	290	PRO	5.0
1	B	301	TRP	5.0
1	B	309	THR	4.8
1	B	299	PRO	4.7
1	B	298	LYS	4.6
1	B	308	ASN	4.6
1	B	344	GLY	4.5
1	B	314	VAL	4.3
1	B	339	TRP	4.3
1	B	310	PHE	4.1
1	B	420	LYS	4.1
1	B	307	TYR	4.1
1	B	285	ALA	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	288	MET	4.0
1	B	295	MET	3.9
1	B	233	ASN	3.9
1	B	306	ALA	3.8
1	B	291	GLN	3.7
1	B	232	PRO	3.6
1	A	2	THR	3.5
1	B	289	TRP	3.5
1	B	315	ASP	3.5
1	B	286	ALA	3.5
1	B	346	PRO	3.4
1	B	424	ALA	3.3
1	A	421	GLU	3.2
1	B	342	GLN	3.1
1	A	419	ILE	3.1
1	B	287	ASN	3.1
1	B	343	TYR	3.0
1	A	427	VAL	3.0
1	A	415	ASP	2.9
1	B	304	ASP	2.9
1	A	413	GLY	2.7
1	B	317	LEU	2.7
1	B	313	ARG	2.7
1	B	330	ASP	2.6
1	B	418	LYS	2.6
1	B	239	ASP	2.5
1	A	207	LEU	2.5
1	B	327	LYS	2.5
1	B	413	GLY	2.5
1	B	415	ASP	2.5
1	A	199	ALA	2.4
1	B	426	GLN	2.4
1	B	311	GLU	2.4
1	A	114	GLU	2.4
1	A	77	PRO	2.4
1	A	420	LYS	2.4
1	B	241	ASN	2.3
1	A	425	LYS	2.3
1	B	284	ILE	2.2
1	B	282	PHE	2.2
1	A	167	LEU	2.2
1	B	414	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	326	THR	2.2
1	A	424	ALA	2.2
1	B	166	ALA	2.2
1	A	414	LEU	2.1
1	B	421	GLU	2.1
1	A	17	ALA	2.1
1	B	238	ARG	2.0
1	B	17	ALA	2.0
1	B	425	LYS	2.0
1	B	81	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	OXL	A	1430	6/6	0.78	0.24	3.84	55,57,57,57	0
5	EPE	A	1431	15/15	0.92	0.16	0.69	32,33,34,34	15
2	COA	B	1169	48/48	0.91	0.14	-0.25	30,45,48,48	0
2	COA	A	1169	48/48	0.93	0.12	-0.90	25,42,46,46	0
3	CL	B	1429	1/1	0.99	0.17	-0.92	21,21,21,21	0
3	CL	A	1429	1/1	0.99	0.12	-2.10	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.