



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:59 AM GMT

PDB ID : 2VJV
Title : CRYSTAL STRUCTURE OF THE IS608 TRANSPOSASE IN COMPLEX
WITH LEFT END 26-MER DNA HAIRPIN AND A 6-MER DNA REPRESENTING
THE LEFT END CLEAVAGE SITE
Authors : Barabas, O.; Ronning, D.R.; Guynet, C.; Hickman, A.B.; Ton-Hoang, B.;
Chandler, M.; Dyda, F.
Deposited on : 2007-12-13
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	<div> <div>3%</div> <div>67% 10% 21%</div> </div>
1	B	159	<div> <div>5%</div> <div>70% 7% 21%</div> </div>
2	C	26	<div> <div>62% 38%</div> </div>
2	D	26	<div> <div>42% 54%</div> </div>
3	E	6	<div> <div>17%</div> <div>67% 33%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	6	<div><div></div><div></div><div></div></div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3622 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSPOSASE ORFA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	125	Total	C	N	O	S	0	0	0
			1023	659	181	177	6			
1	B	125	Total	C	N	O	S	0	0	0
			1023	659	181	177	6			

- Molecule 2 is a DNA chain called 5'-D(*DA*DA*DA*DG*DC*DC*DC*DC*DT*DA*DG*DC*DTP*DT *DT*DT*DA*DG*DC*DT*DA*DT*DG*DG*DG*DGP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	26	Total	C	N	O	P	0	0	0
			531	254	97	155	25			
2	D	26	Total	C	N	O	P	0	0	0
			531	254	97	155	25			

- Molecule 3 is a DNA chain called 5'-D(*DT*DA*DT*DT*DA*DCP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	6	Total	C	N	O	P	0	0	0
			118	59	19	35	5			
3	F	6	Total	C	N	O	P	0	0	0
			118	59	19	35	5			



- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		
4	E	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	63	Total 63	O 63	0	0
5	B	70	Total 70	O 70	0	0
5	C	53	Total 53	O 53	0	0
5	D	59	Total 59	O 59	0	0
5	E	20	Total 20	O 20	0	0
5	F	10	Total 10	O 10	0	0

- Molecule 1: TRANSPOSASE ORFA





- Molecule 3: 5'-D(*DT*DA*DT*DT*DA*DCP)-3'

Chain F: 17% 83% 17%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.29 Å 72.41 Å 110.17 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.61 – 1.90 24.61 – 1.90	Depositor EDS
% Data completeness (in resolution range)	87.8 (24.61-1.90) 87.9 (24.61-1.90)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.49 (at 1.90 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.194 , 0.227 0.193 , 0.218	Depositor DCC
R_{free} test set	1131 reflections (3.04%)	DCC
Wilson B-factor (Å ²)	24.9	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	2 of 37219 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3622	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/1044	0.78	1/1407 (0.1%)
1	B	0.46	0/1044	0.73	1/1407 (0.1%)
2	C	0.52	0/595	0.75	0/917
2	D	0.57	0/595	0.78	0/917
3	E	0.43	0/131	0.65	0/200
3	F	0.47	0/131	0.63	0/200
All	All	0.49	0/3540	0.75	2/5048 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
2	D	0	2
All	All	0	3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	20	HIS	N-CA-C	-5.27	96.76	111.00
1	B	61	ASP	N-CA-C	-5.25	96.83	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	18	DA	Sidechain
2	D	18	DA	Sidechain
2	D	37	DT	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1023	0	1064	13	0
1	B	1023	0	1064	7	0
2	C	531	0	295	14	0
2	D	531	0	295	16	0
3	E	118	0	71	1	0
3	F	118	0	71	1	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
5	A	63	0	0	1	0
5	B	70	0	0	0	0
5	C	53	0	0	0	0
5	D	59	0	0	0	0
5	E	20	0	0	0	0
5	F	10	0	0	0	0
All	All	3622	0	2860	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:40:DG:H2'	2:D:41:DG:H5''	1.20	1.10
2:D:40:DG:C2'	2:D:41:DG:H5''	1.82	1.08
2:D:25:DA:H2''	2:D:26:DG:H5'	1.44	0.97
2:C:31:DT:H5'	2:C:31:DT:H6	1.33	0.90
1:A:62:LYS:H	1:A:62:LYS:HD2	1.46	0.81
2:D:40:DG:H2'	2:D:41:DG:C5'	2.09	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:31:DT:C6	2:C:31:DT:H5'	2.18	0.78
2:C:31:DT:C5'	2:C:31:DT:H6	1.96	0.77
2:D:25:DA:H2''	2:D:26:DG:C5'	2.20	0.72
2:C:26:DG:H2''	2:C:27:DC:H5''	1.75	0.68
2:D:41:DG:H8	2:D:41:DG:H5'	1.63	0.63
1:A:129:GLU:O	1:A:130:ASN:ND2	2.32	0.63
1:B:29:ARG:HG3	1:B:103:LEU:HD21	1.82	0.61
1:A:90:ARG:HG3	2:C:30:DT:H1'	1.85	0.58
1:A:29:ARG:HG3	1:A:103:LEU:HD21	1.84	0.58
2:C:29:DT:H3'	2:C:30:DT:H5'	1.85	0.58
2:C:31:DT:C6	2:C:31:DT:C5'	2.81	0.57
2:D:27:DC:H2'	2:D:28:DT:H71	1.87	0.56
2:D:30:DT:H4'	2:D:32:DA:O4'	2.06	0.54
2:D:40:DG:C3'	2:D:41:DG:H5''	2.36	0.54
2:D:23:DC:H2''	2:D:24:DT:H5'	1.89	0.54
2:C:30:DT:H2''	2:C:31:DT:O5'	2.08	0.53
1:A:82:LYS:HB3	2:C:35:DT:H4'	1.93	0.50
2:D:27:DC:H2''	2:D:28:DT:O5'	2.13	0.48
1:A:121:LEU:O	1:A:125:LYS:HG3	2.13	0.47
1:B:93:ARG:HB3	1:B:100:LYS:HE3	1.97	0.46
1:A:40:LEU:O	1:A:44:ILE:HG12	2.15	0.46
1:B:121:LEU:HA	1:B:124:VAL:HG22	1.96	0.46
1:A:127:TYR:O	3:F:-1:DC:H2'	2.16	0.46
1:A:30:LYS:HB3	1:A:62:LYS:HB3	1.99	0.44
1:B:61:ASP:HB2	1:B:64:HIS:CE1	2.51	0.44
1:A:93:ARG:HB3	1:A:100:LYS:HE2	1.98	0.44
2:D:36:DA:H3'	2:D:37:DT:H5''	1.99	0.43
2:D:41:DG:C8	2:D:41:DG:H5'	2.49	0.43
2:C:32:DA:N3	2:C:32:DA:H5''	2.34	0.43
1:B:128:ILE:HG13	1:B:129:GLU:HG3	2.00	0.43
2:C:27:DC:C6	2:C:28:DT:H72	2.54	0.42
1:B:74:SER:HB2	2:C:37:DT:H4'	2.00	0.42
2:C:26:DG:H2''	2:C:27:DC:C5'	2.48	0.42
2:C:32:DA:N3	2:C:32:DA:H2'	2.35	0.42
1:A:33:VAL:CG1	1:A:62:LYS:HG3	2.50	0.41
1:B:82:LYS:HE3	2:D:34:DC:O2	2.20	0.41
1:A:74:SER:HB2	2:D:37:DT:H4'	2.01	0.41
3:E:-5:DA:H1'	3:E:-4:DT:H71	2.02	0.40
2:D:37:DT:H3'	2:D:38:DG:C5'	2.52	0.40
1:A:34:GLY:HA3	5:A:2027:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	123/159 (77%)	121 (98%)	2 (2%)	0	100	100
1	B	123/159 (77%)	121 (98%)	2 (2%)	0	100	100
All	All	246/318 (77%)	242 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/144 (80%)	112 (97%)	3 (3%)	54	45
1	B	115/144 (80%)	113 (98%)	2 (2%)	68	64
All	All	230/288 (80%)	225 (98%)	5 (2%)	60	53

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	ARG
1	A	100	LYS
1	A	130	ASN
1	B	68	LEU
1	B	82	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	130	ASN
1	B	64	HIS
1	B	97	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	125/159 (78%)	0.20	5 (4%) 42 46	16, 24, 52, 63	0
1	B	125/159 (78%)	0.31	8 (6%) 23 25	13, 23, 55, 71	0
2	C	26/26 (100%)	-0.17	0 100 100	18, 28, 44, 64	0
2	D	26/26 (100%)	-0.24	0 100 100	18, 27, 44, 56	0
3	E	6/6 (100%)	0.07	1 (16%) 2 2	24, 26, 33, 80	0
3	F	6/6 (100%)	-0.04	1 (16%) 2 2	23, 24, 32, 81	0
All	All	314/382 (82%)	0.17	15 (4%) 34 37	13, 25, 55, 81	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	122	ASN	6.1
1	A	6	LEU	4.5
1	A	122	ASN	4.0
3	E	-6	DT	3.2
1	A	98	HIS	3.1
1	B	6	LEU	3.0
3	F	-6	DT	2.7
1	B	67	ILE	2.5
1	B	126	GLN	2.4
1	A	130	ASN	2.3
1	B	21	ILE	2.3
1	A	102	LYS	2.2
1	B	101	THR	2.2
1	B	23	TRP	2.1
1	B	123	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	MG	C	1042	1/1	0.99	0.06	-1.51	20,20,20,20	0
4	MG	D	1042	1/1	0.99	0.03	-2.48	16,16,16,16	0
4	MG	E	1042	1/1	0.97	0.15	-	41,41,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.