



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:58 PM GMT

PDB ID : 3VP1  
Title : Crystal structure of human glutaminase in complex with L-glutamate and BPTES  
Authors : Thangavelu, K.; Sivaraman, J.  
Deposited on : 2012-02-23  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtrriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:


Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	603	-	X	-	-





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	10	5	1	4	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
4	A	1	5	4	1	0	0
4	A	1	5	4	1	0	0


- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	103	Total 103	O 103	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutaminase kidney isoform, mitochondrial



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	139.64Å 139.64Å 156.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30 40.03 – 2.30	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-2.30) 94.4 (40.03-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.13 (at 2.29Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.205 , 0.232 0.211 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	39.7	Xtrriage
Anisotropy	0.309	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 40.1	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Outliers	0 of 34688 reflections	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2552	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 04A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	3/2470 (0.1%)	0.65	1/3340 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	321	LEU	CG-CD2	-8.52	1.20	1.51
1	A	321	LEU	C-O	-5.61	1.12	1.23
1	A	320	LYS	CA-CB	-5.07	1.42	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	315	GLY	N-CA-C	-6.11	97.82	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2411	0	2344	56	1
2	A	18	0	10	2	2
3	A	10	0	5	0	0
4	A	10	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	103	0	0	1	0
All	All	2552	0	2359	57	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:319:ASN:O	1:A:320:LYS:HB2	1.39	1.14
1:A:320:LYS:HD3	1:A:320:LYS:N	1.90	0.85
1:A:507:LYS:HD2	1:A:507:LYS:H	1.47	0.80
1:A:316:LEU:HB2	1:A:318:PHE:CE1	2.22	0.75
1:A:268:VAL:HG13	1:A:436:ASN:OD1	1.90	0.71
1:A:318:PHE:HB3	1:A:321:LEU:HD12	1.72	0.71
1:A:313:PRO:HA	1:A:463:CYS:SG	2.32	0.70
1:A:315:GLY:O	1:A:317:ARG:N	2.26	0.69
1:A:316:LEU:HB2	1:A:318:PHE:CD1	2.31	0.66
1:A:228:THR:HG23	1:A:273:HIS:CE1	2.30	0.66
1:A:318:PHE:CB	1:A:321:LEU:HD12	2.26	0.65
1:A:507:LYS:CD	1:A:507:LYS:H	2.08	0.65
1:A:498:MET:HE3	1:A:517:PHE:CE2	2.33	0.64
1:A:507:LYS:HD2	1:A:507:LYS:N	2.15	0.62
1:A:319:ASN:O	1:A:320:LYS:CB	2.20	0.60
1:A:234:LEU:HD22	1:A:520:ASP:HB3	1.82	0.60
1:A:435:ALA:HB2	1:A:491:VAL:HG13	1.82	0.59
1:A:317:ARG:O	1:A:317:ARG:HG2	2.03	0.58
1:A:260:LEU:HD13	1:A:501:TRP:CH2	2.39	0.57
1:A:320:LYS:HD3	1:A:320:LYS:H	1.66	0.57
1:A:432:ALA:HB1	1:A:441:PRO:HG2	1.89	0.55
1:A:323:LEU:HD21	1:A:395:LEU:CD2	2.37	0.54
1:A:316:LEU:HB2	1:A:318:PHE:HE1	1.69	0.53
1:A:318:PHE:CB	1:A:330:HIS:HE2	2.21	0.53
1:A:227:PHE:CZ	1:A:231:ILE:HD11	2.44	0.53
1:A:265:VAL:HG22	1:A:498:MET:HG2	1.91	0.52
1:A:251:PRO:O	1:A:255:LYS:HG3	2.09	0.51
2:A:601:04A:NAV	2:A:601:04A:OAB	2.42	0.51
1:A:260:LEU:HD13	1:A:501:TRP:HH2	1.74	0.51
1:A:285:GLN:O	1:A:288:VAL:HG12	2.13	0.48
1:A:498:MET:HE3	1:A:517:PHE:HE2	1.74	0.48
1:A:382:ARG:HD2	5:A:710:HOH:O	2.12	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:318:PHE:HA	1:A:321:LEU:HD12	1.96	0.47
1:A:498:MET:HB3	1:A:498:MET:HE2	1.68	0.47
1:A:316:LEU:HD12	1:A:316:LEU:N	2.29	0.47
1:A:274:SER:HB3	1:A:278:THR:HG21	1.96	0.47
1:A:265:VAL:HG22	1:A:498:MET:CG	2.46	0.46
1:A:320:LYS:HD2	1:A:320:LYS:HA	1.30	0.46
1:A:314:SER:HB3	1:A:316:LEU:H	1.80	0.46
1:A:318:PHE:CA	1:A:321:LEU:HD12	2.46	0.45
1:A:267:THR:OG1	1:A:269:ASP:HB2	2.16	0.45
1:A:223:ASP:OD2	1:A:226:SER:HB2	2.17	0.45
1:A:479:PRO:HD2	1:A:491:VAL:O	2.17	0.44
1:A:427:ALA:HB3	1:A:499:MET:HG2	1.99	0.44
1:A:356:ASP:O	1:A:360:GLN:HG3	2.17	0.44
1:A:313:PRO:CB	1:A:462:SER:HB2	2.48	0.43
1:A:317:ARG:CG	1:A:317:ARG:O	2.65	0.43
1:A:278:THR:O	1:A:425:GLU:HG3	2.19	0.43
1:A:323:LEU:HD21	1:A:395:LEU:HD23	2.00	0.42
1:A:453:VAL:O	1:A:457:LEU:HG	2.19	0.42
1:A:314:SER:HB2	1:A:316:LEU:HG	2.01	0.42
1:A:317:ARG:HD3	2:A:601:04A:HAH	2.01	0.42
1:A:318:PHE:HB2	1:A:330:HIS:NE2	2.36	0.41
1:A:318:PHE:HA	1:A:321:LEU:CD1	2.50	0.40
1:A:412:ASP:O	1:A:416:GLN:HG3	2.20	0.40
1:A:318:PHE:CE2	1:A:325:GLU:HG3	2.56	0.40
1:A:492:VAL:HG12	1:A:495:VAL:HB	2.03	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:601:04A:CAN	2:A:601:04A:SAY[8_555]	1.21	0.99
1:A:317:ARG:NH1	1:A:317:ARG:NH1[15_555]	1.90	0.30
2:A:601:04A:CAN	2:A:601:04A:CAN[8_555]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	311/315 (99%)	295 (95%)	14 (4%)	2 (1%)	30 36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	316	LEU
1	A	317	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	266/268 (99%)	259 (97%)	7 (3%)	54 71

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	241	GLN
1	A	312	GLU
1	A	316	LEU
1	A	317	ARG
1	A	320	LYS
1	A	361	PHE
1	A	507	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	360	GLN
1	A	529	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	04A	A	601	-	17,19,38	4.97	11 (64%)	13,24,49	6.51	6 (46%)
3	GLU	A	602	-	3,9,9	0.21	0	2,11,11	0.13	0
4	SO4	A	603	-	4,4,4	5.23	4 (100%)	6,6,6	1.15	1 (16%)
4	SO4	A	604	-	4,4,4	0.29	0	6,6,6	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	04A	A	601	-	-	0/8/11/24	0/1/2/4
3	GLU	A	602	-	-	0/3/9/9	0/0/0/0
4	SO4	A	603	-	-	0/0/0/0	0/0/0/0
4	SO4	A	604	-	-	0/0/0/0	0/0/0/0

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	04A	CAN-SAY	-17.13	1.21	1.80
2	A	601	04A	NAT-NAV	-4.53	1.28	1.37
2	A	601	04A	CBG-SBA	-3.23	1.63	1.73
2	A	601	04A	CAR-CBE	2.02	1.55	1.51
2	A	601	04A	CBG-NAT	2.16	1.36	1.33
2	A	601	04A	CAL-CBE	2.18	1.43	1.38
2	A	601	04A	CAD-CAG	2.63	1.44	1.38
2	A	601	04A	CBI-NAX	2.74	1.41	1.36
2	A	601	04A	CAH-CAL	3.95	1.47	1.38
2	A	601	04A	CAK-CBE	4.58	1.48	1.38
2	A	601	04A	CAD-CAH	4.68	1.50	1.38
4	A	603	SO4	O2-S	4.93	1.64	1.47
4	A	603	SO4	O1-S	5.11	1.64	1.47
4	A	603	SO4	O3-S	5.25	1.66	1.47
4	A	603	SO4	O4-S	5.63	1.67	1.47

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	04A	OAB-CBC-CAR	-3.88	111.03	121.88
2	A	601	04A	CAR-CBE-CAK	-3.42	115.73	120.86
2	A	601	04A	CAR-CBE-CAL	2.63	124.81	120.86
4	A	603	SO4	O2-S-O1	2.72	118.12	109.50
2	A	601	04A	CBE-CAR-CBC	2.91	122.33	112.84
2	A	601	04A	CAR-CBC-NAX	4.37	127.96	114.81
2	A	601	04A	CAP-CAN-SAY	21.98	141.52	113.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	04A	2	2

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	313/315 (99%)	0.26	19 (6%) 25 33	25, 38, 60, 85	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	315	GLY	9.4
1	A	318	PHE	7.8
1	A	314	SER	7.3
1	A	317	ARG	5.1
1	A	316	LEU	4.8
1	A	222	PRO	4.1
1	A	223	ASP	3.4
1	A	348	GLY	2.8
1	A	319	ASN	2.8
1	A	269	ASP	2.7
1	A	320	LYS	2.6
1	A	403	GLU	2.5
1	A	225	MET	2.5
1	A	313	PRO	2.4
1	A	219	SER	2.3
1	A	230	HIS	2.1
1	A	220	MET	2.1
1	A	524	LEU	2.0
1	A	227	PHE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GLU	A	602	10/10	0.85	0.19	1.48	43,54,56,57	0
2	04A	A	601	18/35	0.93	0.16	-0.51	31,38,45,46	1
4	SO4	A	604	5/5	0.81	0.33	-	24,25,28,28	5
4	SO4	A	603	5/5	0.94	0.20	-	19,24,25,26	5

### 6.5 Other polymers [i](#)

There are no such residues in this entry.