



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:01 PM GMT

PDB ID : 3VR2
Title : Crystal structure of nucleotide-free A3B3 complex from *Enterococcus hirae* V-ATPase [eA3B3]
Authors : Arai, S.; Saijo, S.; Suzuki, K.; Mizutani, K.; Kakinuma, Y.; Ishizuka-Katsura, Y.; Ohsawa, N.; Terada, T.; Shirouzu, M.; Yokoyama, S.; Iwata, S.; Yamato, I.; Murata, T.
Deposited on : 2012-04-03
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	600	<div> <div>0%</div> <div>78% 18% ..</div> </div>
1	B	600	<div> <div>4%</div> <div>78% 19% ..</div> </div>
1	C	600	<div> <div>8%</div> <div>69% 25% ..</div> </div>
2	D	465	<div> <div>3%</div> <div>67% 24% 5% .</div> </div>
2	E	465	<div> <div>6%</div> <div>74% 21% ..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	465	<div><div></div><div>13%</div><div>72%</div><div>23%</div><div>• •</div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 24053 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called V-type sodium ATPase catalytic subunit A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	586	Total	C	N	O	S	Se	0	0	0
			4559	2864	766	904	3	22			
1	B	586	Total	C	N	O	S	Se	0	0	0
			4555	2862	766	902	3	22			
1	C	583	Total	C	N	O	S	Se	0	0	0
			4501	2829	757	890	3	22			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	GLY	-	EXPRESSION TAG	UNP Q08636
A	-5	SER	-	EXPRESSION TAG	UNP Q08636
A	-4	SER	-	EXPRESSION TAG	UNP Q08636
A	-3	GLY	-	EXPRESSION TAG	UNP Q08636
A	-2	SER	-	EXPRESSION TAG	UNP Q08636
A	-1	SER	-	EXPRESSION TAG	UNP Q08636
A	0	GLY	-	EXPRESSION TAG	UNP Q08636
B	-6	GLY	-	EXPRESSION TAG	UNP Q08636
B	-5	SER	-	EXPRESSION TAG	UNP Q08636
B	-4	SER	-	EXPRESSION TAG	UNP Q08636
B	-3	GLY	-	EXPRESSION TAG	UNP Q08636
B	-2	SER	-	EXPRESSION TAG	UNP Q08636
B	-1	SER	-	EXPRESSION TAG	UNP Q08636
B	0	GLY	-	EXPRESSION TAG	UNP Q08636
C	-6	GLY	-	EXPRESSION TAG	UNP Q08636
C	-5	SER	-	EXPRESSION TAG	UNP Q08636
C	-4	SER	-	EXPRESSION TAG	UNP Q08636
C	-3	GLY	-	EXPRESSION TAG	UNP Q08636
C	-2	SER	-	EXPRESSION TAG	UNP Q08636
C	-1	SER	-	EXPRESSION TAG	UNP Q08636
C	0	GLY	-	EXPRESSION TAG	UNP Q08636

- Molecule 2 is a protein called V-type sodium ATPase subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	447	Total 3447	C 2179	N 596	O 658	Se 14	0	0	0
2	E	449	Total 3448	C 2180	N 596	O 658	Se 14	0	0	0
2	F	451	Total 3454	C 2177	N 598	O 665	Se 14	0	0	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-6	GLY	-	EXPRESSION TAG	UNP Q08637
D	-5	SER	-	EXPRESSION TAG	UNP Q08637
D	-4	SER	-	EXPRESSION TAG	UNP Q08637
D	-3	GLY	-	EXPRESSION TAG	UNP Q08637
D	-2	SER	-	EXPRESSION TAG	UNP Q08637
D	-1	SER	-	EXPRESSION TAG	UNP Q08637
D	0	GLY	-	EXPRESSION TAG	UNP Q08637
E	-6	GLY	-	EXPRESSION TAG	UNP Q08637
E	-5	SER	-	EXPRESSION TAG	UNP Q08637
E	-4	SER	-	EXPRESSION TAG	UNP Q08637
E	-3	GLY	-	EXPRESSION TAG	UNP Q08637
E	-2	SER	-	EXPRESSION TAG	UNP Q08637
E	-1	SER	-	EXPRESSION TAG	UNP Q08637
E	0	GLY	-	EXPRESSION TAG	UNP Q08637
F	-6	GLY	-	EXPRESSION TAG	UNP Q08637
F	-5	SER	-	EXPRESSION TAG	UNP Q08637
F	-4	SER	-	EXPRESSION TAG	UNP Q08637
F	-3	GLY	-	EXPRESSION TAG	UNP Q08637
F	-2	SER	-	EXPRESSION TAG	UNP Q08637
F	-1	SER	-	EXPRESSION TAG	UNP Q08637
F	0	GLY	-	EXPRESSION TAG	UNP Q08637

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	39	Total 39	O 39	0	0
3	B	8	Total 8	O 8	0	0
3	C	14	Total 14	O 14	0	0
3	D	8	Total 8	O 8	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	12	Total	O	0	0
			12	12		
3	F	8	Total	O	0	0
			8	8		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: V-type sodium ATPase catalytic subunit A




- Molecule 1: V-type sodium ATPase catalytic subunit A




- Molecule 1: V-type sodium ATPase catalytic subunit A





• Molecule 2: V-type sodium ATPase subunit B



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	121.80Å 121.50Å 128.26Å 90.00° 90.06° 90.00°	Depositor
Resolution (Å)	44.18 – 2.80 44.18 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (44.18-2.80) 99.3 (44.18-2.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.11 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.198 , 0.243 0.198 , 0.242	Depositor DCC
R_{free} test set	4573 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	64.4	Xtriage
Anisotropy	0.204	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 51.5	EDS
Estimated twinning fraction	0.009 for k,h,-l 0.011 for -k,-h,-l 0.021 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 91450 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	24053	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/4613	0.74	1/6206 (0.0%)
1	B	0.49	0/4609	0.61	0/6201
1	C	0.49	0/4555	0.59	0/6131
2	D	0.57	0/3493	0.68	0/4698
2	E	0.58	0/3494	0.68	0/4700
2	F	0.48	0/3501	0.63	0/4706
All	All	0.55	0/24265	0.66	1/32642 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	209	MSE	CG-SE-CE	-5.76	86.22	98.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	124	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4559	0	4521	98	0
1	B	4555	0	4517	75	0
1	C	4501	0	4435	145	0
2	D	3447	0	3416	112	0
2	E	3448	0	3400	89	0
2	F	3454	0	3394	87	0
3	A	39	0	0	3	0
3	B	8	0	0	0	0
3	C	14	0	0	0	0
3	D	8	0	0	4	0
3	E	12	0	0	1	0
3	F	8	0	0	1	0
All	All	24053	0	23683	578	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 578 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:102:GLU:CB	2:E:103:ILE:HA	1.63	1.24
1:A:139:VAL:CG2	1:A:187:MSE:HE1	1.72	1.19
1:A:320:MSE:HE2	1:A:322:TYR:CE2	1.84	1.12
2:E:79:LEU:HD13	2:E:227:MSE:CE	1.80	1.12
2:D:361:THR:HG21	2:D:365:HIS:CD2	1.85	1.11

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	584/600 (97%)	556 (95%)	27 (5%)	1 (0%)	52	84
1	B	584/600 (97%)	540 (92%)	41 (7%)	3 (0%)	34	69
1	C	581/600 (97%)	537 (92%)	42 (7%)	2 (0%)	46	79
2	D	443/465 (95%)	407 (92%)	31 (7%)	5 (1%)	17	50
2	E	445/465 (96%)	411 (92%)	33 (7%)	1 (0%)	52	84
2	F	449/465 (97%)	407 (91%)	39 (9%)	3 (1%)	26	62
All	All	3086/3195 (97%)	2858 (93%)	213 (7%)	15 (0%)	34	69

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	174	SER
2	D	392	SER
1	C	280	ASN
2	D	172	ASP
1	B	391	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	501/487 (103%)	476 (95%)	25 (5%)	30	64
1	B	500/487 (103%)	476 (95%)	24 (5%)	31	66
1	C	489/487 (100%)	451 (92%)	38 (8%)	16	41
2	D	358/372 (96%)	322 (90%)	36 (10%)	9	27
2	E	355/372 (95%)	329 (93%)	26 (7%)	17	44
2	F	356/372 (96%)	336 (94%)	20 (6%)	26	59
All	All	2559/2577 (99%)	2390 (93%)	169 (7%)	21	51

5 of 169 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	476	LEU
2	D	73	LEU
2	F	77	LEU
1	C	494	LYS
1	C	555	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 56 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	485	ASN
2	D	112	ASN
2	F	112	ASN
1	C	502	GLN
1	C	520	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	563/600 (93%)	-0.18	3 (0%) 91 88	24, 41, 83, 117	0
1	B	563/600 (93%)	0.14	26 (4%) 36 25	38, 71, 129, 155	0
1	C	561/600 (93%)	0.35	47 (8%) 14 6	38, 76, 175, 198	0
2	D	433/465 (93%)	-0.09	13 (3%) 54 41	34, 60, 133, 196	0
2	E	435/465 (93%)	0.15	30 (6%) 20 11	32, 66, 167, 221	0
2	F	437/465 (93%)	0.58	61 (13%) 4 2	42, 91, 191, 212	0
All	All	2992/3195 (93%)	0.15	180 (6%) 25 15	24, 66, 169, 221	0

The worst 5 of 180 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	581	ILE	9.4
2	F	388	VAL	8.0
1	C	565	GLU	7.1
2	F	447	ASP	7.0
2	D	397	ILE	6.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.