



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:00 PM GMT

PDB ID : 3VRA
Title : Mitochondrial rhodoquinol-fumarate reductase from the parasitic nematode *Ascaris suum* with the specific inhibitor Atpenin A5
Authors : Shimizu, H.; Shiba, T.; Inaoka, D.K.; Osanai, A.; Kita, K.; Sakamoto, K.; Harada, S.
Deposited on : 2012-04-07
Resolution : 3.44 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1007 (3.54-3.34)
Clashscore	102246	1044 (3.52-3.36)
Ramachandran outliers	100387	1013 (3.52-3.36)
Sidechain outliers	100360	1014 (3.52-3.36)
RSRZ outliers	91569	1012 (3.54-3.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	645	<div> <div>6%</div> <div> <div>45%</div> <div>41%</div> <div>9%</div> <div>5%</div> </div> </div>
1	E	645	<div> <div>2%</div> <div> <div>42%</div> <div>46%</div> <div>7%</div> <div>5%</div> </div> </div>
2	B	282	<div> <div>%</div> <div> <div>36%</div> <div>42%</div> <div>11%</div> <div>12%</div> </div> </div>
2	F	282	<div> <div>2%</div> <div> <div>40%</div> <div>34%</div> <div>14%</div> <div>12%</div> </div> </div>
3	C	188	<div> <div>4%</div> <div> <div>32%</div> <div>39%</div> <div>10%</div> <div>19%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	188	
4	D	156	
4	H	156	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	AT5	C	202	-	-	X	X
11	AT5	G	202	-	-	X	X
12	EPH	D	201	-	-	-	X
12	EPH	H	201	-	-	-	X
5	MLI	A	701	-	-	X	-
5	MLI	E	701	-	-	X	-
8	SF4	B	302	-	-	X	-
9	F3S	F	303	-	-	X	-

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 18262 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Flavoprotein subunit of complex II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	613	Total	C	N	O	S	0	0	0
			4758	2983	851	896	28			
1	E	613	Total	C	N	O	S	0	0	0
			4758	2983	851	896	28			

- Molecule 2 is a protein called Iron-sulfur subunit of succinate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	249	Total	C	N	O	S	0	0	0
			1973	1254	337	359	23			
2	F	249	Total	C	N	O	S	0	0	0
			1973	1254	337	359	23			


- Molecule 3 is a protein called Cytochrome b-large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	153	Total	C	N	O	S	0	0	0
			1217	813	204	194	6			
3	G	153	Total	C	N	O	S	0	0	0
			1217	813	204	194	6			

- Molecule 4 is a protein called Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	129	Total	C	N	O	S	0	0	0
			994	658	165	166	5			
4	H	129	Total	C	N	O	S	0	0	0
			994	658	165	166	5			

- Molecule 5 is MALONATE ION (three-letter code: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	3	4		
5	E	1	Total	C	O	0	0
			7	3	4		

- Molecule 6 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 7 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	Fe	S	0	0
			4	2	2		
7	F	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 8 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			8	4	4		
8	F	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 9 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe_3S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	Fe	S	0	0
			7	3	4		
9	F	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 10 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
10	G	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 11 is 3-[(2S,4S,5R)-5,6-DICHLORO-2,4-DIMETHYL-1-OXOHXYL]-4-H YDROXY-5,6-DIMETHOXY-2(1H)-PYRIDINONE (three-letter code: AT5) (formula: $C_{15}H_{21}Cl_2NO_5$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	C	1	Total	C	Cl	N	O	0	0
			23	15	2	1	5		
11	G	1	Total	C	Cl	N	O	0	0
			23	15	2	1	5		

- Molecule 12 is L-ALPHA-PHOSPHATIDYL-BETA-OLEOYL-GAMMA-PALMITOYL-PHOSPHATIDYLETHANOLAMINE (three-letter code: EPH) (formula: $C_{39}H_{68}NO_8P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	D	1	Total	C	N	O	P	0	0
			44	34	1	8	1		
12	H	1	Total	C	N	O	P	0	0
			44	34	1	8	1		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Flavoprotein subunit of complex II




• Molecule 1: Flavoprotein subunit of complex II






• Molecule 3: Cytochrome b-large subunit




• Molecule 3: Cytochrome b-large subunit



• Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial



• Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	122.82Å 132.25Å 220.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.72 – 3.44 49.72 – 3.44	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.72-3.44) 89.5 (49.72-3.44)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.26 (at 3.40Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.205 , 0.281 0.203 , 0.278	Depositor DCC
R_{free} test set	2217 reflections (5.39%)	DCC
Wilson B-factor (Å ²)	103.9	Xtriage
Anisotropy	0.079	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 43.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	0 of 43349 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	18262	wwPDB-VP
Average B, all atoms (Å ²)	107.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, MLI, F3S, FES, EPH, HEM, AT5, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	0/4859	0.77	2/6564 (0.0%)
1	E	0.68	0/4859	0.81	0/6564
2	B	0.72	1/2016 (0.0%)	0.83	0/2723
2	F	0.77	1/2016 (0.0%)	0.89	1/2723 (0.0%)
3	C	0.72	0/1255	0.83	2/1709 (0.1%)
3	G	0.71	0/1255	0.84	2/1709 (0.1%)
4	D	0.74	0/1026	0.83	1/1402 (0.1%)
4	H	0.75	0/1026	0.80	0/1402
All	All	0.70	2/18312 (0.0%)	0.81	8/24796 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	187	CYS	CB-SG	-5.08	1.73	1.81
2	B	97	CYS	CB-SG	-5.04	1.73	1.81

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	439	LEU	CA-CB-CG	6.64	130.58	115.30
4	D	52	LEU	CA-CB-CG	6.52	130.30	115.30
2	F	179	LEU	CA-CB-CG	6.16	129.46	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	184	LEU	CA-CB-CG	5.49	127.92	115.30
3	G	165	LEU	CA-CB-CG	5.24	127.35	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	425	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4758	0	4692	260	0
1	E	4758	0	4692	284	0
2	B	1973	0	1994	135	0
2	F	1973	0	1994	133	0
3	C	1217	0	1265	86	0
3	G	1217	0	1265	67	0
4	D	994	0	977	48	0
4	H	994	0	977	59	0
5	A	7	0	2	3	0
5	E	7	0	2	4	0
6	A	53	0	30	12	0
6	E	53	0	31	16	0
7	B	4	0	0	0	0
7	F	4	0	0	1	0
8	B	8	0	0	5	0
8	F	8	0	0	0	0
9	B	7	0	0	0	0
9	F	7	0	0	2	0
10	C	43	0	30	14	0
10	G	43	0	30	13	0
11	C	23	0	20	12	0
11	G	23	0	21	11	0
12	D	44	0	53	4	0
12	H	44	0	53	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	18262	0	18128	1009	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 1009 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:79:HIS:CD2	6:E:702:FAD:HM82	1.71	1.26
1:E:79:HIS:NE2	6:E:702:FAD:HM82	0.92	1.24
3:C:114:TRP:HB2	3:C:175:ARG:HH11	1.09	1.14
2:F:115:GLN:HA	2:F:115:GLN:HE21	1.14	1.08
3:C:107:ILE:HD11	4:D:156:LEU:HD22	1.35	1.07

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	611/645 (95%)	493 (81%)	92 (15%)	26 (4%)	3	30
1	E	611/645 (95%)	499 (82%)	82 (13%)	30 (5%)	3	26
2	B	247/282 (88%)	204 (83%)	34 (14%)	9 (4%)	4	36
2	F	247/282 (88%)	192 (78%)	42 (17%)	13 (5%)	2	23
3	C	151/188 (80%)	110 (73%)	34 (22%)	7 (5%)	3	28
3	G	151/188 (80%)	118 (78%)	25 (17%)	8 (5%)	2	23
4	D	127/156 (81%)	96 (76%)	27 (21%)	4 (3%)	5	40
4	H	127/156 (81%)	87 (68%)	32 (25%)	8 (6%)	2	19
All	All	2272/2542 (89%)	1799 (79%)	368 (16%)	105 (5%)	3	28

5 of 105 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	489	THR
1	A	614	ARG
2	B	89	CYS
3	C	97	PRO
4	D	49	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	498/526 (95%)	415 (83%)	83 (17%)	3	15
1	E	498/526 (95%)	424 (85%)	74 (15%)	4	20
2	B	219/242 (90%)	173 (79%)	46 (21%)	1	6
2	F	219/242 (90%)	172 (78%)	47 (22%)	1	6
3	C	127/158 (80%)	94 (74%)	33 (26%)	0	3
3	G	127/158 (80%)	106 (84%)	21 (16%)	3	15
4	D	96/119 (81%)	75 (78%)	21 (22%)	1	5
4	H	96/119 (81%)	75 (78%)	21 (22%)	1	5
All	All	1880/2090 (90%)	1534 (82%)	346 (18%)	2	10

5 of 346 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	165	LEU
1	E	162	LYS
3	G	149	ILE
3	C	177	GLU
4	D	121	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 66 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	140	ASN
1	E	159	ASN
2	F	165	GLN
1	E	88	ASN
1	E	120	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	MLI	A	701	-	0,6,6	0.00	-	0,7,7	0.00	-
6	FAD	A	702	1	48,58,58	1.34	7 (14%)	54,89,89	2.44	16 (29%)
7	FES	B	301	2	0,4,4	0.00	-	0,4,4	0.00	-
8	SF4	B	302	2	0,12,12	0.00	-	0,24,24	0.00	-
9	F3S	B	303	2	0,9,9	0.00	-	0,15,15	0.00	-
10	HEM	C	201	3,4	30,50,50	2.11	9 (30%)	24,82,82	2.51	10 (41%)
11	AT5	C	202	-	19,23,23	1.57	3 (15%)	16,32,32	2.44	5 (31%)
12	EPH	D	201	-	42,43,48	1.98	8 (19%)	43,48,53	2.65	9 (20%)
5	MLI	E	701	-	0,6,6	0.00	-	0,7,7	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	FAD	E	702	1	48,58,58	1.37	6 (12%)	54,89,89	2.60	13 (24%)
7	FES	F	301	2	0,4,4	0.00	-	0,4,4	0.00	-
8	SF4	F	302	2	0,12,12	0.00	-	0,24,24	0.00	-
9	F3S	F	303	2	0,9,9	0.00	-	0,15,15	0.00	-
10	HEM	G	201	3,4	30,50,50	1.97	7 (23%)	24,82,82	2.29	8 (33%)
11	AT5	G	202	-	19,23,23	1.47	3 (15%)	16,32,32	2.60	5 (31%)
12	EPH	H	201	-	42,43,48	2.12	12 (28%)	43,48,53	2.95	10 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MLI	A	701	-	-	0/0/4/4	0/0/0/0
6	FAD	A	702	1	-	0/30/50/50	0/6/6/6
7	FES	B	301	2	-	0/0/4/4	0/1/1/1
8	SF4	B	302	2	-	0/0/48/48	0/6/5/5
9	F3S	B	303	2	-	0/0/24/24	0/0/3/3
10	HEM	C	201	3,4	-	0/10/54/54	0/0/8/8
11	AT5	C	202	-	-	2/22/22/22	0/1/1/1
12	EPH	D	201	-	-	0/47/47/52	0/0/0/0
5	MLI	E	701	-	-	0/0/4/4	0/0/0/0
6	FAD	E	702	1	-	0/30/50/50	0/6/6/6
7	FES	F	301	2	-	0/0/4/4	0/1/1/1
8	SF4	F	302	2	-	0/0/48/48	0/6/5/5
9	F3S	F	303	2	-	0/0/24/24	0/0/3/3
10	HEM	G	201	3,4	-	0/10/54/54	0/0/8/8
11	AT5	G	202	-	-	0/22/22/22	0/1/1/1
12	EPH	H	201	-	-	0/47/47/52	0/0/0/0

The worst 5 of 55 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	C	201	HEM	C3B-C4B	-6.31	1.46	1.51
10	G	201	HEM	C3B-C4B	-6.12	1.46	1.51
10	C	201	HEM	C2C-C1C	-4.40	1.44	1.52
10	G	201	HEM	C2C-C1C	-4.10	1.44	1.52
10	C	201	HEM	C3D-C4D	-3.81	1.46	1.51

The worst 5 of 76 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	E	702	FAD	N3A-C2A-N1A	-12.58	119.26	128.89
6	A	702	FAD	N3A-C2A-N1A	-11.06	120.43	128.89
11	G	202	AT5	CM3-O3-C3	-7.78	109.58	117.31
11	C	202	AT5	C6-C5-N4	-5.99	119.65	124.19
11	C	202	AT5	C11-C12-C13	-4.77	101.92	110.28

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	C	202	AT5	CM3-O3-C3-C2
11	C	202	AT5	CM3-O3-C3-N4

There are no ring outliers.

13 monomers are involved in 97 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	701	MLI	3	0
6	A	702	FAD	12	0
8	B	302	SF4	5	0
10	C	201	HEM	14	0
11	C	202	AT5	12	0
12	D	201	EPH	4	0
5	E	701	MLI	4	0
6	E	702	FAD	16	0
7	F	301	FES	1	0
9	F	303	F3S	2	0
10	G	201	HEM	13	0
11	G	202	AT5	11	0
12	H	201	EPH	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	613/645 (95%)	0.29	41 (6%) 21 19	84, 111, 132, 145	0
1	E	613/645 (95%)	-0.06	11 (1%) 71 65	78, 105, 126, 141	0
2	B	249/282 (88%)	-0.15	3 (1%) 81 75	78, 99, 125, 146	0
2	F	249/282 (88%)	0.04	6 (2%) 62 56	71, 92, 114, 130	0
3	C	153/188 (81%)	-0.05	8 (5%) 31 27	89, 107, 148, 165	0
3	G	153/188 (81%)	0.16	12 (7%) 16 15	88, 101, 172, 199	0
4	D	129/156 (82%)	-0.19	4 (3%) 52 47	92, 107, 131, 142	0
4	H	129/156 (82%)	-0.03	5 (3%) 43 38	90, 109, 145, 151	0
All	All	2288/2542 (90%)	0.05	90 (3%) 43 38	71, 106, 134, 199	0

The worst 5 of 90 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	185	PRO	12.1
3	G	184	LEU	9.0
4	H	49	PHE	7.0
3	G	186	THR	6.4
3	G	181	LYS	5.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
12	EPH	D	201	44/49	0.88	0.45	8.17	114,133,158,158	0
12	EPH	H	201	44/49	0.85	0.72	7.53	101,138,160,161	0
11	AT5	C	202	23/23	0.92	0.25	2.84	110,115,144,151	0
11	AT5	G	202	23/23	0.90	0.30	2.22	115,121,134,148	0
10	HEM	G	201	43/43	0.94	0.28	1.55	111,117,126,129	0
5	MLI	A	701	7/7	0.92	0.37	1.00	99,101,101,102	0
10	HEM	C	201	43/43	0.97	0.21	0.69	105,108,121,127	0
6	FAD	A	702	53/53	0.95	0.36	0.13	69,90,95,96	0
5	MLI	E	701	7/7	0.97	0.24	0.07	95,96,96,96	0
8	SF4	B	302	8/8	1.00	0.20	-0.10	67,72,75,75	0
8	SF4	F	302	8/8	1.00	0.21	-0.30	57,65,68,68	0
6	FAD	E	702	53/53	0.98	0.26	-0.34	63,76,88,90	0
9	F3S	F	303	7/7	1.00	0.17	-0.38	79,82,83,83	0
7	FES	B	301	4/4	1.00	0.25	-0.80	78,78,80,82	0
7	FES	F	301	4/4	1.00	0.24	-0.90	82,85,88,89	0
9	F3S	B	303	7/7	0.99	0.15	-0.97	89,90,92,93	0

6.5 Other polymers

There are no such residues in this entry.